Poster (M16)

Preinvasive Disease of Cervix, Vulva, and Vagina

<https://doi.org/10.3802/jgo.2021.32.S1.M16>

Efficacy of lidocaine spray for pain reduction during colposcopy-directed cervical biopsy: a randomized controlled trial

Charuwan Saeteng,^{*} Ornwitsanate Mongkolmafai, Kittipat Charoenkwan, Prapaporn Suprasert, Jatupol Srisomboon

Chiang Mai University, Chiang Mai, Thailand (nokctantipa@gmail.com)

Objective: To evaluate the efficacy of lidocaine spray in reducing pain during colposcopy-directed cervical biopsy (CDB).

Methods: From December 2017 through February 2019, 312 women undergoing CDB were enrolled. The participants were randomized to 3 groups: group 1 (lidocaine spray), lidocaine spray was applied thoroughly to the cervix; group 2 (placebo), normal saline was applied thoroughly to the cervix; group 3 (control), no anesthetic agent was applied to the cervix. Each woman completed a 10-cm visual analog scale for subjective pain experience at three-time points: baseline, immediately after biopsy, and 10 minutes after the procedure. The primary outcome of this study was the biopsy pain score.

Results: Of the 312 women, 104 were randomly assigned to each study group. The clinic-pathological and procedure-related characteristics of the participants in all groups were similar. The baseline, the biopsy, and the post-procedure pain scores were comparable among the three groups. We found a significant difference in the pain scores changes from baseline to biopsy and from baseline to postprocedure among the three groups. However, when compared to the lidocaine spray group directly to the control group, there was no statistically significant difference in the pain scores changes from baseline to biopsy ($p=0.06$) and from baseline to postprocedure ($p=0.29$). There was no complication with the intervention observed.

Conclusion: The application of lidocaine spray to the cervix has a small benefit in reducing pain associated with CDB. However, the intervention is safe and may be considered in nulliparous and/or overly anxious women undergoing the procedure.

Poster (M17)

Preinvasive Disease of Cervix, Vulva, and Vagina

<https://doi.org/10.3802/jgo.2021.32.S1.M17>

Treatment outcomes of patients with cervical intraepithelial neoplasia or invasive carcinoma who underwent loop electrosurgical excision procedure (LEEP)

Jakkapan Khunnarong,^{1,*} Nitinan Bunyasontikul,² Siriwan Tangitgamol³

¹Faculty of Medicine Vajira Hospital, Navamindradhiraj University, Bangkok, Thailand (jak020@hotmail.com)

²Obstetrics and Gynecology Section, Somdejprapinklao Hospital, Bangkok, Thailand

³Women's Health Center, MedPark Hospital, Bangkok, Thailand

Objective: To study treatment outcomes of cervical intraepithelial neoplasia (CIN) or cancer patients who underwent loop electrosurgical excision procedure (LEEP) in terms of primary outcome and factors associated with persistent/recurrence.

Methods: Patients with CIN or cancer who underwent LEEP from January 2007 to December 2015 were reviewed. Data collected were age, parity, menopausal status, human immunodeficiency virus (HIV) infection, smoking, cervical cytology, histopathology from cervical biopsy and LEEP including margin status, final histopathology, and follow-up data.

Results: The mean age of 385 patients was 41.9 + 10.8 years (range, 18–79 years). Majority were multiparous (81.6%) and premenopausal (78.2%). There were 15.3% of patients had HIV infection. The most common cervical cytology was HSIL (44.1%) followed by ACS-US (21%). Minor complications of bleeding or infection from LEEP were encountered 7.3 %. Among 153 patients (39.7%) who had positive margin(s), 43 underwent second LEEP whereas 76 had hysterectomy. From all patients, 47 had failure aftertreatment (12.2%): being either persistence (30 patients; 7.8%) or recurrence (17 patients; 4.4%). Factors associated with persistence or recurrence by multivariate analysis were age ≥ 55 years old, HIV infection, final diagnosis of invasive cancer, and positive of endocervical margin or both ecto- and endo-cervical margins.

Conclusion: LEEP had low rate of persistence/recurrence. Age ≥ 55 years old, HIV infection, final diagnosis of cancer, and positive of endocervical or both endo- and ecto-surgical margin(s) were significantly associated with persistent or recurrent diseases.

Poster (M18)

Quality of Life, Nursing & Supportive Care

<https://doi.org/10.3802/jgo.2021.32.S1.M18>

Quality of life and psychological aspects among ovarian cancer survivors compare to normal female population at Bhumibol Adulyadej Hospital

Wanlaya Onwatanasrikul,^{1,*} Worrawan Sirichai,¹ Komsan Suwanalerk²

¹Bhumibol Adulyadej Hospital, Bangkok, Thailand (wanlaya.onw@gmail.com)

²Thammasat University Hospital, Pathum Thani, Thailand

Objective: This study intended to compare quality of life (QOL) and psychological aspects among ovarian cancer survivors to healthy female population who presented at Bhumibol Adulyadej Hospital, using standardized questionnaire.

Methods: A cross-sectional analytical study, conducted at Bhumibol Adulyadej Hospital, Thailand from August 2020 to May 2021. This study enrolled healthy Thai female who came to hospital for their health check-up at gynecology clinic as a control group and ovarian cancer survivors on their follow up at tumor clinic as a study group. Questionnaire by Functional Assessment of Cancer Therapy (FACT) was used to measure QOL. Anxiety and depression were assessed by Hospital Anxiety and Depression Scale.

Results: Results shown ovarian cancer survivors and healthy participants had an equal QOL in all four domains of FACT general (FACT-G). Total FACT-G score was similar between two groups (50.2±9.7 vs. 49.5±8.5). QOL of nonepithelial ovarian cancer survivors were better than control group (56.1±5.4 vs. 49.5±8.5) in clinical significance. Psychological aspects were also comparable between two groups.

Conclusion: Ovarian cancer survivors experienced deteriorating in physical function or emotional related to treatment. But after completed the course of treatment, QOL will improve over time along with mental status, approaching level that comparable to the healthy peers.

Poster (M19)

Quality of Life, Nursing & Supportive Care

<https://doi.org/10.3802/jgo.2021.32.S1.M19>

Sexual dysfunction in gynecologic cancer survivors

Apirak Nguanboonmak,^{*} Marut Yanaranop

Rajavithi Hospital, Bangkok, Thailand (oporakung@hotmail.com)

Objective: To evaluate the sexual dysfunction in gynecologic cancer survivors and to explore the risk factors associated between gynecologic cancer survivors and sexual dysfunction.

Methods: A cross-sectional study was conducted with Thai gynecologic cancer survivors in Rajavithi Hospital. To assess the sexual dysfunction, a Thai version of the Female Sexual Function Index (FSFI) questionnaire was utilized. The inclusion criteria were Thai women who were gynecologic cancer survivors scheduled at the Gynecologic Oncology Clinic, Rajavithi Hospital, and had active sexual intercourse (SI). Data were collected between September 1, 2020, to January 31, 2021.

Results: A total of 180 gynecologic cancer survivors who had active SI participated in this study. Those consisted of cervical

cancer (43.9%), endometrial cancer (28.3%), and ovarian/tubal/peritoneal cancer (25.6%) and gestational trophoblastic neoplasia (2.2%). Ninety-nine out of 180 women (55%) had sexual dysfunction (FSFI score <26.5). Cervical cancer was found in the majority with sexual dysfunction, but the mean FSFI score was the lowest in endometrial cancer. Among the six domains, desire had the lowest mean score and pain had the highest mean score. Furthermore, postmenopausal status was the significant risk factors that had an impact on sexual dysfunction.

Conclusion: The prevalence of sexual dysfunction in gynecologic cancer survivors was 55%, and cervical cancer was found in the majority. The factors that significantly affected sexual dysfunction were postmenopausal status.

Poster (M20)

Quality of Life, Nursing & Supportive Care

<https://doi.org/10.3802/jgo.2021.32.S1.M20>

Symptom experience of cancer-related cognitive impairment in women with ovarian cancer

Liying Wang,¹ Yi Zhang,¹ Mei Xue,¹ Zhenqi Lu,¹ Xiaojun Zhang,¹ Yan Ding^{2,*}

¹Fudan University Shanghai Cancer Center, Fudan University, Shanghai, China

²Department of Nursing, Obstetrics & Gynecology Hospital of Fudan University, Shanghai, China (ding_yan@fudan.edu.cn)

Objective: To explore Chinese ovarian cancer survivors' symptom experience of cancer-related cognitive impairment (CRCI).

Methods: Semi-structured interviews were conducted with 34 women aged 22–65 in Cancer Center of Fudan University, Shanghai, China. This study was guided by symptom management theory and the conceptual model of preliminary cognition model among cervical cancer survivors. Qualitative content analysis was used to refine themes.

Results: The average age of participants (n=34) was 51.38 years (range, 25–65 years) and median time since diagnosis was 1 year (range, 0.2–42.8 years). Three themes and 9 subthemes were identified from the research. For perception of cognitive changes: Become stupid, not as smart as before; not noticing/having the chance to notice any cognition change. Possible influence events in the evaluation of cognitive changes include that comfortable, monotonous, isolation and narrow-focused life cause the brain to “retire”; Surgical anesthesia and chemotherapy were blamed for the change; Fatigue, sleep disorders and anxiety were also considered to affect cognition. For the impact of CRCI, most participants expressed that it was not an issue for them, and they did not particularly care, only a few said that their self-confidence was affected a little.