Poster (E29)

Gynecologic Cancer Screening

<https://doi.org/10.3802/jgo.2021.32.S1.E29>

Profiling of menopausal symptoms and therapeutic effects of hormone replacement therapy (HRT) in endometrial cancer survivors

Megumi Yokota,* Wataru Yamagami, Shiyoko Kitazawa, Satoko Tanimoto, Takuma Yoshimura, Kensuke Sakai, Takayuki Chiyoda, Takashi Iwata, Koji Banno, Daisuke Aoki

Keio University School of Medicine, Tokyo, Japan
(megu_ayu2726@yahoo.co.jp)

Objective: This study aimed to investigate the profile of menopausal symptoms in endometrial cancer survivors and analyze the effects and risks of hormone replacement therapy (HRT).

Methods: This was a retrospective study in 33 endometrial cancer survivors who underwent hysterectomy and bilateral oophorectomy (BSO) at Keio University Hospital from January 2013 to June 2019 and received HRT. We administered HRT to patients with postsurgical stage I or II endometrial cancer who complained of menopausal symptoms. Some patients with severe menopausal symptoms diagnosed with stage III disease were also treated after obtaining informed consent after being informed of the risks. Patients filled out a questionnaire on menopausal symptoms before and after HRT. We analyzed patient characteristics, clinicopathological factors, adverse events, and oncologic outcomes, which were extracted from medical records.

Results: The median age at surgical menopause was 44 years, and the median body mass index was 21.2 kg/m². Using the The International Federation of Gynaecology and Obstetrics 2008 staging criteria, 30 patients were diagnosed with postsurgical stage I disease, and 3 patients were with diagnosed with stage III disease. The histological grade of endometrial cancer was grade 1 in 22 patients, grade 2 in 8 patients, and grade 3 in 2 patients. The median duration of HRT was 963 days (641–2,681 days). Our results revealed that 31/33 patients (96%) showed improvement of menopausal symptoms. During the entire observation period, there were no cases of endometrial cancer recurrence.

Conclusion: This study shows that HRT can be given for menopausal symptoms in endometrial cancer survivors in Japan.

Poster (E30)

Endometrial Hyperplasia, Endometrial Intra-epithelial Neoplasia, and Endometrial Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.E30>

Sentinel lymph node mapping in endometrial cancer

Atthapon Jaishuen,* Pisutt Srichaikul, Khemanat Khemworapong

Siriraj Hospital, Bangkok, Thailand (ajaishuen@hotmail.com)

A 57-year-old female presented with postmenopausal bleeding. Fractional curettage showed grade 1 endometrioid carcinoma. She underwent laparoscopic surgical staging with sentinel lymph node (SLN) biopsy using indocyanine green (ICG) and near infrared camera. ICG was injected at 2, 4, 8, and 10 o'clock of cervix, 1 mL superficially and 1 mL at 1 cm depth. Ten minutes after ICG injection, the laparoscopic surgery was begun. Theoretically, ICG will come out of the cervix through the lymph chain, cross the internal iliac artery and 90% drain at obturator or external iliac LN. Otherwise, the SLN will be found in internal iliac, presacral or para-aortic area. This patient has ICG SLN at both obturator area. The patient underwent total laparoscopic hysterectomy, bilateral pelvic lymphadenectomy and para-aortic LN dissection to the level of inferior mesenteric artery. The SLNs were processed with ultrastaging, which was positive micrometastasis on the right side. The full lymphadenectomy revealed negative of other 26 pelvic LNs on the right, 18 pelvic LNs of the left and 5 para-aortic LNs. The postoperative period was uneventful. The patient was discharge in 2 days. We conducted the prospective study to see the detection rate of SLN in endometrial cancer (20% conventional laparoscopy and 80% laparotomy). The detection rate of unilateral SLN was 99%. The detection rate of bilateral SLN was 90%. We performed 100% pelvic lymphadenectomy (mean LN of 17) and 80% para-aortic LN removal (mean LN of 5). The sensitivity of SLN compared to full lymphadenectomy was 86% with 98% negative predictive value and 98% accuracy. The false negative rate was still high at 14%. However, this study is not designed to achieve the false negative rate. We are recruiting more patients to get the real false negative rate. If it is acceptable, we will not do the full lymphadenectomy anymore.

Ovarian cancer

Poster (O01)

Epithelial Ovarian Cancer including Borderline Tumor

<https://doi.org/10.3802/jgo.2021.32.S1.O01>

A retrospective review of mucinous borderline ovarian tumor at Siriraj Hospital, Thailand: oncologic outcome including important risk factors

Vitcha Poonyakanok,* Pattama Chaopotong, Malee Warnnissorn

Siriraj Hospital, Mahidol University, Bangkok, Thailand
(vpoonyakanok@gmail.com)

Objective: A borderline ovarian tumor (BOT) was an important group of tumors of the ovary which needs to be distinguished from invasive ovarian cancer. The most common subtype of BOT in East Asia was the mucinous subtype (mBOT). mBOT was still controversial in terms of management due to different oncologic outcomes and difficulty in preoperative diagnosis. The objectives of the current study were to determine the prognosis and risk factors for recurrence, focusing only on mucinous subtype.

Methods: This retrospective study included patients with mucinous BOT, diagnosed between January 2004 and December 2019, who were operated at Siriraj Hospital. Patients with other subtypes, invasive carcinoma, and other primary carcinomas were excluded.

Results: Two hundred and thirty-five patients were diagnosed as mBOT. The median follow-up time was 54 months. There were 9 patients (3.8%) who had tumor recurrence. Risk factors of recurrence were a ruptured tumor, presence of ascites, residual tumor, and stage of the tumor. The recurrence rate between fertility-sparing surgery and radical surgery was not different. Detailed surgical staging, a pathological finding of intraepithelial carcinoma, and microinvasion were not associated with recurrence of the disease.

Conclusion: mBOT had an excellent prognosis. Nowadays, fertility-sparing surgery had more roles with no difference in the oncologic outcomes. Patients with risk factors had to be observed closely. Moreover, pre/perioperative diagnosis and accuracy of the final pathological results were also important.

Poster (O02)

Epithelial Ovarian Cancer including Borderline Tumor
<https://doi.org/10.3802/jgo.2021.32.S1.O02>

Bowel surgery by gynecologic oncologists during maximal cytoreductive surgery for advanced ovarian cancer

Chi-Son Chang, Chel Hun Choi, Tae-Joong Kim, Jeong-Won Lee, Byoung-Gie Kim, Yoo-Young Lee*

Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea (yooyoung.lee@samsung.com)

Objective: We report the oncological outcomes in patients with advanced ovarian cancer who had bowel surgery which was performed by gynecologic oncologist (GO) during maximal cytoreductive surgery and compared the outcomes with those of bowel surgery performed by general surgeons (GS).

Methods: Patients who were stage III–IV ovarian cancer and had bowel surgery during maximal cytoreductive surgery were eligible. Patients were divided into two groups according to

whether bowel resection was performed by GO (group 1) or GS (group 2). In both groups, GO were mainly involved in debulking procedures. Perioperative and survival outcomes were compared between 2 groups.

Results: A total of 268 patients were eligible. Fifty-five patients received bowel surgery by GO (group 1), and 213 patients by GS (group 2). The rate of R0 resection (no residual tumor) after maximal cytoreductive surgery did not differ between 2 groups (45.5% vs. 40.8%, $p=0.529$). Among extra-uterine procedures, the rate of diaphragm stripping was higher in group 1 (72.7% vs. 40.8%, $p<0.001$). Operation time (331 minutes [212–608] vs. 362 minutes [173–677], $p=0.031$) and hospital stay (12 days [8–62] vs. 14 days [7–80], $p<0.001$) were slightly longer in group 2. Bowel surgery specific complications did not differ between 2 groups (ileus, 36.4% vs. 35.7%, $p=0.925$; leakage/perforation/fistula, 3.6% vs. 6.1%, $p=0.478$). There were no significant differences in progression-free survival and overall survival between 2 groups.

Conclusion: Bowel resection performed by GO was feasible and showed equivalent oncological outcomes when compared with those by GS during maximal cytoreductive surgery for advanced ovarian cancer.

Poster (O03)

Imaging in Gynecologic Cancer
<https://doi.org/10.3802/jgo.2021.32.S1.O03>

Comparison of 2D USG based IOTA simple rules and 3D USG in preoperative assessment of adnexal masses

Rishu Goel, Seema Singhal,* Jyoti Meena, Sunesh Kumar, Smita Manchanda, Kallol Kumar Roy

All India Institute of Medical Sciences, New Delhi, India
(drseemasinghal@gmail.com)

Objective: To compare the diagnostic accuracy of “International Ovarian Tumor Analysis (IOTA) simple rules” and 3-dimensional (3D) ultrasound (USG) to discriminate benign from malignant adnexal masses.

Methods: A prospective observational study was conducted on 84 patients with adnexal masses. Both 2D USG and 3D USG with power doppler were done and IOTA simple rules applied using 2D parameters. Findings were correlated with final histopathology report.

Results: Eighty-four cases were recruited, 41 benign and 43 malignant. IOTA simple rules were conclusive in 88.1% (74/84) of cases, and the sensitivity and specificity of IOTA simple rules in conclusive cases were 83.78% (95% confidence interval [CI]=67.99–93.81%) and 89.19% (95% CI=74.58–96.97%),