depended on the age or number of chemotherapy cycles. The 19 of 26 patients who attempted for pregnancy were leading to delivery (73%). No one required infertility treatments. Mean of chemotherapy cycles and age is related to successful pregnancy. **Conclusion:** we showed patients with MOGCT could become pregnant and give birth if they desire. Advanced tumor stage was not the convincing factor for avoiding fertility preservation. Fertility sparing surgery with adjuvant chemotherapy is a safe treatment and results in high fertility rate.

Poster (O10)

Basic/Translational Science

<https://doi.org/10.3802/jgo.2021.32.S1.O10>

Identification of the therapeutic and prognostic significance of STAU2 in ovarian cancer

Xiaoshi Liu*

Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University Shanghai, China (xiaoshiliu@sjtu.edu.cn)

Objective: Of all the gynecologic cancers, ovarian cancer (OC) accounts for the most deaths worldwide. Factors such as chemotherapy and extensive abdominal metastasis may be important reasons leading to the high mortality of OC patients. Staufen homolog 2 (STAU2) is a member of the family of double-stranded RNA-binding proteins involved in cell fate decision by regulating mRNA transport, mRNA stability, translation, and ribonucleoprotein assembly. Past research have shown that STAU2, an anti-apoptotic protein, can participate in DNA replication and/or maintain genome integrity. However, the role of STAU2 in OC remains unknown.

Methods: GEPIA2 (<http://gepia2.cancer-pku.cn/#index>) is an analysis tool containing RNA sequence expression data of 9,736 tumors and 8,587 normal samples from TCGA and the GTEx projects, using a standard processing pipeline. In this study, we used the “Single Gene Analysis” module of GEPIA2 to analyze the differential mRNA expression of tumor and normal tissues, perform pathological stage analysis, and analyze the prognosis value of claudins. The p-value cutoff was 0.05. The Human Protein Atlas (<https://www.proteinatlas.org/>) provides analysis of the secreted and membrane proteins, as well as an analysis of the expression profiles for all proteins targeted by pharmaceutical drugs and proteins implicated in cancer. In this study, we confirmed the expression of STAU2 in OC via immunohistochemistry using The Human Protein Atlas. Metascape (<https://metascape.org/gp/index.html>) includes analysis of bio-pathway enrichment, protein interaction network structure, and abundant gene annotation functions based on

more than 40 public database. In this study, we performed an enrichment analysis of STAU2 and its most related genes by the Express Analysis module. TIMER (<https://cistrome.shinyapps.io/timer/>) is a comprehensive resource for systematical analysis of immune infiltrates across diverse cancer types. In this study, “Gene” module was used to evaluate the correlation between STAU2 level and the infiltration of immune cells. And we evaluated the correlation among clinical outcome and abundance of immune infiltrates expression using “Survival” module.

Results: STAU2 expression among several cancers were analyzed, and we found the transcription level of STAU2 in OC tissues was significantly higher than normal tissues. We further confirmed that STAU2 in OC tissues overexpressed compared with normal tissues via immunohistochemistry. OC patients with higher expression of STAU2 were significantly associated with poorer overall survival ($p=0.0014$) and disease-free survival ($p=0.039$). Besides, the pathological stage of OC patients was closely related to the expression of STAU2 ($p=0.0295$), concluding that STAU2 may promote the progress of OC. GO/KEGG analysis showed the differential expression of STAU2 and its related genes were mainly associated with regulation of mRNA stability, endocytosis and ubiquitin mediated proteolysis. Then we found that CD8+ T cells expression (coefficient=0.175, $p=5.64e-3$), CD4+ T cells (coefficient=0.205, $p=1.13e-3$) and macrophages (coefficient=0.353, $p=9.66e-9$) infiltration increased, while B cells (coefficient=-0.201, $p=1.36e-3$) and dendritic cells (coefficient=-0.13, $p=3.98e-2$) infiltration decreased with high STAU2 expression. The Cox proportional hazard model showed CD4+ T cells ($p=0.001$), macrophages ($p=0.001$), neutrophils ($p=0.030$) and STAU2 expression ($p=0.038$) were significantly associated with the clinical outcome of OC patients.

Conclusion: In summary, STAU2 may play a great influence on immune cell infiltration and may represent a valuable prognostic biomarker and therapeutic target for ovarian cancer.

Poster (O11)

Gynecologic Pathology, Genetics and Epidemiology

<https://doi.org/10.3802/jgo.2021.32.S1.O11>

Molecular characterization and its correlation with tumor mutation burden in Chinese ovarian cancer patients

Jia Jia Xu, Shoutai Ding, Huiling Chen, Chaoran Xia, Zhixiang Yan, Peng Luo

Topgen Biological Medicine Technology Co., Ltd., Shanghai, China (xujiajia@topgen.com.cn)

Objective: The breakthrough have been brought in therapy of ovarian cancer by poly(ADP-ribose) polymerase PARP inhibitor

and immune checkpoint inhibitor, but predictive biomarkers are lacking. We performed genomic profiling and characterized specific genes associated with an increased tumor mutational burden (TMB) in Chinese patients with ovarian cancer (OC). OC accounts for high rates of relapse and mortality among people with solid tumors worldwide. As traditional treatments are limited and often intolerant for patients with advanced ovarian cancer, PARP inhibitors and immunotherapy may breakthrough therapies patients immunotherapy could be a better option.

In order to explore the potential use of immunotherapy biomarkers, our study aimed to assess gene alteration and its correlations with microsatellite instability (MSI), TMB and tumor neoantigens burden (TNB) in Chinese CLC populations.

Methods: A total of 201 female Chinese patients with OC (including primary peritoneal cancer and fallopian tube cancer) were involved from February 2020 to July 2021. All paired specimens were detected by OncoDrug-Seq™ 603-gene panel assay through next generation sequencing using Illumina NovaSeq 6000.

Results: TP53 (62.7%) was the gene with highest mutation frequency, followed by *BRCA1* (18.9%), *KRAS* (11.9%), *ARID1A* (9.5%), *PIK3CA* (9.0%), *NFI* (6.0%), and *BRCA2* (6.0%). Further analysis showed that TP53 alterations (84.6%, n=11/13) were significantly more common in high-grade serous carcinoma than other histological subtypes. Variations detected included fusions (1.4%), duplication (4.5%), and deletion (4.0%). Among them the fusions were RET-CCDC6, BRAF-KIAA1549, NRG1-DIP2B, ERBB2-MACROD2, especially, the latter 2 fusions occurred simultaneously in one patient. Mutational incidences of significant pathway were also analyzed. Five signal pathways, including JAK-STAT (1.4%), MAPK (74.6%), EGFR (28.9%), WNT (63.7%), and TGFβ (1.0%) were related to mutational incidences. Correlation analysis between TMB and mutation status in OC (n=110) showed that the OC with RAD50 (n=4) or PALB2 (n=3) mutation patterns had higher TMB (mean values 20.3, 25.5, respectively, p<0.05), while the MSI status showed no significant difference (p>0.05).

Conclusion: The landscape of mutation patterns and MSI, and TMB among Chinese OC populations in this study will further assist the utilization of these biomarkers to immunotherapy strategies.

Poster (O12)

Rare Tumors & Metastatic Tumors

<https://doi.org/10.3802/jgo.2021.32.S1.O12>

Mucinous ovarian cancer metastasis to cervix: a rare case report

Gatot Purwoto, Raditya Perdhana, Tantri Hellyanti, Trifonia Pingkan Siregar, Siti Rafiqah Fajri

Faculty of Medicine, Indonesia University, Jakarta, Indonesia
(raditya.perdhana@gmail.com)

Objective: Ovarian cancer is the second most common gynecological malignancy, but the most lethal. Mucinous ovarian cancer (MOC) is a rare subtype of epithelial ovarian carcinoma. MOC was believed to constitute around 12% of ovarian malignancies. However, recent estimations show the true incidence to be at around 3%. It is difficult to diagnose multiple malignancies in genital organ due to almost similar spreading pattern.

Methods: Women, 45 years old with vaginal bleeding and abdominal enlargement. The diagnose of the malignancies was made from Histopathology and advanced imaging. This is a rare case of MOC spreading to cervix has been treat as double primary cancer with radiotherapy and chemotherapy.

Results: After completion the therapy (radiotherapy and chemotherapy), patient shows good response without no evidence of disease. the close follow up are need to early detection of the recurrence.

Conclusion: Radiotherapy and Chemotherapy in this rare case of MOC that spread to cervix can treat both for locally advanced cervix metastasis and MOC. A good modality for diagnose is a key to detect a pathologic disease in gynecology malignancies.

Poster (O13)

Epithelial Ovarian Cancer including Borderline Tumor

<https://doi.org/10.3802/jgo.2021.32.S1.O13>

Predictors of outcome for relapsed epithelial ovarian cancer

Rudrika Chandra, Neerja Bhatla, Sachin Khurana, MD Ray, Sarita Kumar, Lalit Kumar, Prabhat S. Malik, Sanjay Thulkar

All India of Medical Sciences, New Delhi, India (rudrika21@gmail.com)

Objective: To determine the predictors of overall survival (OS) in relapsed epithelial ovarian cancer (EOC).

Methods: Case records of patients with EOC treated between January 2010 and December 2019 were reviewed to identify cases with relapse who received treatment. Primary treatment included upfront cytoreductive surgery followed by adjuvant paclitaxel + carboplatin in 116 (45.7%) patients and neoadjuvant chemotherapy followed by interval cytoreduction in 138 (54.3%) patients. Following first relapse, 215 (84.6%) received salvage chemotherapy, 13 (5.1%) underwent surgery while 27 (10.6%) patients with biochemical relapse were kept on observation. An intent to treat analysis was done.

Results: Total 254 patients, median age 50 (range=18–81) years had relapse; median relapses-2 (range=1–9). Median progression