PHYTOCHEMICAL SCREENING AND *IN-VIVO* ANTIDIABETIC ACTIVITY OF HYDROALCOHOLIC EXTRACT OF SEEDS OF *LEPIDIUM SATIVUM* LINN

Bhagwant Mewada*, Rajendra Chouksey, Harish Pandey, Prabhakar Budholiya

Sri Satya Sai University of Technology & Medical Sciences, Sehore (M.P.)

*Corresponding Author's E mail: bhagwanmewada9@gmail.com

Received 11 Sept. 2019; Revised 17 Sept. 2019; Accepted 21 Sept. 2019, Available online 15 Oct. 2019

ABSTRACT

Diabetes mellitus (DM) is a global health problem and the incidence of DM is increasing at alarming rate all over the world. Many Indian medicinal plants have been reported to possess potential antidiabetic activity and could play important role in the management diabetes. This study sought to evaluate the antidiabetic potentials of hydroalcoholic extract of seed of *L. sativum* in diabetic rats. Qualitative analysis of various phytochemical constituents and quantitative analysis of total phenolics and flavonoids were determined by the well-known test protocol available in the literature. Quantitative analysis of phenolic and flavonoids was carried out by folins ciocalteau reagent and aluminium chloride method. Diabetes was induced in rats by Alloxan monohydrate (120 mg/kg; *i.p.*) administration. Oral treatment of hydroalcoholic extract of *L. sativum* using rat oral needle at 100 and 200mg/kg doses significantly ($P < 0.001$) decreased blood glucose and cholesterol levels in diabetic rats than control rats. In this study, Body weight, blood glucose, HDL, Total Protein, SGOT, SGPT, cholesterol and triglyceride were measured. The results showed that there was a significant decrease ($P < 0.01$) in the blood glucose, cholesterol, total protein, SGOT, SGPT and triglyceride levels in diabetic rats when compared to the normal (control) rats and increase the level of HDL and body wt. In conclusion, the above actions might be responsible for the antidiabetic activity of extract due to presence of gallic acid and other biomarkers.

Keywords: Diabetes mellitus, *Lepidium sativum*, Phytochemical screening, Alloxan monohydrate.

INTRODUCTION

Diabetes mellitus is an epidemic occurring in adults throughout the world and is the leading cause of kidney failure, heart attack, blindness and lower limb amputation. It is the fourth main cause of death in most developed countries. The prevalence of diabetes is estimated to reach 330 million by the year 2025, according to International Diabetes Federation, with the greatest potential increase being in Africa and Asia. This numerical increase will occur in developing countries. By the year 2025, over 75% of people with diabetes will reside in developing countries, as compared to 62% in 1995¹. Currently available therapies for diabetes include insulin and various oral antidiabetic agents such as sulfonylureas, biguanides and glinides. Many of them have a number of serious adverse effects; therefore, the search for more effective and safer hypoglycemic agents is one of the important areas of investigations ². Many herbs and plants have been described as possessing hypoglycemic activity when taken orally ³. According

to the World Health Organization, there are more than 1200 plant species worldwide used in the treatment of diabetes mellitus and substantial number of plant showed effective hypoglycemic activity after laboratory testing ⁴. Recently, some medicinal plants have been reported to be useful in diabetes worldwide and have been used empirically in antidiabetic and anti-hyperlipidemic remedies. More than 400 plant species having hypoglycemic activity have been available in literature, however, searching for new antidiabetic drugs from natural plants is still attractive because they contain substances which demonstrate alternative and safe effects on diabetes mellitus. Most of the plants contain glycosides, alkaloids, terpenoids, flavonoids, carotenoids etc. that are frequently implicated as having antidiabetic effect ⁵. However, the study of plant for hypoglycemic, antioxidant and hypolipidemic activities may give new pharmacological approaches in the treatment of diabetes mellitus ⁶. Alloxan is a toxic glucose analogue, which selectively destroys insulin-producing cells in the pancreas (that is beta cells) when administered to rodents and many other animal species. This causes an insulin-dependent diabetes mellitus (called "Alloxan Diabetes") in these animals, with characteristics similar to type 1 diabetes in humans. Alloxan is selectively toxic to insulin producing pancreatic beta cells because it preferentially accumulates in beta cells through uptake via the GLUT2 glucose transporter ⁷. *Lepidium sativum*, popularly named as Garden cress, member of the Brassicaceae family. It is edible herb which is growing fast and has been cultivated as a culinary vegetable in North America, Europe and all through Asia. *L. sativum* is a small, annual, herbaceous plant of 15-45 cm height, cultivated throughout India. The reddish coloured seeds are oblong, angular, slightly curved on one side having rugous surface. Seeds are odorless, pungent and mucilaginous used as salad supplement. The seeds are used in chronic enlargement of liver and spleen, as carminative adjunct to purgatives, in skin diseases, dysentery, diarrhoea, asthma and in liver complaints ⁸⁻¹⁰. Literature search revealed tachyphylactic ¹¹, diuretic ¹², oral contraceptive ¹³ antihypertensive ¹⁴ effect of *L. sativum* seeds aqueous extract. The aqueous extract of *L. sativum* seed showed hypoglycemic activity in human subjects and are used commonly in the treatment of bronchial asthma ¹⁵. Preliminary phytochemical study of *L. sativum* following standard procedure showed presence of flavonoids, coumarins, sulphur containing glycosides, triterpenes, sterols and various imidazole alkaloids ¹⁶. The major secondary metabolites of this plant are glucosinolates ¹⁷. *L. sativum* contain rare imidazole alkaloids known as lepidine and semilepidine ¹⁸. From methanolic extract of defatted seeds sinapic acid and sinapin were isolated ¹⁹. Despite the wide spread traditional and edible uses of *L. sativum*, very few pharmacological studies have been done so far. Therefore, the aim of the present study is to determine the antidiabetic activity of hydroalcoholic extract of *L. sativum*.

MATERIALS AND METHODS:

Seeds of *Lepidium sativum* were collected from local area of Bhopal in the month of August, 2018. The collected plant materials are cleaned, shade dry, coarsely powdered for further studies.

Chemical reagents

All the chemicals used in this study were obtained from HiMedia Laboratories Pvt. Ltd. (Mumbai, India), SigmaAldrich Chemical Co. (Milwaukee, WI, USA), SD Fine-Chem Chem. Ltd. (Mumbai, India) and SRL Pvt. Ltd. (Mumbai, India). All the chemicals used in this study were of analytical grade. Metformin was used as standard drugs. Streptozotocin (Sigma-Aldrich), biochemical estimation kit (Transasia Bio Medical Limited, Mumbai, India) and other chemicals and solvent obtained from Qualigens, India were used. Alloxan (Central Drug House Pvt.Ltd.,India), Glibenclamide tablets (Daonil; Aventis Pharma. Ltd., India) were procured from the medical store.

Preparation of extracts:

Defatting of plant material:

Seeds of *L. sativum* were shade dried at room temperature. The shade dried plant material was coarsely powdered and subjected to extraction with petroleum ether by maceration. The extraction was continued till the defatting of the material had taken place.

Extraction

89 gm of dried powdered seeds of *L. sativum* has been extracted with hydroalcoholic solvent (ethanol: water: 70:30) using maceration process for 48 hrs, filtered and the extracts were evaporated above their boiling points and stored in an air tight container free from any contamination until it was used. Finally the percentage yields were calculated of the dried extracts ²⁰.

Qualitative phytochemical analysis of plant extract

The *L. sativum* seeds extract obtained was subjected to the preliminary phytochemical analysis following standard methods by Khandelwal and Kokate ^{21, 22}. The extract was screened to identify the presence or absence of various active principles like phenolic compounds, carbohydrates, flavonoids, glycosides, saponins, alkaloids, fats or fixed oils, protein, amino acid and tannins.

Quantification of secondary metabolites

Quantitative analysis is an important tool for the determination of quantity of phytoconstituents present in plant extracts. For this TPC and TFC are determined. Extracts obtained from seeds of *L. sativum* plant material of subjected to estimate the presence of TPC and TFC by standard procedure.

Total phenol determination

The total phenolic content was determined using the method of Olufunmiso *et al* ²³. A volume of 2ml of each extracts or standard was mixed with 1 ml of Folin Ciocalteau reagent (previously diluted with distilled water 1:10 v/v) and 1 ml (7.5g/l) of sodium carbonate. The mixture was allowed to stand for 15 min under room temperature. The colour developed was read at 765 nm using UV/visible spectrophotometer. The total phenolic content was calculated from the standard graph of gallic acid and the results were expressed as gallic acid equivalent (mg/100mg).

Total flavonoids determination

The total flavonoid content was determined using the method of Olufunmiso *et al* ²³. 1ml of 2% AlCl₃ methanolic solution was added to 3 ml of extract or standard and allowed to stand for 15 min at room temperature; the absorbance of the reaction mixture was measured at 420 nm using UV/visible spectrophotometer. The content of flavonoids was calculated using standard graph of quercetin and the results were expressed as quercetin equivalent (mg/100mg).

Animals

Wistar rats (150-200 g) were group housed (n= 6) under a standard 12 h light/dark cycle and controlled conditions of temperature and humidity (25±2°C, 55-65%). Rats received standard rodent chow and water *ad libitum*. Rats were acclimatized to laboratory conditions for 7 days before carrying out the experiments. All the experiments were carried in a noise-free room between 08.00 to 15.00 h. Separate group (n=6) of rats was used for each set of experiments. The animal studies were approved by the Institutional Animal Ethics Committee (IAEC), constituted for the purpose of control and supervision of experimental animals by Ministry of Environment and Forests, Government of India, New Delhi, India.

Acute oral toxicity

Acute toxicity study of the prepared extracts was carried out according to the Organization for Economic Co-Operation and Development (OECD) Guidelines-423 ²⁴ the animals were fasted for 4 h, but allowed free access to water throughout. As per the OECD recommendations, the starting dose level should be that which is most likely to produce mortality in some of the dosed animals; and when there is no information available on a substance to be tested in this regard; for animal welfare reasons, The dose level to be used as the starting dose is selected from one of three fixed levels 50, 150, 300 and 2000 mg/kg body weight. Acute toxicity was determined as per reported method ²⁵.

Induction of experimental diabetes in rats

After fasting, diabetes was induced by a single intraperitoneal injection of 120 mg/kg body weight of 'Alloxan monohydrate' in distilled water. The animals were allowed to drink 5% glucose solution

overnight to overcome the drug-induced hypoglycaemia. These animals were tested for diabetes after 15 days and animals with blood glucose (fasting) were selected for experimentation ²⁶.

Standard drug used

Glibenclamide tablet (Daonil Tab.) manufactured by Aventis Pharma was used as standard drug. It was purchased from local medical store. The tablets were suspended in distilled water using tween-80 as suspending agent & used for study.

Experimental protocol

Five groups of rats were employed in the present study and each group contains six animals, as follows

Group I: Rats served as normal-control and received the vehicle (0.5 ml distilled water/day/rat)

Group II: Rats served as diabetic-control and received the vehicle (0.5 ml distilled water/day/rat)

Group III: Rats (diabetic) were administered of hydroalcoholic extract of *L. sativum* (100 mg/kg p.o.) for 15 days

Group IV: Rats (diabetic) were administered of hydroalcoholic extract of *L. sativum* (200 mg/kg p.o.) for 15 days

Group V: Rats (diabetic) were administered Glibenclamide (600µg/kg p.o.) for 15 days

Bioassay

On 15th day of treatment, blood samples were collected by retro-orbital plexus puncture method under mild ether anesthesia and serum was separated by centrifugation. Serum glucose, total protein, cholesterol (TC) and total triglyceride (TG) levels were evaluated using a commercial kit ²⁷. Body weights of rats were taken before and after treatment ²⁸.

RESULTS AND DISCUSSIONS

The crude extracts so obtained after maceration extraction process was concentrated on water bath by evaporation the solvents completely to obtain the actual yield of extraction. The yield of hydroalcoholic extracts was found to be 8.9 %. The results of qualitative phytochemical analysis of the crude powder of seed of *L. sativum* were shown in Table 1. Hydroalcoholic extracts of *L. sativum* showed the presence of flavonoids, phenols, carbohydrate and saponins. The determination of the total phenolic content, expressed as mg gallic acid equivalents and per 100 mg dry weight of sample. The total flavonoids content of the extracts was expressed as percentage of quercetin equivalent per 100 mg dry weight of sample. TPC and TFC of hydroalcoholic extract of *L. sativum* showed the content values of 0.587 and 0.874 respectively Table 2.

Table 1: Result of phytochemical screening of extracts of *L. sativum*

S. No.	Constituents	Hydroalcoholic extract
1.	Alkaloids	
	Mayer's Test	-ve
	Wagner's Test	-ve
	Dragendroff's test	-ve
	Hager's test	-ve
2.	Glycosides	
	Modified Borntrager's Test	-ve
	Legal's test	-ve
3.	Flavonoids	
	Lead acetate	+ve
	Alkaline test	+ve
4.	Phenolics	
	Ferric Chloride Test	+ve
5.	Proteins and Amino acids	
	Xanthoproteic test	+ve
	Ninhydrin Test	-ve
6.	Carbohydrates	
	Molisch's Test	+ve
	Benedict's Test	-ve
	Fehling's test	+ve
7.	Saponins	
	Froth Test	+ve
	Foam test	+ve
8.	Diterpins	
	Copper acetate test	-ve

Table 2 Total phenolic and total flavonoid content of seeds of *L. sativum*

S. No.	Extract	Total Phenol (mg/100mg)	Total flavonoid (mg/100mg)
1.	Hydroalcoholic extract	0.587	0.874

It is well known fact that alloxan monohydrate induces diabetes mellitus in rats by selective necrotic action on the beta cells of pancreas leading to insulin deficiency. Insulin deficiency leads to various metabolic aberrations in animals like increased blood glucose level, increased levels of cholesterol and triglyceride and decreased protein content ²⁹. As expected in alloxan treated rats, there was significant increase in blood glucose, cholesterol (TC) and triglyceride (TG) levels. The diabetic animals showed significant decrease in blood glucose level after 15 days treatment. Moreover, it also decreased the levels of cholesterol (TC) and triglyceride (TG) increased by alloxan treatment. Alloxan treatment of the rats has showed the loss in body weight as compared to normal rats. However, the hydroalcoholic extract was

more effective and results are comparable with that of reference drug, glibenclamide. Hence, we can say that presence of flavonoid in the ethyl acetate fraction may be responsible for antidiabetic activity.

As shown in Table 3 Blood glucose level of animals in all groups was recorded at 0, 8th and 15th day. Progressive decrease in blood glucose level was found in all treatment groups during study. At the end of experiment Glibenclamide 600µg/kg, *L. sativum* 100 and 200 mg/kg/p.o. (114.77; 122.50 and 120.00) treated group blood glucose level was decrease significantly ($p < 0.01$) at 15th days, respectively.

Table 3 Effect of hydroalcoholic extract of *L. sativum* treatment on blood glucose (mg/dl) in normal and diabetic rats

Group	Treatment	Blood glucose (mg/dl)		
		Days 0	Days 8	Days 15
I	Normal	91.65 ± 5.50	96.00 ± 5.50	100.89 ± 5.50
II	Diabetic Control	270.50 ± 10.35	285.10 ± 9.35 [#]	289.50 ± 8.35 [#]
III	Diabetic + hydroalcoholic extract of <i>L. sativum</i> (100 mg/kg)	245.50 ± 4.50	159.50 ± 4.10 ^{***}	122.50 ± 3.50 ^{***}
IV	Diabetic + hydroalcoholic extract of <i>L. sativum</i> (200 mg/kg)	254.35 ± 3.00	149.40 ± 4.00 ^{***}	120.90 ± 4.52 ^{***}
V	Diabetic + Glibenclamide (600µg/kg)	250.00 ± 4.90	135.50 ± 4.90 ^{***}	114.77 ± 4.90 ^{***}

Values are expressed as mean ± S.E.M ($n = 6$). Values are statistically significant at [#] $p < 0.001$ vs. normal group; * $P < 0.001$, ** $P < 0.01$ vs. diabetic control group (Two-way ANOVA test).

As shown in Table 4 Glibenclamide 600µg/kg, *L. sativum* 100 and 200 mg/kg/p.o. treated group biochemical parameters like TC, TG level was decrease significantly ($p < 0.01$) and total protein level was increase as compared to diabetic control group.

Table 4 Effect of hydroalcoholic extract of *L. sativum* treatment on biochemical parameters in normal and diabetic rats

Group	Treatment	TC (mg/dL)	TG (mg/dL)	Total protein(g/dl)
I	Normal	92.00 ± 3.00	87.50 ± 3.00	9.00 ± 1.50
II	Diabetic Control	183.0 ± 5.00	124.0 ± 6.00	4.70 ± 1.50
III	Diabetic + hydroalcoholic extract of <i>L. sativum</i> (100 mg/kg)	112.5 ± 5.55 ^{**}	89.50 ± 6.00 [*]	8.00 ± 2.50 ^{**}
IV	Diabetic + hydroalcoholic extract of <i>L. sativum</i> (200 mg/kg)	107.5 ± 5.50 ^{**}	86.57 ± 6.00 [*]	8.50 ± 2.50 ^{**}
V	Diabetic + Glibenclamide (600µg/kg)	101.1 ± 5.50 ^{**}	82.50 ± 6.00 [*]	8.32 ± 2.50 ^{**}

Values are expressed as mean ± S.E.M ($n = 6$). Values are statistically significant at [#] $p < 0.001$ vs. normal group; * $P < 0.001$, ** $P < 0.01$ vs. diabetic control group (One-way ANOVA followed by Tukey's post hoc test).

As represented in Table 5 body weights of animals in all groups were performed at the initial and end of the study. Body weight of animals was significantly ($p < 0.05$) maintained in all treatment groups (Glibenclamide 600 μ g/kg, *L. sativum* 100 and 200 mg/kg/p.o. 151, 155, 153) during study as compared to control group (150).

Table 5 Effects of hydroalcoholic extract of *L. sativum* on body weight

Group	Treatment	Initial weight (g)	Final weight (g)
I	Normal	155.00 $\pm$ 8.00	180.10 $\pm$ 9.00
II	Diabetic Control	160.00 $\pm$ 8.00	150.00 $\pm$ 9.00
III	Diabetic + hydroalcoholic extract of <i>L. sativum</i> (100 mg/kg)	165.00 $\pm$ 8.00	155.50 $\pm$ 8.00
IV	Diabetic + hydroalcoholic extract of <i>L. sativum</i> (200 mg/kg)	170.00 $\pm$ 8.00	153.20 $\pm$ 9.00
V	Diabetic + Glibenclamide (600 μ g/kg)	175.00 $\pm$ 8.00	151.20 $\pm$ 9.00

Values are expressed as mean $\pm$ SD of six samples from each group. (Two-way ANOVA test).

CONCLUSION

From the above results, it was concluded that the present study seems to support the claims by traditional medicine practitioners about the usefulness of *L. sativum* L seed for the treatment of diabetes. From the phytochemical investigations studies the correlation of antidiabetic activity can be made with the flavonoid compounds.

REFERENCES

1. Eseyin O, Ebong P, Eyong E, Awofisayo O and Agboke A. Effect of Telfairia occidentalis on oral glucose tolerance in rats. Afr J Pharm Pharmacol. 2010; 4: 368-372.
2. Patel DK, Prasad SK, Kumar R and Hemalatha S. An overview on antidiabetic medicinal plants having insulin mimetic property. Asian Pac J Trop Biomed. 2012; 2: 320-330.
3. Rajan M, Kumar VK, Kumar PS, Swathi KR and Haritha S. Antidiabetic, antihyperlipidaemic and hepatoprotective activity of methanolic extract of Ruellia tuberosa Linn. leaves in normal and alloxan induced diabetes. J Chem Pharm Res. 2012; 4: 2860-2868.
4. Rajasekar V, Kirbanandan S and Lakshmi BS. Assessment of antidiabetic activity of Syzygium jambolanum using in vitro model. Am J Infect Dis. 2010; 5: 99-105.
5. Malviya N, Jain S and Malviya S. Antidiabetic potential of medicinal plants. Acta Pol Pharm 2010; 67: 113-118.
6. Dangi KS and Mishra SN. Antihyperglycemic, antioxidant and hypolipidemic effect of Capparis aphylla stem extracts in streptozotocin induced diabetic rats. Biol Med 2010; 2: 35-44.

7. Lenzen, S. The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia*. 2008; 51, 216-226.
8. Khory R. *Materia Medica of India and Their Therapeutics*. Komal Prakashan, New Delhi, India, 1999; pp: 63-64.
9. Kirtikar KR and Basu BD. *Indian Medicinal Plants*. Vol. 1, Popular Prakashan Pvt. Ltd., Allahabad, Uttar Pradesh, India, 2006; pp: 174-175.
10. *The Wealth of India*. Raw Material. Vol. 6, CSIR Publication, New Delhi, India, 1962; pp: 71.
11. Vohra SB and Khan MS. Pharmacological studies on *Lepidium sativum* linn. *Ind. J. Physiol. Pharmacol.* 1977; 21: 118-120.
12. Navarro E, Alonso J, Rodriguez R, Trujillo J and Boada J. Diuretic action of an aqueous extract of *Lepidium latifolium* L. *J. Ethnopharmacol.* 1994; 41: 65-69.
13. Sharief M and Gani ZH. Garden cress *Lepidium sativum* seeds as oral contraceptive plant in mice. *Saudi Med. J.* 2004; 25: 965-966.
14. Eddouks M, Maghrani M, Zeggwagh NA and Michel JB. Study of the hypoglycaemic activity of *Lepidium sativum* L. aqueous extract in normal and diabetic rats. *J. Ethnopharmacol.* 2005; 97: 391-395.
15. Eddouks M, Maghrani M, Zeggwagh NA and Michel JB. Study of the hypoglycaemic activity of *Lepidium sativum* L. aqueous extract in normal and diabetic rats. *J. Ethnopharmacol.* 2005; 97: 391-395.
16. Patel U, Kulkarni M, Undale V and Bhosale A. Evaluation of diuretic activity of aqueous and methanol extracts of *Lepidium sativum* garden cress (Cruciferae) in rats. *Trop. J. Pharmaceut. Res.* 2009; 8: 215-219.
17. Gill V and MacLeod AJ. Studies on glucosinolate degradation in *Lepidium sativum* seed extract. *Phytochemistry*. 1980; 19: 1369-1374.
18. Maier UH, Gundlach H and Zenk MH. Seven imidazole alkaloids from *Lepidium sativum*. *Phytochemistry*. 1998; 49: 1791-1795.
19. Nayak PS, Upadhyaya SD and Upadhyaya A. A HPTLC densitometer determination of synaptic acid in chandrashur (*Lepidium sativum*). *J. Sci. Res.* 2009; 1(10): 1196.
20. Mukherjee PK. *Quality control of herbal drugs*. 2nd Ed. Business Horizons; 2007.
21. Khandelwal KR. *Practical pharmacognosy technique and experiments*. 23rd Ed. Nirali Prakashan; 2005.
22. Kokate CK. *Practical pharmacognosy*. 4th Ed. Vallabh Prakashan; 1994.

23. Olufunmiso OO and Afolayan AJ. Phenolic content and antioxidant property of the bark extract of *Ziziphus mucronata* willd. Subsp. *mucronata* willd, BMC Complement. Alternative Medicine. 2011; 11:130.
24. Guideline Document on Acute oral Toxicity Testing, Series on Testing and Assessment No. 423. Paris: Organization for Economic Co-Operation and Development, OECD Environment, Health and Safety Publications; 1996. Available from: <http://www.oecd.org/ehs>.
25. Jonsson, M, Jestoi M, Nathanail AV, Kokkonen UM, Anttila M, Koivisto P and Peltonen, K. Application of OECD Guideline 423 in assessing the acute oral toxicity of moniliformin. Food and chemical toxicology. 2013; 53, 27-32.
26. Shankar MB, Parikh JR, Geeta M, Mehta RS and Saluja AK. Anti-diabetic activity of novel andrusterane derivatives from *Syzygium cumini* Linn. J of Natural Remedies. 2007; 7(2): 214-219.
27. Adeneye AA, Ajagbonna OP and Ayodele OW. Hypoglycemic & antidiabetic activities on the stem bark aqueous & ethanol extracts of *Musanga cecropioides* in normal & alloxan induced diabetic rats. Fitoterapia. 2007; 78:502-505.
28. Roy S, Sehgal R, Padhy BM, Kumar VL. Antioxidant and protective effect of latex of *Calotropis procera* against alloxan induced diabetic rats. J of Ethnopharmacology. 2005; 102: 470-3.
29. Dhanabal SP, Mohan Maruga Raja MK, Ramanathan M, Suresh B. Hypoglycemic activity of *Nymphaea stellata* leaves ethanolic extract in alloxan induced diabetic rats. Fitoterapia 2007; 78 : 288-291.