Aluminum phosphide poisoning and development of hemolysis and methemoglobinemia

Sir,

I read with interest a case, presenting with hemolysis and methemoglobinemia associated with aluminum phosphide (ALP) poisoning presented by Soltaninejad and colleagues, published in your journal.^[1] Two major points should not be missed about this case; firstly, although the authors have stated that methemoglobinemia is a rare finding following ALP poisoning, it seems that in all patients with this poisoning, methemoglobinemia is present to some degrees.^[2-6] Therefore, it should be emphasized that symptomatic methemoglobinemia-but not methemoglobinemia itself- is a rare finding in this setting. Secondly, in a previously performed study on two patients, presented by two authors of this same article, it has been stated that the ALP-poisoned patients with hemolysis and methemoglobinemia seem to be resistant to methylene blue and ascorbic acid,^[4] while in the current case report, they have suggested that they may play a role in the successful management of these patients. The authors hypothesize that this difference in response to therapy may be due to the severity of toxicity, inappropriate dose of methylene blue, and the physiological differences between the patients.^[1] The point they may have missed is gastric lavage with potassium permanganate. As you know, potassium permanganate is a strong hemolytic and oxidizing agent that can induce hemolysis and methemoglobinemia.^[6-8] Therefore, lavage by the recommended concentration (1:10,000) of this agent may not only cause methemoglobinemia and hemolysis, but also it may cause methemoglobinemia unresponsive to the treatment. This needs further considerations in the future studies. Thanks for this interesting case.

Hossein Sanaei-Zadeh

Medical School, Shiraz University of Medical Sciences, Shiraz, Iran

Correspondence:

Assoc. Prof. Hossein Sanaei-Zadeh, Medical School, Shiraz University of Medical Sciences, Emergency Room/Division of Medical Toxicology, Hazrat Ali-Asghar (p) Hospital, Meshkinfam Street, 7143918796 Shiraz, Iran.
E-mail: h-sanaiezadeh@tums.ac.ir

References

1. Soltaninejad K, Nelson LS, Khodakarim N, Dadvar Z, Shadnia S. Unusual complication of aluminum phosphide poisoning: Development of hemolysis and methemoglobinemia and its successful treatment. *Indian J Crit Care Med* 2011;15:117-9.
2. Lall SB, Peshin SS, Mitra S. Methemoglobinemia in aluminium phosphide poisoning in rats. *Indian J Exp Biol* 2000;38:95-7.
3. Lakshmi B. Methemoglobinemia with aluminum phosphide poisoning. *Am J Emerg Med* 2002;20:130-2.
4. Shadnia S, Soltaninejad K, Hassanian-Moghadam H, Sadeghi A, Rahimzadeh H, Zamani N, *et al.* Methemoglobinemia in aluminum phosphide poisoning. *Hum Exp Toxicol* 2011;30:250-3.
5. Mostafazadeh B, Pajoumand A, Farzaneh E, Aghabiklooei A, Rasouli MR. Blood levels of methemoglobin in patients with aluminum phosphide poisoning and its correlation with patient's outcome. *J Med Toxicol* 2011;7:40-3.
6. Sanaei-Zadeh H. Response to "blood levels of methemoglobin in patients with aluminum phosphide poisoning and its correlation with patient's outcome". *J Med Toxicol* 2012;8:86-7.
7. Mahomed MC, Mahomed YH, Canham PA, Downing JW, Jeal DE. Methaemoglobinemia following treatment dispensed by witch doctors. Two cases of potassium permanganate poisoning. *Anaesthesia* 1975;30:190-3.
8. Jetter WW, Hunter FT. Death from attempted abortion with a potassium permanganate douche. *N Engl J Med* 1949;240:794-8.

Access this article online

Quick Response Code:

Website:

www.ijccm.org

DOI:

10.4103/0972-5229.106519