The Utility of Spirometry in Assessment of Presumptive Diagnosis of Bronchial Asthma in a Nigerian Tertiary Hospital

Sir,

I read with interest the study by Nwosu *et al.*^[1] published in July–December 2019 issue of the *Indian Journal Respiratory Care*. The authors studied the role of spirometry in the diagnostic evaluation of bronchial asthma (BA) and defined the ventilatory patterns of patients with a presumptive diagnosis of BA using spirometry and the proportion of those who showed positive reversibility test (RT). They found that normal, obstructive, restrictive, and mixed ventilatory patterns were found in 43.8%, 34.8%, 11.2%, and 10.1% of participants, respectively. Moreover, RT test was positive in 53.6% of the studied cohort with a prebronchodilator obstructive pattern of 47.4% and mixed ventilatory pattern of 66.7%.^[1] They concluded that spirometry is useful in confirming the diagnosis of asthma, but clinicians must be aware that normal spirometry does not exclude asthma and that follow-up spirometry might be in need to further assess asthmatic patients with normal spirometric findings.^[1] Apart from a few study limitations addressed by the authors, I assume that the following methodological limitation might cast further suspicions on the study results. In the methodology, the authors mentioned that spirometry maneuver and interpretation were performed according to the Global Initiative on Chronic Lung Disease and American Thoracic Society and European Respiratory Society (ERS) guidelines. Ventilatory functions were accordingly classified as normal, obstructive, and restrictive patterns. The ERS reference values were employed, and a correction factor of 10% was considered to adjust for the African study population.^[1] It is obvious that interpreting ventilatory patterns requires reference values obtained from healthy subjects of the same ethnic origin for that particular population. Actually, the guidelines employed by the authors were constructed for the Caucasian population. They are not suitable to be employed for the Nigerian population as applying prediction formula derived for the Caucasian population was noticed to overestimate the values for African ethnic population.^[2] It is worthy to mention that spirometric standards for healthy general adult Nigerian populations have been already constructed to be employed in the researches and clinical settings.^[3] I assume that employing these national standards in the study methodology could yield

a more precise idea on the role of spirometry in the diagnostic assessment of BA.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Mahmood Dhahir Al-Mendalawi

Department of Paediatrics, Al-Kindy College of Medicine, University of Baghdad, Baghdad, Iraq

Address for correspondence: Prof. Mahmood Dhahir Al-Mendalawi, P. O. Box 55302, Baghdad Post Office, Baghdad, Iraq.
E-mail: mdalmendalawi@yahoo.com

REFERENCES

1. Nwosu NI, Chukwuka CJ, Onyedum CC, Odilinye HC, Nlewedim PI, Ayuk AC. The utility of spirometry in assessment of presumptive diagnosis of bronchial asthma in a Nigerian tertiary hospital. *Indian J Respir Care* 2019;8:102-6.
2. Njoku CH, Anah CO. Reference values for peak expiratory flow rate in adults of African descent. *Trop Doct* 2004;34:135-40.
3. Fawibe AE, Odeigah LO, Saka MJ. Reference equations for spirometric indices from a sample of the general adult population in Nigeria. *BMC Pulm Med* 2017;17:48.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.ijrconline.org
	DOI: 10.4103/ijrc.ijrc_31_19

How to cite this article: Al-Mendalawi MD. The utility of spirometry in assessment of presumptive diagnosis of bronchial asthma in a Nigerian tertiary hospital. *Indian J Respir Care* 2020;9:126.

Received: 08-07-2019 **Accepted:** 17-07-2019 **Published:** 08-01-2020

© 2020 Indian Journal of Respiratory Care | Published by Wolters Kluwer - Medknow