Nipple and areolar reconstruction with tattoo pigments, grafts and local flaps

John M Drever MD FRCSC

Department of Plastic Surgery, Etobicoke General Hospital, Etobicoke, Ontario

JM Drever. Nipple and areolar reconstruction with tattoo pigments, grafts and local flaps. Can J Plast Surg 1993;1(2):84-90.

In order to obtain symmetry when reconstructing the nipple and areola, the breast mounds are worked on first. Having reconstructed one side, the volume and shape of the opposite breast is modified as needed; then the areola can best be imitated on the reconstructed breast by placing tattoo pigments on the dermis of a thin skin graft. If the areola that is to be matched is light in colour, it is preferable to tattoo the skin directly. This is done as the initial procedure, as an improvement or as 'maintenance' to an already tattooed areola. The nipple reconstructed by grafting part of the opposite nipple or section of areola produces the most natural looking results. Another method involves lifting a tulip-shaped, centrally based dermis flap and then covering the areola and this flap with a thin skin graft with tattoo pigments smeared on its dermis. A nipple is made of the breast skin when the mound is revised by taking a segment off its lower quadrants.

Key Words: Breast reconstruction, Mastectomy and reconstruction, Nipple and areolar reconstruction

Reconstruction du mamelon et de l'aréole avec tatouage, greffes et lambeaux

RÉSUMÉ : Pour obtenir une symétrie lors de la reconstruction du mamelon et de l'aréole, la glande mammaire est remodelée d'abord. Après reconstruction d'un côté, le volume et la forme de l'autre sein sont appariés, puis l'aréole peut être imitée en disposant les pigments de tatouage sur la mince peau de la greffe. Si l'aréole qui doit être assortie est de couleur pâle, il est préférable de tatouer directement la peau. Il s'agit de la première étape de l'amélioration ou du «maintien» d'une aréole déjà tatouée. C'est le mamelon reconstruit à partir d'une greffe partielle du mamelon ou d'une section de l'aréole du sein opposé qui donne l'aspect le plus naturel. Une autre méthode consiste à soulever un lambeau en forme de tulipe à la base du derme central et à recouvrir l'aréole et ce lambeau avec un greffon de peau dont le derme est tatoué au préalable. Un mamelon est construit à partir de la peau du sein lorsque la glande a été remodelée, à partir d'un segment prélevé aux quadrants inférieurs.

TATTOOING DIRECTLY ON THE SKIN SURFACE

Indications

Tattooing directly on the skin surface (1-7) is most useful for imitating an areola with a soft transition on the perimeter and with a smooth surface (Figure 1). I also use it to improve previously reconstructed nipples that have the white scar 'target' appearance, to camouflage the areola perimeter scars and for general 'maintenance' on areolas that have depigmented with time. Tattooing the opposite nipple and areola with the same pigments as the reconstructed one is useful for achieving symmetry.

Instruments

There are two basic types of machines used to drive the tattooing needles up and down. One is the electromagnet type that moves a shaft by drawing two contact points. The other is the mechanical type of a camshaft that is driven by an electric motor or by pneumatic power. There are many models of these instruments on the market, available either from 'medical' suppliers or from the less expensive and just as

effective outfitters for the professional tattoo artists (eg, Spaulding & Rogers Mfg Inc, Voorheesville, New York).

I have tried many different machines and found that a pneumatic reciprocating saw with a chuck fitted with a six needle flat shader is a very effective tool. This type of saw is usually available in the operating rooms of nearly every hospital in either the plastic surgery or the orthopedic departments. I use a tattoo machine purchased from the professional tattoo artists who also carry the entire line of 'skin shades' pigments.

Tattoo pigments

The pigments used for imitating natural skin colour are inorganic substances also called 'earths'. The brown colour contains brown iron oxide and titanium dioxide, dissolved in a mixture of glycerin, alcohol and water.

Tattooing technique

The tattooing technique provides a way of permanent body painting, with the intention of creating an aesthetic symmetry. Pigments must be placed so as to copy the opposite areola in its symmetrical location, colour tones, shades and, if possible, its surface variations.

The markings of the position and dimensions should be made with the patient in an upright position marking the

Correspondence and reprints: Dr John M Drever, 135 Queen's Plate Drive, # 560, Toronto, Ontario M9W 6V1. Telephone/Fax (416) 745-8898

Figure 1 (Left) Before and three years after a lower abdominal transverse rectus abdominus myocutaneous (LATRAM) reconstruction. (Above) The areola was created by tattooing on the skin surface; the nipple was made by sharing with the opposite, cutting it transversely across

Figure 2 (Left) Before and two years after a breast reconstruction created with a LATRAM flap. (Above) Nipple created by sharing half of the opposite nipple, and the areola by tattooing under a thin skin graft. This technique gives a more natural wrinkled look

perimeter of the areola. The model areola that has not been operated upon is seldom perfectly round, so I copy its contour by taking the maximum and minimum diameters, then marking a roundish shape as a mirror image on the reconstructed breast. Consideration should be given to the fact that a reconstructed mound with a rectus abdominus myocutaneous (RAM) flap tends to drop 1 or 2 cm in the first year after the reconstruction and the nipple of a reduced breast occasionally tends to rise from the initial surgical markings. There can also be quite a variation in the areola between its contracted and relaxed state, so allowances for these changes are made during these markings.

Once the area to be tattooed is outlined, I coat it with sterile petrolatum gel and start mixing the tattoo pigments. To obtain a shading effect I mix three colours; beige (also called

'tan' or 'flesh' by different manufacturers) (8), light brown and a very small amount of red (commonly found around the nipple in Caucasians). Black is used only in small amounts in people with very dark skin. These pigments can be placed directly over the skin or into the receptacle located in the tube that encases the needle shaft. The darker colours are placed in the centre, getting lighter towards the perimeter. It is important to hold the breast skin quite taught as the needles are applied. The edges of these should be held parallel to the surface of the skin, ensuring one corner of the 'comb' of needles does not penetrate deeper than the other, to avoid causing undesired lacerations. One should also avoid 'over-working' an area (passing the needles several times over the same area). This can create a deeper abrasion that tends to bleed and scab with consequent pigment loss. The needles

Figure 3) Biopsy of an areola made by tattooing under a thin skin graft. Notice how the pigment clumps are located in a deeper area than when tattooing directly on the skin surface. Pigments located in this relatively avascular area tend to last longer than when tattooing is done the traditional way

should be passed in short semicircular motions working from distant to near and frequently wiping the excess off to observe the results. The immediately observed colour will not be the end result because the abrasion effect of the needles produces a reddish tone. Therefore it is very important to select the correct shades of pigments and their location before applying the needles. It is preferable to err on the light than on the dark side, for retouching the area two weeks later is much easier than trying to remove the tattoo. I prefer to place the selected pigments on the skin surface rather than in the receptacle of the needle shaft, so I draw and mix them inside a 3 mL syringe.

Dressings and activities are the usual for any minor abrasion.

TATTOOING UNDER A THIN SKIN GRAFT

Indications

This technique (9,10) gives the most natural results when I want to create an areola with an abrupt demarcation at its perimeter and with a somewhat wrinkled surface. Therefore, the most common situation that this technique applies to is when I am planning to do a contralateral reduction on a breast that has a darkish and wrinkled areola. Another indication is when performing a nipple and areolar reconstruction with the central dermal flap.

Technique

The donor area of a skin graft will be tender for several days and may be a source of complications. Therefore, whenever possible, I take the skin graft from the discarded skin of the reduced breast. After doing the standard markings for a breast reduction with the patient in the upright position, and marking the new nipple placement on the reconstructed breast, the graft is removed with a power dermatome. This dermatome cuts through the dermis very evenly without leaving the stroke marks that show up as definite lines when we use a manual dermatome. The removed graft should be thin and translucent enough to permit the pigments to show through it. This thin skin graft is placed dermis side upwards on an oily gauze and the selected shade of pigments are smeared on it. Care should be taken to wipe off any clumps of pigments with a dry gauze and then, if necessary, reapply darker tones until the best match is acquired, leaving no excess pigment that could hinder the take of the graft. A satisfactory cosmetic result is seen when darker pigments are placed towards the centre and coma shaped Montgomery glands are imitated (Figure 2). Then, after cutting a hole for the grafted nipple to come out, we place the graft on the de-epithelialized area, tailor its edges and immobilize it with paper strip tapes. A moulded oily gauze is then positioned and held in place with more paper strips for at least two weeks.

Evolution and comments

Since 1987, when I started using this technique, 158 areolas have been reconstructed this way. The results are very realistic and have improved as I learned more about mixing pigments to imitate the opposite areola. Complications or partial takes have been very rare, especially since I became careful to remove any pigment clumps. I have noted some degree of fading over the years but much less than the pigment loss that occurs when tattooing directly over the skin. Observing biopsies of areolas performed this way, the pigment can be seen lying in the fibrous interphase between the recipient dermis and the graft (Figure 3). Perhaps this relatively avascular location of the pigment may explain its longevity.

NIPPLE RECONSTRUCTION BY SHARING

Indications

Having reviewed a large number of cases, I have no doubt that the best looking nipples are made with composite grafts of the opposite nipple-areolar complex. Naturally, these are limited by patient refusal, absent donor tissue or for oncological reasons.

Techniques

Donor nipple divided transversally across: This technique (3,11-13) is used when there is a large remaining nipple; removing half of it not only gives an excellent reconstruction but also improves the looks of the donor. I use a #20 scalpel blade after having firmed the nipple with a skin hook and

divided it transversely through the middle. The stump is then closed with a purse-string of 6-0 reabsorbable suture material. Stitches of the same suture material are used to immobilize the graft to the dermis donor area avoiding any crushing of the graft. A moulded bolus dressing of oily medicated gauze is placed over it and left untouched for at least two weeks.

This technique is usually coupled with an areolar reconstruction with the pigmented skin graft so the bolus dressing will cover both.

Donor nipple divided tangentially (3,7,14,15): This is usually performed when the donor nipple has less than 1 cm of projection. I remove the tissue tangentially so the remaining nipple will not lose much height. Usually conducted in combination with a skin tattoo on the surrounding areola, the tattoo is made first, placing the darkest pigments on the centre of the areola. Then a 1 cm diameter U shaped incision is made on the centre of the tattooed area and lifted as a trap door flap from the deep dermal level. This has the great advantage of doubling the bleeding recipient surface which increases the take of the graft. Then a skin hook is placed on the edge of the donor nipple, and with a #20 blade, a composite graft is removed that includes half of the top of the nipple plus one side and part of the areola. This graft is then inserted into the previously tattooed trap door flap and carefully sutured with 6-0 reabsorbable material (Figure 4). The donor site is closed with mattress stitches located on the base of the nipple.

Areolar edge as donor (3,12)(Figure 5): I choose this donor site for nipple reconstruction when the donor areola is smooth and light coloured but not large enough for sharing half its area, and the breast needs a reduction which would leave a scar around the areola anyway. It is also used when the nipple has very little projection or the patient refuses to have it touched.

First, I mark the areolar diameter for the reduction mammoplasty; then a crescent strip along the outer remaining areola, about 2 to 3 cm long by 1 cm wide, is removed. This full thickness graft is folded in half and sutured to itself along the middle and to the de-epithelialized donor site with 6-0 reabsorbable suture material. I use the same dressing as previously mentioned, an oily medicated moulded gauze that also covers the areolar area, and immobilize it with paper tapes for at least two weeks.

NIPPLE AND AREOLAR RECONSTRUCTION WITH THE TUBULAR DERMIS FLAP AND A PIGMENTED SKIN GRAFT

Indications

Bilateral breast reconstruction is the most common indication for the use of this technique (10). Symmetrical and well defined nipple-areolar complexes are the usual result. I also use it in patients who refuse to have the opposite nipple or areola operated on.

Figure 4) Donor nipple removed tangentially. (Top left) With a skin hook on the edge of the nipple, a section is removed that also includes part of the areola. (Top right) This composite nipple and areola graft is placed into a previously tattooed, small trap door flap that will enhance the 'take' and give a better transition between the nipple and the areola. (Bottom left) The composite graft is sutured with 6-0 stitches. (Bottom right) The donor site is closed with one or two mattress stitches at the base of the nipple

Technique

Having marked the areolar perimeter, an incision is made on it down to the deep dermis. Then we proceed to take a thin skin graft out of this circle (Figure 6) using a small manual dermatome, holding the skin tight with one hand and removing thin skin strips with the other. After completing this first de-epithelialization I carve the dermis flap by undermining a second layer off the deep dermis, taking care to preserve some fibrous strands over the fatty layer. This dermis-only flap is left attached to a 1 cm central circle that will become the pedicle of the nipple; the floppy surrounding dermis is now shaped into an upright tube. Two purse-string stitches of reabsorbable material, one half way up and the other at the top, are used to make this tube that can be about 2 cm high. Once this pole or tube is fashioned I proceed to cover it, and the surrounding raw areolar area, with one piece of pigmented skin graft.

The donor site for this graft should be 8 to 10 cm in diameter on each side. I use the traditional concealed areas as the previously removed skin is too small for this purpose. The split-thickness graft is removed with a power dermatome, then the selected pigments are smeared on its dermal surface the same way described for tattooing under a thin skin graft, but the darker centre is made over an area 2 to 3 cm in diameter. The skin graft is then placed evenly over and around the dermal pole, not worrying about the radiating wrinkles but removing all the air pockets. The excess that passes over the edge of the areola is trimmed and the graft is immobilized with multiple paper tapes. A medicated oily gauze shaped into a ring is placed over the grafted pole and further stabilized with more paper tapes.

This dressing is left untouched for two weeks. When uncovered, one has to resist the desire to reduce its size, for it will take one or two months for the dermis to undergo remodelling and finally reduce to a 1 cm high nipple.

Figure 5 (Left) Before and four years after a LATRAM reconstruction; a breast reduction was carried out on the left breast. (Above) The areola was tattooed and the nipple was made by removing a rim from the opposite areola at the same time as the reduction, and setting it into a trap door flap

Figure 6 Nipple and areola reconstruction with a tubular dermis flap and a pigmented skin graft. (Top left) Skin is held taut with one hand and, using a small dermatome, a very thin layer of skin is removed with the other hand. (Top right) A dermis only flap is raised off the deep dermis, leaving it attached only to a central pedicle of about 1 cm in diameter. (Bottom left) The dermis-only flap is shaped into a tube by placing two purse-string stitches of 5-0 absorbable material. (Bottom centre) A thin 8 x 8 cm skin graft is spread out and its dermis side is smeared with tattoo pigments. Darker colours are placed in the centre and Montgomery glands can be imitated. Care is taken to remove any clump of pigment. (Bottom right) The dermal tube is now covered with the prepigmented skin graft, trimmed and a molded oily gauze is placed over it

Figure 7 (Left) Before and five years after a breast reconstructed with a LATRAM flap, and a breast reduction with a free nipple graft on the left. (Above) Close-up of the nipple and areola reconstruction, using a full thickness graft by sharing half of the opposite. The graft on the reconstructed breast had partially depigmented, which is common, but coloration returned after two years

Evolution and comments

This technique has produced very natural and lasting results, though the original colours tend to fade after several years. Also, the nipple projection seems to flatten and acquire a semi-spherical appearance which did not seem to bother any of the patients. So far 48 nipples/areolas have been constructed using this technique, including 17 patients with bilateral reconstruction. In eight cases a scab has formed with eventual loss of part of the nipple.

NIPPLE AND AREOLAR RECONSTRUCTION WITH FULL THICKNESS GRAFTS BY SHARING HALF OF THE OPPOSITE

Indications

This technique (3,15,16) is used when planning the opposite breast reduction which has a very large areola.

Technique

The simplest way of performing this technique involves halving the donor nipple-areolar complex and using each half as a free composite graft placed on the most prominent part of each breast. It is preferable to divide the nipple-areolar complex by drawing a large S or a conjoined spiral across the middle. An alternative approach, which is especially useful when doing a breast reduction with a parenchymal pedicle, is to use as a graft the outer perimeter of areola that is usually discarded. This full thickness areolar graft is cut into a long strip; then one end is folded onto itself to form the nipple and the rest is sutured around it in a snail or 'jelly roll' fashion. The bolus dressing is the same as previously described.

Evolution and comments

Though these full thickness grafts are time consuming and meticulous to perform, the final results are usually very gratifying and after years they look better than many of the previously described techniques. It is not uncommon for these grafts to initially undergo some discoloration with loss of pigment, especially in dark skinned people, but this tends to return in one or two years (Figure 7).

The scars inside the areola can be tattooed if too visible. I have used these full thickness grafts to reconstruct 63 nipple-areola complexes. There have been five cases of minor areas of skin loss, but no total nipple losses. The low incidence of tissue loss can be explained by the fact that the grafting was performed on well vascularized donor areas such as RAM flaps or opposite reductions.

NIPPLE RECONSTRUCTION AS PART OF THE MOUND REVISION

Indications and technique

Breast mounds made with RAM flaps tend to have a rather flat surface that can be transformed into a more conical shape by removing a pie-shaped section of skin and fat from the lower quadrants. When doing this operation, I have reconstructed the nipple with a 1 x 2 cm skin flap left attached at

Figure 8 Nipple reconstruction as part of mound revision. **(Top)** A diamond-shaped skin and fat excision is made in which the cephalad apex will be the nipple. It is marked as a 1 x 2 cm skin flap on the most prominent part of the cone. The area around it, including this rectangular flap, were previously tattooed. On the right is the diamond-shaped tissue which has been removed. **(Centre)** The edges of the excision are brought together at the inframammary fold, which will be much better defined, and the breast will have a more conical shape. The nipple will be made by folding in half the rectangular small skin flap. **(Bottom)** The circle of the areola is completed by tattooing on the skin and the nipple is sutured to form a tube

the top of the excised skin wedge (Figure 8). Before the excision, I tattoo the area that will become the nipple flap, then I excise a diamond-shaped piece of skin and fat down to fascia with its widest point at the inframammary fold that will give the breast a cone shape. The small rectangular flap is folded on itself and sutured, forming a nipple. The wedge edges are sutured in two layers which will better define the inframammary fold, and finally the rest of the areola is tattooed. This technique forms very adequate nipples and at the same time improves the shape of the breast.

REFERENCES

1. Rees, TD. Reconstruction of the breast areola by intradermal tattooing and transfer. *Plast Reconstr Surg* 1975;55:560-8.
2. Rubin LR. The Burned Breast. *NY State J Med* 1975;75:865-71.
3. Gruber RP. Nipple and areola reconstruction: a review of techniques. *Clin Plast Surg* 1979;6:1,71-92.
4. Becker H. The use of intradermal tattoo to enhance the final result of nipple and areolar reconstruction. *Plast Reconstr Surg* 1986;77:673-4.
5. Little JW III, Spear SL. Finishing touches in N-A reconstruction. *Perspect Plast Surg* 1988;2:1.
6. Spear SL, Convit R, Little JW III. Intradermal tattoo as an adjunct to nipple and areolar reconstruction. *Plast Reconstr Surg* 1989;83:907.
7. Bostwick John III. Finishing touches. *Plastic and Reconstructive Breast Surgery*, Vol 2. St Louis: Quality Medical Publishing, 1990:1163.

Drever

8. Spaulding H. Tattooing A to Z – A guide to successful tattooing. ISBN #0929719-00-X.
 9. Drever JM. Tattooing the reconstructed areola without a tattooing machine. *Plast Reconstr Surg* 1988;81:995.
 10. Drever JM. Nipple and areolar reconstruction with the pigmented skin graft technique. EF of the American Society of Plastic and Reconstructive Surgery Video Tape No 9204; 1991.
 11. Cronin TD, Upton J, McDonough JM. Reconstruction of the breast after mastectomy. *Plast Reconstr Surg* 1977;59:1-11.
 12. Gruber RP. Method to produce better areolar and nipples on reconstructed breasts. *Plast Reconstr Surg* 1977;60:505-12.
 13. Millard DR, Devine J, Warren WD. Breast reconstruction: A plea for saving the uninvolved nipple. *Am J Surg* 1971;122:763-71.
 14. Cocke WM Jr. Nipple and Areolar Complex in Breast Reconstruction Following Mastectomy for Cancer. Boston: Little, Brown & Co, 1977.
 15. Wexler NR, O'Neil RM. Areolar sharing to reconstruct the absent nipple. *Plast Reconstr Surg* 1973;51:176-82.
 16. Schwartz AW. Reconstruction of the nipple and areola. *Br J Plast Surg* 1976;29:230-8.
-
-