The tumescent technique for bloodless breast reduction

Massey Beveridge MD, Michael Bell MD FRCSC FACS
Ottawa Civic Hospital, Ottawa, Ontario

M Beveridge, M Bell. The tumescent technique for bloodless breast reduction. Can J Plast Surg 1994;2(3):121-124. The tumescent technique for reduction mammoplasty provides a safe, reliable and inexpensive way to reduce intraoperative blood loss in reduction mammoplasty. The results of a study involving 64 consecutive patients operated on by the same surgeon are presented. Initial subjective experience with the infiltration of large volumes of very dilute anaesthetic solution deep in the breast along the planned incision lines suggested that the technique significantly reduced blood loss. To confirm this, 23 consecutive cases were operated upon without infiltration and the next 20 cases were done with infiltration of 250 mL of 0.1% lidocaine with 1:1,000,000 adrenaline. Preoperative hemoglobin was compared with that obtained at an office visit on the seventh postoperative day. The difference in the drop in hemoglobin observed in the two groups was significant at the $P < 0.02$ level; however, the difference was most pronounced in the group in whom a smaller volume of tissue was removed. Accordingly in the next phase of the study the volume of solution to be injected was calculated in relation to bodyweight. When 10 mL/kg bodyweight of solution was injected there was dramatically less bloodloss compared with the control group ($P < 0.01$). Patients experienced no ill effects attributable to the injection; there were no significant postoperative hematomas nor any adverse effect on wound healing.

Key Words: Bloodloss, Reduction mammoplasty, Surgical technique, Transfusion

Technique d'infiltration dans la réduction mammaire

RÉSUMÉ : La technique d'infiltration utilisée dans la réduction mammaire offre un moyen sûr, fiable et peu coûteux de réduire les pertes sanguines peropératoires. Les résultats d'une étude portant sur 64 patientes consécutives opérées par le même chirurgien sont présentés ici. L'expérience subjective initiale avec infiltration de forts volumes d'une solution très diluée d'anesthésique dans le sein, le long des incisions prévues donne à penser que cette technique réduit significativement les pertes sanguines. Pour confirmer ceci, 23 cas consécutifs ont été effectués sans infiltration et les 20 cas suivants ont été faits avec infiltration de 250 mL de lidocaïne à 0,1 % avec épinéphrine à 1:1 000 000. L'hémoglobine préopératoire a été comparée à l'hémoglobine prélevée lors d'une visite au cabinet médical le septième jour suivant l'opération. La différence quant à la chute d'hémoglobine observée entre les deux groupes était significative $P < 0,02$. Toutefois, nous avons remarqué que la différence était plus marquée dans le groupe chez qui un volume plus petit de tissu avait été extrait. Par conséquent, à la phase suivante de l'étude, le volume de solution injecté a été calculé en rapport avec la masse corporelle. Lorsque 10 mL/kg de poids corporel de solution étaient injectés, on notait une différence significative au plan de la perte sanguine en comparaison avec le groupe témoin ($P < 0,01$). Les patientes n'ont manifesté aucun effet secondaire lié à l'injection; on n'a noté aucun hématome post-opératoire significatif ni de réaction indésirable au niveau de la cicatrisation de la plaie.

Reduction mammoplasty is a commonly performed elective procedure that may result in clinically significant bloodloss. The possibility of disease transmission makes most surgeons reluctant to advise transfusion and there is a considerable financial cost to having autologous blood on hand for each case. Most patients tolerate a transient anemia well enough but undoubtedly they feel better and get back to regular activities faster if operative bloodloss is minimized. The tumescent technique reported here is convenient, inexpensive and provides reliable control of intraoperative blood loss. There is a small body of literature concerned with the use of intramammary infiltration of adrenaline to reduce bloodloss. In 1974 Bretteville-Jensen (1) published a series

of 12 patients each of whom had one breast infiltrated with 80 to 140 mL of a solution containing 0.125% lignocaine and 1:800,000 noradrenaline. Intraoperative bloodloss was measured by weighing sponges, and postoperative drainage was recorded. He found a significant difference and advocated the technique. Brantner (2) published a similar study in 1984 in which 10 patients each had one breast injected with 50 mL of 0.5% lidocaine with 1:200,000 adrenaline and the other with an equal amount of normal saline. Intraoperative bloodloss on sponges and drapes was estimated and the results again showed a significant reduction in bloodloss on the side infiltrated with adrenaline. In 1992 Samdal et al (3) published a very similar study of 12 patients in whom one breast was injected with about 240 mL of 1:800,000 noradrenaline solution. Bloodloss was calculated by swab weighing and measurement of postoperative drainage. Results were again statistically significant. What Klein calls the tumescent technique for liposuction involves the injection of large volumes

Correspondence and reprints: Dr Michael Bell, 340 McLead Street, Suite 306, Ottawa, Ontario K2P 1A4. Telephone (613) 233-1815, Fax (613) 233-5532

TABLE 1: Controls

Patient	Age	Weight excised	Volume injected	HB (Pre-op)	HB (Post-op)	Δ HB
1	23	1100	137	126	11	1.00
2	22	1900	125	115	10	0.53
3	21	800	126	108	18	2.25
4	23	1050	135	114	21	2.00
5	29	725	127	95	32	4.41
6	39	800	137	131	6	0.75
7	25	900	120	103	17	1.89
8	56	3300	128	98	30	0.91
9	32	1025	120	99	21	2.05
10	18	1850	144	126	18	0.97
11	38	1750	139	121	18	1.03
12	24	1400	134	110	24	1.71
13	19	1600	142	130	12	0.75
14	39	1625	144	112	32	1.97
15	21	600	130	116	14	2.33
16	21	3500	135	110	25	0.71
17	53	1000	133	116	17	1.70
18	64	540	124	99	25	4.63
19	32	1075	130	113	17	1.58
20	16	1100	141	125	16	1.45
21	46	1750	143	109	34	1.94
22	27	1875	146	128	18	0.96
23	31	1925	131	118	13	0.68
Means	31	1443	134	114	20	1.66

Standard deviation 1.06; 95% confidence interval 0.43; #=FNG; HB Hemoglobin

of dilute local anaesthetic solution with adrenaline and sodium bicarbonate. He reports that this significantly reduces bloodloss from large volume liposuction (4).

Concerns regarding the use of catecholamine solutions have been raised following reports in 1963 of two cases of

cardiac arrest in association with halothane anaesthesia. Halothane is no longer commonly used and was not employed in any of our patients. A dose of 10 mL/kg lidocaine and 1:1,000,000 adrenaline, 10 mL/kg was employed. Patients experienced no symptoms referable to either agent and the anaesthetists reported no arrhythmias or episodes of tachycardia. This dose is in line with that used in facelift and rhinoplasty procedures and appears to be perfectly safe. Klein reports the safety of doses of up to 35 mg/kg of lidocaine (4). Some surgeons express fears that the use of vasoconstrictors may predispose to postoperative bleeding and hematoma formation. We did not find this to be a problem. Others express concern that nipple ischemia may result. A superior pedicle procedure was used with all but two patients (who had free nipple grafts) and no nipple loss was experienced in the entire series.

MATERIALS AND METHODS

Sixty-seven consecutive patients were included in the series of whom four were later excluded as no postoperative hemoglobin was available either because the patient was from out of town or because of clerical error. One patient was in the control group and three were in the 10 mg/kg group. Routine preoperative hemoglobin obtained either in the hospital preadmission unit or on admission was compared with hemoglobin drawn at a visit on the eighth postoperative day. We feel this gives a more accurate picture of the overall bloodloss than weighing swabs and estimating the amount of

Figure 1) Comparison of mean drop in hemoglobin per 10 g of breast tissue removed. Hatched areas represent 95% confidence limits either side of the mean

TABLE 2: Treatment - 250 mL/side

Patient	Age	Weight excised	Volume excised	HB (Pre-op)	HB (Post-op)	Δ HB	Δ HB/100 g excised
1	23	1800	250	129	116	13	0.72
2	22	2000	250	129	122	7	0.35
3	24	1900	250	131	104	27	1.42
4	34	1055	250	148	118	30	2.84
5	39	1500	250	131	125	6	0.40
6	22	2150	250	121	111	10	0.47
7	45	1550	250	138	125	13	0.84
8	37	1500	250	143	117	26	1.73
9	17	1350	250	137	111	26	1.93
10	22	1100	250	138	127	11	1.00
11	32	2400	250	137	112	25	1.04
12	34	600	250	123	118	5	0.83
13	56	1000	250	123	120	3	0.30
14	31	650	250	133	129	4	0.62
15	28	1700	250	131	117	14	0.82
16	24	1850	250	137	120	17	0.92
17	24	1300	250	136	131	5	0.38
18	34	1250	250	132	116	16	1.28
19	23	1200	250	134	120	14	1.17
20	29	950	250	125	112	13	1.37
Means	30	1440	250	133	119	14	1.02

Standard deviation 0.63; 95% Confidence interval=0.27; Compared to controls, $P=0.02$

TABLE 3: Treatment: 10 mg/kg bodyweight injection

Patient	Age	Weight excised	Volume injected	HB (Pre-op)	HB (Post-op)	Δ HB	Δ HB/100 g excised
1	23	1025	700	134	127	7	0.68
2	18	1700	0.71	800	134	122	12
3	31	3225	1300	138	112	26	0.81
4	29	1200	780	143	134	9	0.75
5	23	550	570	119	110	9	1.64
6	30	1225	720	142	134	8	0.65
7	32	700	*	131	120	11	1.57
8	40	1550	930	144	130	14	0.90
9	53	3050	1260	143	135	8	0.26
10	27	2025	820	134	124	10	0.49
11	29	1225	640	139	124	15	1.22
12	68	800	660	128	126	2	0.25
13	19	775	*	141	132	9	1.16
14	30	600	510	128	131	-3	0.50
15	48	1075	720	130	121	9	0.84
16	28	1250	680	125	119	6	0.48
17	23	1450	*	136	128	8	0.55
18	36	1100	640	148	137	11	1.00
19	29	1660	720	138	128	10	0.60
20	30	775	640	130	131	-1	-0.13
21	23	1025	*	137	12(0	11	1.07
Means	32	1333		135	126	9	0.72

Standard deviation 0.50; *Volume not recorded; 95% confidence interval 0.22; Compared to controls $P=0.000454$

blood on the drapes. The latter method is also fraught with problems when large volumes of solution are infiltrated, as fluid sponged up during the operation is quite watery. The same surgeon (MSGB), assisted by a resident, performed all the operations using a superior pedicle technique, except for two patients in the control group who required free nipple grafts. The solution was prepared by mixing five 20 mL bottles of 1% lidocaine with 1:100,000 in 1 L of normal saline. This yields a concentration of 0.1% lidocaine with 1:1,000,000 adrenaline. The incision lines were marked with Bonney's Blue and a small volume of solution was infiltrated subcutaneously with a 1.5 inch 21 gauge needle; the remaining solution was infiltrated along the planned incision lines and down to the pectoralis fascia with a 12 gauge blunt tipped Wells Johnson infusion cannula (#20-0008LL) fitted to a large syringe. Twenty patients were infiltrated with 500 mL (250 mL per breast) and 21 patients were injected with 10 mL/kg bodyweight (eg, a 70 kg woman would receive 350 mL per breast.) No drains were left in place and the incisions were dressed with Hypafix and pads. The pads were removed the morning following surgery and most patients went home that day. No problems were experienced with breast reshaping or symmetry despite the large volumes infused.

Tissue removed was weighed in the operating room and recorded. The change in hemoglobin was obtained by comparing the pre- and postoperative values, and the change in hemoglobin per 100 g of tissue removed was calculated. The values for each group of patients are recorded in Tables 1, 2 and 3. The standard deviations and 95% confidence intervals were calculated for each group and compared (Figure 1). Each of the two treatment groups was compared with the control group using Student's *t*-test with a two tailed distribution and two sample equal variance. There was no significant difference between the groups with respect to age or volume of breast removed.

RESULTS

The mean drop in hemoglobin in the control group was 20 mg/dL and the mean drop per 100 g of tissue resected was

1.66 mg/dL with 95% confidence limits of ± 0.43 . In the group treated with 250 mL infiltration per side the mean drop in hemoglobin was 14 mg/dL or 1.02 mg/dL for each 100 g of breast removed, with confidence limits of ± 0.27 mg/dL. The difference was statistically significant at the $P < 0.02$ level. In the group treated with 10 mL infiltration per kg of body weight the mean drop in hemoglobin was 9 mg/dL or 0.72 mg/dL for every 100 g of breast removed. The 95% confidence limits in this group were ± 0.22 and when compared with the control group the difference was significant ($P < 0.01$). There was no significant difference between the groups with regard to age or size of breast reduction.

DISCUSSION

Healthy women presenting for elective surgery should not be placed at risk of requiring homologous blood transfusions. Autologous transfusion is expensive and inconvenient but many surgeons performing reduction mammoplasty have regarded the acquisition of autologous blood as necessary in case of unexpectedly high blood loss during the procedure. We present a series of 64 patients, none of whom required transfusion, and a method of reducing intraoperative blood-loss to an average of 9 mg/dL. The tumescent technique of infiltrating breast tissue along the lines of resection with large volumes of extremely dilute local anaesthetic is a safe, cheap and startlingly effective method of reducing bloodloss during reduction mammoplasty. It also removes the inconvenience and cost of autologous donation while sparing the patient days lost from her usual activities caused by the lethargy of postoperative anemia.

REFERENCES

1. Bretteville-Jensen G. Mammoplasty with reduced blood loss: Effect of noradrenaline. *Br J Plastic Surg* 1974;27:31-4.
2. Brantner Lt-Col JN. The role of vasoconstrictors in control of blood loss in reduction mammoplasty. *Plast Reconstr Surg* 1985;75:339-41.
3. Samdal F, Serra M, Skollebor KC. The effects of infiltration with adrenaline on blood loss during reduction mammoplasty. *Scand J Plast Reconstr Surg* 1992;26:211-5.
4. Klein JA. Tumescent technique for local anaesthesia improves safety in large volume liposuction. *Plast Reconstr Surg* 1993;92:1085-100.