Tissue expansion for closure of the donor site of a distant pedicle flap

Ronald M Zuker MD FRCSC FACS, Craig Fielding MD FRCSC, Steve Beatty MD FRCSC
Division of Plastic Surgery, The Hospital for Sick Children, Toronto, Ontario

RM Zuker, C Fielding, S Beatty. Tissue expansion for closure of the donor site of a distant pedicle flap. Can J Plast Surg 1994; 2(4):155-158. Tissue expansion continues to provide new and exciting applications in reconstructive surgery. Tissue expansion has allowed for an increase in size of specialized tissue, particularly about the head and neck. It has also been demonstrated that the blood supply of pedicle flaps can be augmented, thus extending the surviving length of expanded flaps. Recently tissue expansion has been used with free flap transfer to expand the amount of tissue to be transferred and to facilitate donor site closure. In certain situations free tissue transfer may not be recommended and pedicle flap and particularly cross leg and cross thigh flaps are required. In addition to the prolonged unnatural positioning of the patient, the donor defect of these flaps has been harshly criticized. In this report tissue expansion is used to provide adequate tissue for direct closure of the donor defect thus minimizing the deformity.

Key Words: *Pedicle flap, Tissue expansion*

Expansion tissulaire pour la fermeture du site donneur au moyen d'un lambeau pédiculé à distance

RÉSUMÉ : L'expansion tissulaire continue d'offrir de nouvelles applications prometteuses en chirurgie reconstructive. Ce procédé a permis d'accroître la taille des tissus spécialisés, particulièrement au niveau de la tête et du cou. Il a en outre été démontré que l'apport sanguin au niveau des lambeaux pédiculés peut être accru, ce qui prolonge la survie des lambeaux. Récemment, ce procédé a été utilisé pour le transfert d'un lambeau libre qui a permis d'agrandir la portion de tissu à transférer et qui a facilité la fermeture du site donneur. Dans certaines situations, le transfert de tissu libre ne peut être recommandé et les lambeaux pédiculés, particulièrement les lambeaux de la jambe ou de la cuisse contralatérales sont nécessaires. En plus d'imposer au patient une posture qui n'est pas naturelle, le site donneur fait l'objet de critiques sévères. Dans le présent rapport, l'expansion tissulaire est utilisée afin de produire un segment de tissu suffisant pour une fermeture directe du site donneur, ce qui réduit les risques de déformation.

Tissue expansion continues to see new and innovative applications (1-6). Additional tissue may be provided utilizing this technique for conventional flap reconstructions. Recently tissue expansion of free flap donor sites has not only provided additional tissue for transfer but has also facilitated closure of the donor site (7,8). The cross leg and cross thigh flaps, although not commonly used, suffer from aesthetically unappealing residual donor site deformities. In this report we describe the use of tissue expansion at the time of pedicle flap elevation to facilitate donor site closure.

CASE HISTORY

A six-year-old boy sustained a major lawn mower injury to his left foot. He was initially resuscitated at a local hospital. The wound was superficially cleansed and, because of a strong family history of malignant hyperthermia, he was transferred by air ambulance to the major tertiary centre for adequate cleansing under general anaesthesia.

Shortly following arrival at The Hospital for Sick Children he was taken to the operating room where the wound was

thoroughly cleansed (Figure 1). There was a major injury to the posterior medial aspect of the foot with extensive soft tissue loss, as well as fracture of the underlying calcaneus and division of the Achilles tendon. At the initial procedure, following thorough debridement of necrotic tissue, the calcaneal fracture was reduced and held in place with K-wires, the Achilles tendon was repaired, and the exposed tendon itself was covered with a bipedical fasciocutaneous flap. The residual soft tissue defects were skin grafted (Figure 2).

In order to achieve definitive cover, a flap would be required. A free tissue transfer was considered, but after appropriate consultation with the anaesthesia staff it was felt that shorter procedures would be safer for this patient with a strong history of malignant hyperthermia. Five days following the injury he was returned to the operating room where plans were made for a cross thigh pedicle flap to resurface the 7 x 3 cm defect. This flap was designed as a superiorly based fasciocutaneous flap on the medial aspect of the right thigh. The donor defect could not be closed and was covered with Biobrane (Dow Hickam Pharmaceuticals, Texas). Through the anterior incision, a pocket was created on the anteromedial thigh which would house a 150 mL tissue expander (Figure 3). After the required defect on the foot was prepared, the fasciocutaneous flap was secured in place (Figure 4).

As the patient was from a considerable distance away, he

Correspondence and reprints: Dr Ronald M Zuker, The Hospital for Sick Children Head, Division of Plastic Surgery 1524-555 University Avenue, Toronto, Ontario M5G 1X8. Telephone (416) 598-1202, Fax (416) 813-6147.

Figure 1) Initial appearance of the wound

Figure 4) Flap secured in place

Figure 2) Appearance of the wound after initial debridement, repair and temporizing skin graft

Figure 5) Primary closure of the donor site

Figure 3) Appearance of the heel defect and outline of pedicle flap, tissue expansion for closure of donor site

Figure 6) Uneventful wound healing in the foot and in the thigh

remained hospitalized during the time of pedicle attachment. This allowed for injection of the expander on alternate days. After three weeks the expander was fully inflated and the pedicle flap was ready for division. He was returned to the operating room where the pedicle was divided and the flap

was inset in the foot. The tissue expander was removed as was the Biobrane over the donor site defect. With relative ease the donor site was closed primarily (Figure 5).

The postoperative course was uneventful both in the thigh and in the foot with primary healing taking place (Figure 6).

Figure 7) Normal range of motion in ankle and forefoot

Figure 8) Stable flap coverage

With gradual mobilization the patient returned to full activity after three months.

At two year follow-up the patient was taking part in all activities with no limitation. He had a full range of motion of his ankle and forefoot with normal strength in dorsiflexion and plantarflexion (Figure 7). He wears normal footwear and has had no breakdown problems (Figure 8). The flap exhibits protective and light touch sensation throughout. Imprint

Figure 9) Imprint analysis demonstrating weight bearing on the flap

Figure 10) Satisfactory long term donor site scarring

analysis demonstrates that the flap is taking the majority of the weight on the patient's left heel (Figure 9). The donor site on the right thigh remains satisfactory (Figure 10).

DISCUSSION

The continued use of tissue expansion provides reconstructive solutions for difficult problems. In addition to providing additional specialized tissue and augmenting the blood supply of traditional random flaps, tissue expansion also may play a role in improving donor site morbidity. This has been demonstrated in free tissue transfer to provide not only additional tissue for transfer, but also to facilitate donor flap closure. This same concept can be used for closure of the donor site of pedicle flaps with no additional surgical procedures required. The expanders are inserted in the tissue adjacent to the flap at the time of flap elevation. It is removed at the time of flap division. The expanded tissue can provide for primary donor site closure.

REFERENCES

1. Antonyshyn O, Gruss JS, Zuker RM, et al. Tissue expansion in head and neck reconstruction. *Plast Reconstr Surg* 1988;82:58.
 2. Cherry GW, Austad E, Pasyk K, et al. Increased survival and vascularity of random pattern skin flaps elevated in controlled expanded skin. *Plast Reconstr Surg* 1983;72:680.
 3. Hallock GG. Refinement of radio forearm flap donor site using skin expansion. *Plast Reconstr Surg* 1988;81:21.
 4. Leighton WD, Russel RC, Marcus DE, et al. Experimental free transfer expansion of free flap donor sites. 1. Flap viability and expansion characteristics. 2. Physiology, histology and clinical correlation. *Plast Reconstr Surg* 1988;82:69.
 5. Manders EK, Schenden MJ, Furrey JA, et al. Soft tissue expansion: Concepts and complications. *Plast Reconstr Surg* 1984;74:493.
 6. Radovan C. Breast reconstruction after mastectomy using a temporary tissue expander. *Plast Reconstr Surg* 1982;69:195.
 7. Sasaki GH, Pang CY. Pathophysiology of skin flaps raised on expanded pig skin. *Plast Reconstr Surg* 1984;74:59.
 8. Zuker RM. The use of tissue expansion in pediatric scalp burn reconstruction. *J Burn Care Rehab* 1987;103:106.
-