Evidence-based medicine: The *Canadian Journal of Plastic Surgery's* future

Over the past few decades, evidence-based medicine has been rated as one of the most significant medical advances in patient care (1). The ability to systematically rate research evidence should direct patient care so that the most beneficial interventions for patients are implemented. In the absence of intervention evaluation, significant portions of our health care dollars continue to be wasted on care that does not benefit the patient or has never been appraised for its supporting evidence. In the Spring 2012 issue of the *Canadian Journal of Plastic Surgery* (CJPS), the article by Thoma et al (2) provided a brief history and detailed the importance of evidence-based medicine. Currently, more and more scientific journals are focusing on evidence-based medicine and level of evidence (LOE) rating scales when evaluating journal articles. These scales enable researchers and authors to produce better articles, and enable readers to easily understand the LOE for each published article.

There are many LOE rating scales in the literature. The CJPS has decided to use the rating system and standardized icons currently used by *Plastic and Reconstructive Surgery*. The editorial by Sullivan et al (3) highlights the importance of evidence-based medicine and the need for journals to embrace LOE rating scales. The LOE tables to be implemented by the CJPS were developed by the American Society of Plastic Surgeons and are presented here. It is with their full support and encouragement that we begin duplicating their requirements for authors to rate their articles based on the LOE in one of three categories: THERAPEUTIC, DIAGNOSTIC or PROGNOSTIC/RISK. Each article will have its level rated based on the set criteria established by the American Society of Plastic Surgeons. The CJPS's Editorial Board will then confirm the level and publish the level with the article. The same triangular rating logo will be used in a similar manner to what is currently published in *Plastic and Reconstructive Surgery* (Figure 1).

Hopefully, this new initiative will enhance the quality of the work published in the CJPS and enhance the educational experience of our readers.

Sincerely,
Edward W Buchel MD
Editor-in-Chief
Canadian Journal of Plastic Surgery

Figure 1 The level of evidence pyramid, showing level of evidence (I through V) and clinical question addressed by the article. **Left** Diagnostic clinical question addressed, with a level of evidence of II. **Centre** Therapeutic clinical question addressed, with a level of evidence of III. **Right** Risk clinical question addressed, with a level of evidence of II. Reproduced with permission

Evidence rating scale for therapeutic studies

Level of evidence	Qualifying studies
I	High-quality, multicentred or single-centred, randomized controlled trial with adequate power; or systematic review of these studies
II	Lesser-quality, randomized controlled trial; prospective cohort or comparative study; or systematic review of these studies
III	Retrospective cohort or comparative study; case-control study; or systematic review of these studies
IV	Case series with pre/post test; or only post test
V	Expert opinion developed via consensus process; case report or clinical example; or evidence based on physiology, bench research or 'first principles'

Evidence rating scale for diagnostic studies

Level of evidence	Qualifying studies
I	High-quality, multicentred or single-centred, cohort study validating a diagnostic test (with 'gold' standard as reference) in a series of consecutive patients; or a systematic review of these studies
II	Exploratory cohort study developing diagnostic criteria (with 'gold' standard as reference) in a series of consecutive patient; or a systematic review of these studies
III	Diagnostic study in nonconsecutive patients (without consistently applied 'gold standard' as reference); or a systematic review of these studies
IV	Case-control study; or any of the above diagnostic studies in the absence of a universally accepted 'gold' standard
V	Expert opinion developed via consensus process; case report or clinical example; or evidence based on physiology, bench research or 'first principles'

Evidence rating scale for prognostic/risk studies

Level of evidence	Qualifying studies
I	High-quality, multicentred or single-centred, prospective cohort or comparative study with adequate power; or a systematic review of these studies
II	Lesser-quality prospective cohort or comparative study; retrospective cohort or comparative study; untreated controls from a randomized controlled trial; or a systematic review of these studies
III	Case-control study; or systematic review of these studies
IV	Case series with pre/post test; or only post test
V	Expert opinion developed via consensus process; case report or clinical example; or evidence based on physiology, bench research or 'first principles'

REFERENCES

- Godlee F. Milestones on the long road to knowledge. *BMJ* 2007;334(Suppl 1):S2-3.
- Thoma A, Ignacy TA, Li YK, Coroneous CJ. Reporting the level of evidence in the *Canadian Journal of Plastic Surgery*: Why is it important? *Can J Plast Surg* 2012;20:12-6.
- Sullivan D, Chung KC, Eaves FF III, Rohrich RJ. The level of evidence pyramid: Indicating levels of evidence in *Plastic and Reconstructive Surgery* articles. *Plast Reconstr Surg* 2011;128:311-4.