Corrigendum to "On the Diophantine Equation $x^2 = 4q^n - 4q^m + 9$ "

Amal Al-Rashed AND Fadwa S. Abu Muriefah

*Mathematical Sciences Department, College of Science, Princess Nourah Bint Abdulrahman
University*

aaalrashed@pnu.edu.sa , fsabumuryifah@pnu.edu.sa & abu.muriefah@yahoo.com

Abstract

This article is a corrigendum to Theorem 1, of our paper ^[1] on the solution of the Diophantine equation $x^2 = 4q^n - 4q^m + 9$.

Key Words

Diophantine equation, Lucas sequence, primitive divisor.

I. Introduction

The Diophantine equation

$$x^2 = 4q^n - 4q^m + D^2, \quad (1)$$

where x, D, q are positive integers, D is any odd integer and q is any prime, is a generalization of Ramanujan equation ($m = 1, q = 2, D = 1$), which has the form

$$x^2 = 2^{n+2} - 7.$$

He conjectured in 1913 that this equation had exactly five solutions in integers, given by

$$(n, x) = (1, 1), (2, 3), (3, 5), (5, 11), (13, 181).$$

For the history of this conjecture one can see ^[2,3]. The case $D = 1$, in Eq. (1) is solved completely by Luca ^[4]. In this paper we suppose $D = 3$, and we solve the diophantine equation

$$x^2 = 4q^n - 4q^m + 9, \quad n > m > 0 \quad (2)$$

in positive integers and q a prime number.

II. Preliminaries

In this section we introduce some useful results. We start by the following definition.

Definition ^[5]

A **Luca pair** is a pair (α, β) of algebraic integers such that $(\alpha + \beta)$ and $\alpha\beta$ are non-zero coprime rational integers and α/β is not a root of unity.

Given a Luca pair (α, β) , the corresponding **Lucas numbers** are defined by

$$u_n(\alpha, \beta) = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n = 0, 1, 2, \dots$$

A prime number p is a **primitive divisor** of $u_n(\alpha, \beta)$ if p divides $u_n(\alpha, \beta)$, but does not divide $(\alpha - \beta)^2 u_1 u_2 \dots u_{n-1}$.

For any integer $n > 30$, the n -th term of any Lucas sequence has a primitive divisor. Further, for any positive integer $n \leq 30$, all Lucas pairs which have no primitive divisor are determined in Table 1 and Table 3 ^[5].

III. Main Result

In this paper we correct the proof of Theorem (1) in our paper ^[1].

If $(n, m) = (1, 0)$, we get the equation $x^2 = 4q + 5$, which has a solution $q = 5$, $x = 5$. Also, if $d = \gcd(n, m)$, suppose $m = dm_1$ and $n = dn_1$, where $\gcd(m_1, n_1) = 1$, then rewrite Eq. (2), as

$$\begin{aligned} x^2 &= 4q^{dn_1} - 4q^{dm_1} + 9 \\ &= 4q_1^{n_1} - 4q_1^{m_1} + 9, \end{aligned}$$

where $q_1 = q^d$. This equation on q_1 has the same type as Eq. (2), where the exponents are coprime. So, we may assume that m and n are coprime and q is a prime power.

Theorem

For $m = 1$, The diophantine equation (2) has at most two solutions (x, n) ; while for $m > 1$, the only solutions of diophantine equation (2) with q a prime power are

$$\begin{aligned} &(x, n, m, q) \\ &= (93, 7, 3, 3), (2 \times 3^{M+1} - 3, 2M + 2, M \\ &\quad + 2, 3), (9, 3, 2, 3), \\ &(5, 3, 2, 2), (11, 5, 2, 2), (181, 13, 2, 2). \end{aligned}$$

for any $M \in \mathbb{N}$.

Proof

i. $m = 1$

Put $q = 2$, in Eq (2), we get $x^2 - 1 = 2^{n+2}$, which has the solution $(x, n) = (3, 1)$. If $q = 3$, then Eq. (2) becomes $x^2 = 4 \times 3^n -$

3, which is impossible modulo 9, unless $n = 1$ which is impossible since $n > m$. Otherwise the equation $x^2 + 4q - 9 = 4q^n$ has at most two solutions (x, n) ^[6].

ii. $m > 1$

Put $q = 3$, in Eq. (2), we get

$$y^2 = 4 \times 3^{n-2} - 4 \times 3^{m-2} + 1, \quad (3)$$

where $y = x/3$. If $(n-2, m-2) \neq (1,0)$, the solution of Eq. (3) were found by Luca ^[4], as follows:

- $(y, n, m) = (31, 7, 3)$ if $n-2 \neq 2(m-2)$,
- $(y, n, m) = (2 \times 3^M - 1, 2M+2, M+2)$ with $M \in \mathbb{N}$, if $n-2 = 2(m-2)$.

Finally, when $(n-2, m-2) = (1,0)$, then $y = 3$. Consequently, when $q = 3$, Eq. (2) has the following solutions:

$$(x, n, m) = (93, 7, 3), (2 \times 3^{M+1} - 3, 2M+2, M+2), (9, 3, 2) \text{ with } M \in \mathbb{N}.$$

Thus, from now on we assume that $q \neq 3$. Rewrite Eq. (2), as

$$4q^m - 9 = 4q^n - x^2 = Aa^2 \quad (4)$$

where A is an odd square free number. It is clear that $A \neq 1$, because -1 is a quadratic nonresidue modulo 4. Also $\gcd(3, A) =$

$\gcd(3, a) = 1$, if so then $q = 3$. Hence $A \geq 5$. Let $q = p^f$, where p is a prime and $f \geq 1$. Since $A \equiv 3 \pmod{4}$ and $q > 3$. It is easy to prove that p splits in $\mathbb{Q}[\sqrt{-A}]$. Write $\langle p \rangle = \pi \bar{\pi}$, where π is a prime ideal in the field $\mathbb{Q}[\sqrt{-A}]$. From (4), we get

$$p^{fm} = q^m = \left(\frac{3 + a\sqrt{-A}}{2} \right) \left(\frac{3 - a\sqrt{-A}}{2} \right), \quad (5)$$

which implies that

$$\pi^{fm} \bar{\pi}^{fm} = \left\langle \frac{3 + a\sqrt{-A}}{2} \right\rangle \left\langle \frac{3 - a\sqrt{-A}}{2} \right\rangle. \quad (6)$$

It is easy to prove that the two ideals in the right-hand side of (6) are coprime. Hence from the unique factorization property for ideals, it follows that

$$\pi^{fm} = \left\langle \frac{3 + a\sqrt{-A}}{2} \right\rangle. \quad (7)$$

Since π^{fm} is principal, therefore the order of the ideal class of π divides fm .

Now from (4), we also have

$$q^n = \left(\frac{x + a\sqrt{-A}}{2} \right) \left(\frac{x - a\sqrt{-A}}{2} \right), \quad (8)$$

which implies that

$$\pi^{fn} \bar{\pi}^{fn} = \left\langle \frac{x + a\sqrt{-A}}{2} \right\rangle \left\langle \frac{x - a\sqrt{-A}}{2} \right\rangle \quad (9)$$

It is easy to prove that the two ideals appear in the right-hand side of (9) are coprime.

Hence

$$\pi^{fn} = \left\langle \frac{x + a\sqrt{-A}}{2} \right\rangle$$

Since π^{fn} is principal, therefore the order of the ideal class of π divides fn . But m and n are coprime, so the order of the ideal class of π divides f . Hence, π^f is principal. Let $z = \frac{c+b\sqrt{-A}}{2}$, where $c \equiv b \pmod{2}$, be a generator of π^f , then we have

$$\langle q \rangle = \langle p^f \rangle = \pi^f \bar{\pi}^f = \langle z \rangle \cdot \langle \bar{z} \rangle,$$

therefore, from (6), we get

$$\langle z^m \rangle \cdot \langle \bar{z}^m \rangle = \langle q^m \rangle = \left\langle \frac{3+a\sqrt{-A}}{2} \right\rangle \left\langle \frac{3-a\sqrt{-A}}{2} \right\rangle. \quad (10)$$

The two ideals in the right-hand sides of (10) are coprime. It follows that we may assume

$$\langle z^m \rangle = \left\langle \frac{3+a\sqrt{-A}}{2} \right\rangle \text{ and } \langle \bar{z}^m \rangle = \left\langle \frac{3-a\sqrt{-A}}{2} \right\rangle.$$

Since the units in the field $\mathbb{Q}(\sqrt{-A})$ are ± 1 , therefore

$$\pm z^m = \frac{3+a\sqrt{-A}}{2} \text{ and } \pm \bar{z}^m = \frac{3-a\sqrt{-A}}{2} \quad (11)$$

Hence

$$\pm 3 = z^m + \bar{z}^m = \frac{z^{2m} - \bar{z}^{2m}}{z^m - \bar{z}^m} = \frac{u_{2m}}{u_m} \quad (12)$$

where $\{u_m\}$ denotes the Lucas sequence corresponding to $(z, \bar{z})$. Using the value of z in (12), we get

$$\pm 3 \times 2^m = (c + b\sqrt{-A})^m + (c - b\sqrt{-A})^m$$

If c and b are both even, then 2^{m+1} divides the left-hand side which is not true, so c and b are both odd. It is clear that $\gcd(c, b) = \gcd(c, A) = 1$ and so $\gcd(c^2, Ab^2) = 1$. Therefore u_{2m} is a Lucas number which has no primitive divisors.

We start with $m = 2$, then substitute z in (12), we get

$$z^2 + \bar{z}^2 = \frac{c^2 - Ab^2}{2} = \pm 3 \quad (13)$$

Now from (11) and the value of z , we have

$$\begin{aligned} \frac{3 + a\sqrt{-A}}{2} &= \pm z^2 \\ &= \pm \left(\frac{c^2 - Ab^2}{4} + \frac{cb\sqrt{-A}}{2} \right), \end{aligned}$$

so $a = \pm cb$. On the other hand, Eq. (8) can be written in this form

$$z^n \bar{z}^n = q^n = \left(\frac{x + a\sqrt{-A}}{2} \right) \left(\frac{x - a\sqrt{-A}}{2} \right).$$

Since $q \neq 3$, as above we can conclude

$$\frac{x+a\sqrt{-A}}{2} = \pm z^n \text{ and } \frac{x-a\sqrt{-A}}{2} = \pm \bar{z}^n,$$

which implies

$$z^n - \bar{z}^n = \pm a\sqrt{-A} = \pm cb\sqrt{-A},$$

hence

$$\begin{aligned} \pm cb &= \frac{b}{2^{n-1}} (nc^{n-1} + \dots \\ &\quad + (-1)^{(n-1)/2} A^{(n-1)/2} b^{n-1}) \end{aligned}$$

This implies $c|A^{(n-1)/2}b^{n-1}$, but $\gcd(c, Ab) = 1$, so $c = \pm 1$. Substitute in (13), we get $A = 7, b = 1$, hence $q = 2$, and then from (4), we get

$$x^2 = 2^{n+2} - 7. \quad (14)$$

Eq. (14) has the following solutions^[3]

$$(x, n) = (5, 3), (11, 5), (181, 13).$$

Now suppose $m = 3$, in (11), then we have

$$\pm 12 = c(c^2 - 3Ab^2). \quad (15)$$

This implies that $c = \pm 1, \pm 3$. Again from (15), this is impossible.

Finally, for other values of m , we have $2m \geq 8$. Since u_{2m} has no primitive divisor, then from Table 2 and 3 in^[4] we conclude that Eq. (2) has no solutions.

◆References

- [1] **F. S. Abu Muriefah, and Amal AL-Rashed**, On the diophantine equation $x^2 = 4q^n - 4q^m + 9$. J. KAU: Sci., 21 (1), (2009), 135-141.
- [2] **L. J. Mardell**, diophantine equations, Academic Press, New York (1969), 205-206
- [3] **T. Nagell**, The diophantine equation $x^2 - 7 = 2^{n+2}$, Nordisk. Mat. Tidsker, 30, 62-64, Ark. Mat. (1960), 185-187.
- [4] **F. Luca**, On the diophantine equation $x^2 = 4q^n - 4q^m + 1$, Proc. Amer. Math. Soc. 131(2002), 1339-1345.
- [5] **Y. Bilu, G. Hanrot and P.M. Voutier** "Existence of primitive divisors of Lucas and Lehmer numbers" J. Reine Angew. Math., 539(2001), 75-122.
- [6] **L. Maohua**, On the diophantine equation $x^2 + D = 4y^n$, J. Number Theory, 41(1992), 87-97.

$$x^2 = 4q^n - 4q^m + 9 \text{ تصويب معادلة ديوفانتين}$$

أمل الرشيد و فدوى أبومريفة

قسم العلوم الرياضية - كلية العلوم

جامعة الأميرة نورا بنت عبد الرحمن

المستخلص. في هذا البحث قدمنا تصحيح لبرهان المبرهنة 1، الواردة في المرجع (1) حول حل المعادلة الديوفانتية

$$x^2 = 4q^n - 4q^m + 9$$