The use of bioinformatics tools to characterize a hypothetical protein from *Penicillium rubens*

F.F. Silva, D.B. Gonçalves and D.O. Lopes

Universidade Federal de São João del Rei, Divinópolis, MG, Brasil

Corresponding author: D.O. Lopes

E-mail: debora@ufsj.edu.br

Genet. Mol. Res. 19 (2): gmr18574

Received February 16, 2020

Accepted May 23, 2020

Published June 30, 2020

DOI <http://dx.doi.org/10.4238/gmr18574>

ABSTRACT. Advances in biotechnology and molecular biology fields have allowed the sequencing of species genomes to improve the understanding of their genetics. As a consequence of the large number of whole genome sequencing projects, various DNA and protein sequences predicted by bioinformatics still without an identified function and are annotated in data banks as hypothetical proteins (HPs). Approximately 30% of the genome of recently sequenced organisms is composed of HPs, and these proteins are not included in functional studies due to the inexistence of full annotation. The characterization of HPs through bioinformatics is an important step to improve the understanding of genes and proteins and aims to assign some information to these sequences. Here, we used bioinformatics tools to predict the function of a hypothetical protein using as a model a protein from the filamentous fungus *Penicillium rubens* (XP_002569027.1). This fungus has been extensively studied due its ability to produce a wide range of natural products, many of them with biotechnological and pharmaceutical applications. The first step was to characterize the physicochemical properties of this HP. This protein has a molecular weight of 79.4 kDa and a theoretical pI of 5.06; 44.1% of its sequence is composed of hydrophilic amino acid residues and it was predicted as an extracellular protein. Prediction of 3D structure showed 42.21% of this sequence with an alpha galactosidase from *Geobacillus stearothermophilus*. In addition, XP_002569027.1 showed the presence of a glycosylase domain and most important amino acids residues in a catalytic site. Thus, we characterized

a HP from *P. rubens* using computational tools and were able to produce useful information for future *in vivo* and *in vitro* studies, highlighting the importance of bioinformatics as a preliminary tool for functional studies.

Key words: Bioinformatics; Hypothetical Proteins; *Penicillium rubens*; Functional prediction; Alpha-galactosidase

INTRODUCTION

After the idealization and realization of the Human Genome Project, a massive amount of data was generated and a lot of information was revealed to the scientific world (Lander et al., 2001; Schmutz et al., 2004). This was a milestone in scientific history, which is called the Genomic Era. Genomics deals with not just genes, but the whole genetic composition of a particular organism; with the dramatic improvement of sequencing technologies and the impressive reduction in costs, scientists are facing a “data avalanche” (Papageorgiou et al., 2018).

Sequencing projects started with small genomes and simple organisms, such as the bacteriophage MS2, which was the first organism that had its genome elucidated in 1976 (Fiers et al., 1976). As the techniques improved, it became possible to sequence genomes of more complex organisms, such as the human genome. Currently, there are over 50,000 genomes from eukaryotes, viruses, plasmids and organelles deposited in the National Center for Biotechnology Information (NCBI) genome database (<https://www.ncbi.nlm.nih.gov/genome/>).

The genomic era was driven by scientific interests and a tremendous expectation of new opportunities created by availability of genetic information, since the data produced by genome sequencing projects could be used by scientists around the world for different purposes and in different areas. The large amount of data produced through genome sequencing demanded and pushed the improvement of an important area called bioinformatics (Hesper and Hogeweg, 1970). This new science field put together two important areas: informatics with algorithms and computer software and molecular biology. This new field started to solve biological problems as the first demand emerged from genomic era: to assemble the genome puzzle and process all genomic information with quality and curation (Poliakov et al., 2015; Ge and Guo, 2020). By definition, bioinformatics is a science that uses computational tools to analyze genetic material and information about the function of genes and proteins in organisms. Nowadays, this interdisciplinary science encompasses various areas such as computing, mathematics, physics and biology, with the primary objective to answer biological questions (Bayat, 2002; Srinivasa et al., 2020).

The availability of genome sequences of a huge number of organisms associated to knowledge about the original central dogma of molecular biology (DNA, RNA, Proteins) pushed the appearing of two other important eras: proteomic and transcriptomic. Proteomics came with large-scale analyses of all expressed proteins and their isoforms present in a specific cell or tissue (Mallick and Kuster, 2010) and, in the same way, transcriptomic came with analyses of RNA transcripts (Lowe et al., 2017). From this time, the term “omic” started to become popular and took part of all genomic derivative products studies, as metabolomics, lipidomics, epigenomics and interactomics.

All “omic” studies use genes or proteins sequences from public databanks and sometimes face with sequences that have not been characterized and did not have functions attributed to them (Bernstein et al., 1977). These sequences are annotated in databases as unknown or hypothetical genes (HGs) or proteins (HPs), since they were predicted by computational tools, but there is no experimental evidence of its function (Mishra et al., 2013).

Determination of gene and its encoded protein function is one of the challenging problems coped by the post-genome era and demands from bioinformatics to predict functions of these sequences by developing efficient tools (Sivashankari and Shanmughavel, 2006). The sequencing of the complete genome of various organisms has been carried out for a long time, and yet a large part of the genome is still composed by proteins classified as hypothetical. Currently, it is estimated that 30% of all newly sequenced genes of microorganisms ended up annotated as “hypothetical” (Naveed et al., 2018). The bacteria *Mycobacterium tuberculosis*, for example, had its entire genome published in 1998 (Cole et al., 1998) and recent studies show that 27% of genes in *M. tuberculosis* genome are still classified as HPs, and there only few studies relating these proteins to some function (Yang et al., 2019).

Elucidate proteins function is an important step to understand the organism genetics, and the relevance of its characterization lies on the potential of application of this macromolecule in different studies. Nowadays, *in silico* functional studies have an important role in protein function prediction linking a hypothetical protein to one or many cellular processes and helping to elucidate its role in metabolic pathways (Ijaq et al., 2019). HPs characterization has become an attractive subject of study due to its potential importance, leading to creation of some databases where only HPs sequences are described, such as *HypoDB* (Adinarayana et al., 2011).

The scenario of HPs studies in filamentous fungi is no different. It is estimated that there are 1.5 to 5 million species of fungi on earth (Hawksworth and Lücking, 2017) and the vast majority of these organisms has not yet been sequenced and characterized (Uranga et al., 2017). Filamentous fungi have a great ability to secrete substances to the culture medium, such as secondary metabolites and proteins, which is due to the refined secretion machinery that these organisms have. Proteins secreted by filamentous fungi are related to broad range of metabolic processes in these organisms, like nutrition, resistance and pathogenicity (Peberdy, 1994). Such secretion ability is an attraction for commercial production of extracellular proteins in these organisms, functioning as biofactories through heterologous protein production systems. Fungi of the genus *Aspergillus* (*A. niger*, *A. oryzae*) and *Penicillium* (*P. camemberti*, *P. rubens*, *P. glaucum*) are the main representatives of this case (Punt et al., 2002).

The genus *Penicillium* has been extensively studied after the discovery of penicillin by Alexander Fleming. Filamentous fungi of this genus are used in several processes in the food industry, such as cheese, bread and brewery production, for example. With the scientific and biotechnological development, the emergence of new techniques has also made possible to use these filamentous fungi to produce a range of secondary metabolites and proteins of industrial interest (Meena et al., 2018). Specifically, the filamentous fungus *Penicillium rubens*, formerly *P. chrysogenum* and *P. notatum* (Houbraken et al., 2011), is widely used for the production of enzymes (Guzman-chavex et al., 2018), such as proteases (Haq et al., 2005), lipases (Ferrer et al., 2000) and mainly glycosylases, such as invertases

(Yakimova et al., 2017), amylases (Dar et al., 2015), galactosidases (Aleksieva et al., 2010) and xylanases (Chinedu et al., 2008). Although it is an organism widely used for biotechnological purposes, a huge number of its proteins are still annotated as HPs, reinforcing the importance of an initial *in silico* characterization to define its functions. Moreover, few studies related to proteomics and transcriptomics of filamentous fungus have been done (Kim et al., 2007), and the number is even lower for *P. rubens*, which had its complete genome published by Van Den Berg et al in 2008. Most studies with this organism are related only to characterization at an expression / excretion level and not at the genomic level, highlighting the importance of genomic studies in *P. rubens* (Jami et al., 2010; Pohl et al., 2020).

Given the importance of study of HPs few genomic studies have been made of the filamentous fungus *P. rubens*, the goal of this study was to perform, through bioinformatics analysis, the prediction and characterization of the HP XP_002569027.1.

MATERIAL AND METHODS

Nucleotide and amino acid sequences

The nucleotide and protein sequences were retrieved from the complete genome of *P. rubens* Wisconsin available at GenBank (<http://www.ncbi.nlm.nih.gov/genbank/>) under the access number AM920437.1. The sequence XP_002569027.1 was randomly chosen using the term “hypothetical”.

Search for signal peptide

The signal peptide prediction of the HP was performed using the *SignalP 5.0* (<http://www.cbs.dtu.dk/services/SignalP/>). This tool provides the probability the analyzed sequence has a signal peptide when compared to known signal peptides. In addition, this tool provides the cleavage sites of signal peptides and differentiate the types of signal peptide in *Archaea*, gram-positive or gram-negative bacteria and *Eukarya* (Almagro et al., 2019).

Physicochemical characterization

The physicochemical analyses of the HP were performed using two tools: *ProtParam* (<https://web.expasy.org/protparam/>) and *ProtScale* (<https://web.expasy.org/protscale/>). *ProtParam* allows the calculation of molecular weight, theoretical isoelectric point, amino acid composition, atomic composition, estimated half-life, hydropathicity index, among others. *ProtScale* also provides hydropathicity analyses, which is the ratio of hydrophobic and hydrophilic amino acids, where a value < 0 indicates a greater number of hydrophilic amino acids and consequently a protein that tends to be soluble in water, and a value > 0 indicates greater presence of hydrophobic amino acids, indicating water-insoluble protein (Gasteiger et al., 2005).

Structural analyses

The first structural tool used, called *TMHMM*, is based on the *Hidden Markov model* (HMM) for predicting probability that the analyzed sequence presents regions that

have characteristics of transmembrane helices (<http://www.cbs.dtu.dk/services/TMHMM/>), (Krogh et al., 2001). To predict secondary structures of the HP, we used *GOR* (https://npsa-prabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=/NPSA/npsa_gor4.html), a tool that indicates the probability of the presence of alpha helices, beta sheets, turns, random coils, among others (Garnier et al., 1996). The last structural analysis tool used was the *SWISS-MODEL* (<https://swissmodel.expasy.org>). This tool works through the 3D construction of the protein, in this case the HP, comparing with 3D structures already deposited in the database. The tool automatically selects, among the best candidates, templates to create the prediction of the possible 3D structure of the HP (Waterhouse et al., 2018).

Cellular localization prediction

Cellular localization prediction of the HP was performed using *TargetP* (<http://www.cbs.dtu.dk/services/TargetP/>). This tool search, in its database, similarity with specific regions that indicate cellular localization of proteins, such as signal peptides and mitochondrial targeting peptides, for example, presenting the probability of the analyzed sequence to have any of these specific regions (Almagro et al., 2019).

Search for similarity through local alignment

For the similarity analysis, we used the *Basic Local Alignment Search Tool* (*BLASTp*) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). This tool searches for regions of local similarity between the desired sequence with other sequences already deposited in the database and calculates the statistical significance of the correspondences (Altschul et al., 1990).

Characterization of functional domains and catalytic residues

The search for functional domains and important regions of the HP was performed using *InterProScan* (<http://www.ebi.ac.uk/interpro/>). This tool compares the analyzed sequence with proteins already described in its database, to provide information about conserved patterns, families, important domains and catalytic residues (Jones et al., 2014).

Global alignment

Five of the 10 sequences that showed high similarity (higher than 75%) with the selected HP in previous analyses (*BLASTp*, item 2.6) were submitted to *MultAlin* (<http://multalin.toulouse.inra.fr/multalin/>), to align and compare the sequences. Regions with similarity between the analyzed sequences and important residues (Corpet, 1988) were sought.

RESULTS AND DISCUSSION

Selection of a Hypothetical protein from *P. rubens*

The abundance of hypothetical genes and proteins is an important issue coming from Pos-Genomic and Pos-Proteomic Eras. Currently, databases are far from complete and errors in sequences in public databanks are expected, keeping in mind how complex is the

whole process, starting with genome sequencing, data assembly, annotation and curation of sequences (De Simone et al., 2019). Consequently, there is a vast number of HPs waiting to be functionally characterized. Therefore, the initial characterization of HPs through bioinformatics tools will help to speed up these studies, as they can gather information on HPs to guide functional studies. This strategy can help to assign proteins a tentative function promoting an important connection between *in silico* analysis with experimental trials (Lubec et al., 2005).

According to the Genome platform (www.ncbi.nlm.nih.gov/genome), *P. rubens* has a genome of 32.2 Mb, with 12,791 proteins, of which 2,158 are classified as HPs or partially hypothetical, representing approximately 17% of total proteins. The hypothetical gene (Pc21g20400) and its protein XP_002569027.1 from *P. rubens* were randomly chosen from databank to perform its bioinformatic characterization. The selected gene has 2241bp and encodes a protein with 746 amino acids.

Identification of the signal peptide

A signal peptide is a sequence of 16-30 amino acids usually located in the N-terminal region of the proteins and functions as a signal that the protein will be exported to different compartments from which it was synthesized, such as organelles or extracellular, and it is usually cleaved after protein transport. Alternatively, a signal peptide can be located inside the protein and not be cleaved after translocation (Kapp et al., 2009).

SignalP analysis indicates that there is 89.43% chance of HP XP_002569027.1 having an N-terminal signal peptide and that this signal peptide consists of the amino acids 1 to 20, presenting a cleavage site between the amino acids 24-25 (Figure 1), suggesting that HP XP_002569027.1 will be exported. After confirming the presence of a signal peptide, the first 24 amino acids were removed from the sequence for further analysis, leaving a sequence of 722 amino acids. The signal peptide is usually removed in bioinformatics analysis because it is not a sequence that is part of the protein after post-translational processing and can interfere in the following analyses (Choo et al., 2009).

Figure 1. Search for the presence of the signal peptide in hypothetical protein XP_002568981. Analysis done by *SignalP* tool, where (—) indicates probability of finding a signal peptide; (---) probability to be the cleavage site for the signal peptide; (—) probability to not contain any signal peptide.

Physicochemical characterization of HP XP_002569027.1

The physical-chemical analyses show that HP XP_002569027.1 has an estimated molecular weight of 79.4 kDa, a theoretical pI of 5.06, presenting 86 and 59 negatively and positively charged amino acids residues, respectively. The *ProtParam* tool also predicts the probable protein half-life time in different cell types, which was 1.3 hours in *in vitro* mammal reticulocytes, 3 min in yeast and 3 min in *E. coli*. In addition, it was possible to observe an instability index of 32.39, where proteins whose instability index are smaller than 40 are predicted as stable proteins (Guruprasad et al., 1991), indicating that the HP analyzed is a stable protein in test tubes.

The importance of analyzing the physical-chemical parameters of a HP is found in the transition from *in silico* to *in vitro* studies. For example, when analyzing the isoelectric point of a protein it is possible to know the pH at which the net charge of protein is zero, and this analysis is useful on purification methods, such as purification by isoelectric focusing. When knowing the theoretical pI of this HP, *in vitro* experiments that require pH control to be carried out can be done using a 5.06 isoelectric point as a start and control for *in vitro* experiments (Mohan and Venugopal, 2012).

According to the results of the *ProtScale* tool, which provides hydropathicity, HP XP_002569027.1 has a hydrophilic attribute, since a greater number of amino acids is depicted below the red line, with a value < 0 (Figure 2). The result of the hydropathicity from *ProtScale* is similar to that generated by *ProtParam*, which provided a hydropathicity index of -0.227 and has a hydrophilic character. Knowledge about the physicochemical characteristics of HPs is of great importance to produce this protein to be assayed in *in vitro* and to plan future experiments (Skolnick et al., 2000).

Figure 2. Hydropathicity analysis of hypothetical protein XP_002569027.1. Amino acids that are below the red line (value < 0) represent hydrophilic amino acids. Results generated by the ProtScale platform.

Production of a 3-D model of HP XP_002569027.1 using comparative Modeling

The transmembrane helices prediction was performed using *TMHMM* tool. The plot illustrated in Figure 3 indicates the probability of portions of sequence to be intracellular, extracellular or transmembrane, with results ranging from 0 to 1. The line between 1 and 1.2 is an overall line that sums the analysis. In this case, it is shown that the whole sequence (722 aa) is predicted to be outside the cell (thin pink line), and the overall analyses confirms it (thick pink line). According to this results, it is possible to infer that the HP XP_002569027.1 is not a transmembrane protein, and that most of the protein is more likely to be outside of the membrane, being a possible indication of extracellular protein, which agrees with the presence of a signal peptide (Figure 1), likely to be a signal for extracellular export. Although, according to *TMHMM* creators, this tool is not used as a confirmation of cellular location, its only predicts if the analyzed protein contain transmembrane helices and if so, which portion of the protein is likely to be oriented inside or outside the cell membrane. The topology analysis is an important result which can corroborate with other predictions. In this way, each bioinformatic tool adds an additional layer of comprehension about the HP, that can be linked with other results obtained in other programs (Tusnady and Simon, 2010).

Figure 3. Prediction of transmembrane helices in the hypothetical protein XP_002569027.1. Results generated through TMHMM. (—) Probability to be a transmembrane portion; (—) probability to be an intracellular portion; (—) probability to be an extracellular portion.

The GOR tool predicts secondary structures, indicating the probability of the presence of alpha helices, beta sheets, turns and random coils. As shown in Figure 4, of the 722 amino acids residues, 171 are related to alpha helices (23.68%); 145 are related to beta sheets (20.08%) and 406 are related to random coils (56.23%).

The 3D structure prediction of the XP_002569027.1 was performed by *SWISS-MODEL*, a web-based integrated service dedicated to protein structure homology modelling. Building a homology model comprises four main steps: the identification of structural template, the alignment of target sequence and template structure, the model-building, and finally, the model quality evaluation. When analyzing a protein, a protein structure database is searched to identify suitable templates. The similarity between target sequence and protein template library is used to select the best templates, and then, the most similar sequences are selected to estimate the 3D model of the HP (Bordoli et al., 2009). Figure 5 shows the predicted 3D structure of the HP

[illegible]

Understanding the structural properties of a protein is of great importance, since protein primary, secondary, tertiary and quaternary structure are directly linked with its functionality properties (Ji and Li, 2010). The 3D prediction of HPs by comparative modeling is an important tool capable of linking the protein to a specific function, since well known proteins, with similar structures and defined function are used as templates to create the HP model. Thus, a high amino acid identity can indicate a high structure similarity, and proteins with similar structures may have similar biochemical function, therefore proteins with similar amino acid sequence usually have related functions. Structural analyses are an essential step on HP characterization and function prediction (Da Costa et al., 2018).

Figure 5. Predicted 3D structural representation of the hypothetical protein XP_002569027.1 (blue and red) performed by comparison with the alpha-galactosidase protein AgaA of *G. stearothermophilus* (color). Prediction made by SWISS-MODEL tool. (A) 3D Structural prediction of hypothetical protein XP_002569027.1; (B) 3D structural representation of the alpha-galactosidase protein D478A AgaA A355E of *G. stearothermophilus*.

Prediction of HP XP_002569027.1 subcellular localization

TargetP tool was used to predict the subcellular localization of HP XP_002569027.1. The score 1 indicates the presence of N-terminal pre-sequences: MTP refers to the possibility to present the sequence of mitochondrial targeting peptide, SP the possibility of the sequence present signal peptide and others refers to the possibility of the sequence indicate another subcellular localization. The final results of the analyses generate four possible outcomes: C is indicative that the protein is found in the chloroplasts, M is indicative that the protein is mitochondrial, S is indicative that the protein is related to cellular secretion pathways or - is indicative that the protein is related to any other possible location.

Confirming the results from section 3.2, *TargetP* prediction indicates that HP XP_002569027.1 has a signal peptide (value = 0.805), indicating that it is a protein related to secretion pathway, more specifically an extracellular protein (S) (Table 1). As this program evaluates the presence of a signal peptide, we used the complete sequence containing 746 amino acids. In addition to supporting previous results, it also confirms conclusions of previous reports that alpha galactosidases in *P. rubens* and *Aspergillus sp* are extracellular enzymes, used for substrate degradation and obtaining energy (Manzanares et al., 1998; Aleksieva et al., 2010).

Table 1. Subcellular localization prediction of HP XP_002569027.1. Results obtained by *TargetP* platform. Modified. MTP: possibility to present the sequence of mitochondrial targeting peptide; SP: possibility of the sequence present signal peptide; Others: possibility of the sequence indicate another cellular location; Location: (S) indicative that the protein is related to cellular secretion pathways.

Name	Size	MTP	SP	Others	Location
XP_002569027.1	746 aa	0.131	0.805	0.036	S

Search for patterns of similarity with other proteins

The Basic Local Alignment Search Tool (*BLASTp*) finds regions of local similarity between sequences. The program compares nucleotide or protein sequences to sequence databases and calculates the statistical significance of matches. *BLASTp* can be used to infer functional and evolutionary relationships between sequences as well as help identify members of gene families. This is one of the most used and reliable tools on searching for sequence similarity and identity. Sequence similarity comparison, such as for DNA, RNA, and protein, is the base of bioinformatics, and this kind of analysis improve the protein function prediction, being one of the most important analysis in HPs characterization (Pearson, 2013; Yao et al., 2014).

The HP XP_002569027.1 from *P. rubens* has high amino acid identity with other proteins from *Penicillium* genus (Table 2) as two aldolases from *Penicillium expansum* (87.55%) and *Penicillium italicum* (87.42%). Aldolases are glycolytic pathway enzymes that catalyze the conversion of fructose-1,6-diphosphate to dihydroxyacetone phosphate and glyceraldehyde-3-phosphate, thus playing an important role on energetic metabolism (Chang et al., 2018).

The other class of proteins that showed high identity with the HP XP_002569027.1. was alpha-galactosidase, as this protein matches proteins annotated as alpha-galactosidases from other species of *Penicillium* genus as *Penicillium digitatum* (84.05%), *Penicillium janczewskii* (76.87%) and *Penicillium brasilianum* (73.35%) (Table 2). Alpha-galactosidases catalyze the removal of the α -galactose terminal of oligosaccharides and glycoproteins, such as raffinose and stachyose (Garman, 2007). This enzyme has several industrial applications in the food industry and has great potential applications in soybean-based food (Weignerová et al., 2009).

In addition, HP XP_002569027.1 also show identity with other members of the glycosidases family, showing high sequence identity with proteins from *Penicillium roqueforti* (87.94%) and *Penicillium camemberti* (87.69%). Aldolases and alpha galactosidases are included in the glycosidase family, enzymes in this family catalyze the hydrolysis of glycosidic molecules, such as oligosaccharides and glycoconjugates. This group of enzymes play important roles in biological processes, highlighting the importance of studies for elucidate the biological process and for industrial applications (Kotzler et al., 2014).

Besides these proteins that have defined functions, the two proteins that showed the highest identity with the HP XP_002569027.1. were also HP, one from *Penicillium flavigenum*, named PENFLA_c012G03777 with 98.26% of identity and the other from *Penicillium nalgiovense*, named PENNAL_c0022G06902 and presented 97.86% of identity (Table 2).

The 12 sequences that presented higher identity cited above are all proteins that belong to the *Penicillium* genus, which is expected as closely-related organisms tend to share high sequence similarity.

Table 2. Search for identity of the amino acid sequence of HP XP_002569027.1 with other proteins in NCBI database. Results obtained by *BLASTp* platform. Modified.

Reference (NCBI)	Description	Max Score	Total Score	Query Cover	E value	Per. Identity
XP_002569027.1	Pc21g20400 <i>Penicillium rubens</i> Wisconsin 54-1255	1535	1535	100%	0.0	100.00%
OQE22811.1	Hypothetical protein PENFLA_c012G03777	1515	1515	100%	0.0	98.26%
OQE86109.1	<i>Penicillium flavigenum</i> Hypothetical protein PENNAL_c0022G06902	1512	1512	100%	0.0	97.86%
XP_016597363.1	<i>Penicillium nalgioense</i> Aldolase TIM barrel-type <i>Penicillium expansum</i>	1376	1376	100%	0.0	87.55%
CDM31376.1	Glycosidase, clan GH-D <i>Penicillium roqueforti</i> FM164	1375	1375	100%	0.0	87.94%
KGO76349.1	Aldolase TIM barrel-type <i>Penicillium italicum</i>	1372	1372	100%	0.0	87.42%
CRL28250.1	Glycosidase, clan GH-D <i>Penicillium camemberti</i>	1337	1337	96%	0.0	87.69%
XP_014535717.1	Alpha-galactosidase C, <i>Penicillium digitatum</i> putative Pd1	1335	1335	100%	0.0	84.05%
KXG52816.1	Aldolase TIM barrel-type <i>Penicillium griseofulvum</i>	1317	1317	100%	0.0	87.40%
ABC70181.1	Alpha-galactosidase <i>Penicillium</i> sp. F63	1316	1316	95%	0.0	88.67%
ACN38386.1	Alpha-galactosidase <i>Penicillium janczewskii</i>	1213	1213	100%	0.0	76.87%
CEJ57078.1	Alpha-galactosidase putative <i>Penicillium brasiliense</i> C	1163	1163	99%	0.0	73.35%

Identification of Functional domains and important regions in HP XP_002569027.

Search for functional domains and important regions is a significant step on HP function prediction and characterization. Domains are functional and structural portions of a protein, which are usually linked to some specific function or interaction, contributing to the functionality of the protein as a whole (Yegambaram et al., 2013). In this sense, *InterProScan* is a resource that provides functional analysis of protein sequences by classifying them into families and predicting the presence of domains and important sites. To classify proteins in this way, *InterProScan* uses predictive models, known as signatures, provided by several different databases that make up the *InterPro* consortium (Mitchell et al., 2019). Protein families, as the name implies, are groups of proteins that have a similar evolutionary origin, which usually results in similarity in their structures and sequences (Yi et al., 2012).

Searching for conserved domains using *InterProScan* was possible to identify elements that corroborate with *BLASTp* results, showing that HP XP_002569027.1 has

domains that are conserved in different families of glycosylases, such as domains of alpha galactosidases and aldolases (Figure 6). Specifically, the first four lines of the figure show that the HP XP_002569027.1 presents regions of similarity with some proteins of a homologous superfamily (Figure 6 -blue H). This indicates not only that the protein may be related to glycoside metabolism, but it indicates that HP XP_002569027.1 probably shares evolutionary origins with the other proteins in this homologous superfamily. Also, it was possible to predict that the HP XP_002569027 have large similarity with proteins in Glycoside Hydrolase Family 36 (Figure 6- red F), showing high similarity with an alpha-galactosidase family named PIRSF005536 (Figure 6 - pink line). Lastly, the *InterProScan* analyses showed that the HP presented similar N-terminal and C-terminal domains (Figure 6- green D) with proteins in Glycosyl Hydrolase Family 36.

The association of HPs with families of proteins already described and with defined domains works as additional step to conclude the functional prediction, grouping all the previous analyzes in an attempt to link the HP to a probable function. This type of analysis is important because conserved domains are usually linked with the protein function, identify these domains in HPs is an important indication of functionality (Rentzsch and Orenge, 2013). Taken together, these results suggests that the HP XP_002569027.1 may be a protein related with carbohydrate metabolism.

Detailed signature matches

Figure 6. Search for functional domains, families and important regions of the hypothetical protein. Results obtained by InterProScan platform.

Alignment of similar sequences and search for conserved regions and catalytic residues

After verifying the homology among HP XP_002569027.1 with other know proteins from *Penicillium* genus, the next step was to obtain these sequences for a local alignment (item 3.6.) and perform a simultaneous align of them. To perform this analysis, it was used the *Multalin*, a tool which creates a multiple sequence alignment from a group of related sequences using progressive pairwise alignments using the method of multiple

sequence alignment with hierarchical clustering. It was selected the protein alpha-galactosidase from *P. janczewskii* (ACN38386.1, accessed on 25/09/2019), the alpha-galactosidase C (putative) from *P. digitatum* (XP_014535717.1, Accessed on 09/25/2019), alpha-galactosidase C from *A. flavus* (RAQ44733.1, Accessed on 09/25/2019) and aldolase from *P. expansum* (XP_016597363.1, Accessed on 25/09/2019).

After alignment, regions that have a high consensus, that is, sequences in which the amino acids were identical in all proteins, are marked in red. The regions marked in blue indicate low consensus, that is, only a few amino acids are conserved. The gray regions indicate neutrality, that is, almost no amino acids are conserved in this region. Our results indicate that a large number of regions have a high consensus (Figure 7). Moreover, it was verified that important catalytic residues are preserved in these sequences (505 and 567) and substrate-binding amino acids (216, 470 and 545) marked with a green triangle. This important residues were automatic annotated by UniProtKB (www.uniprot.org/uniprot/B6HKN2, Accessed on 09/25/2019), but literature has shown that residues such as serines 216, 470 (Margolles-Clark et al., 1996), glutamine 505 (Moracci et al., 1996; Liou and Grabowski, 2009), methionine 567 (Kachurin et al., 1995) and cysteine 545 (Kumar et al., 2014) are residues that play important roles in catalytic and active sites as well as in protein structure stabilization. In addition, of the few regions that are not red (high consensus), just a small number of amino acids are presented as very low / no consensus, being an indicative of sequence conservation (Figure 7).

Figure 7. Global alignment of sequences of the hypothetical protein XP_002569027.1 with proteins that present defined functions. Results obtained by MultAlin platform. (X) Represent high consensus regions; (X) regions of medium or low consensus; (X) very low or no consensus; () represents amino acids related to the catalytic site of the hypothetical protein XP_002569027.1.

To improve our study and provide new information about these similar proteins, we conducted physicochemical properties comparison using these five proteins. The results presented in Table 3, along with the previous results, show that the HP XP_002569027.1 have similar physicochemical parameters with these proteins, pointing once again for possible glucosidase function. The results showed that the HP XP_002569027.1 probably

plays an important role in energetic metabolism of *P. rubens*, with very similar characteristics of proteins that play this role in other related filamentous fungi.

Table 3. Comparison of physicochemical properties of the HP XP_002569027.1, alpha-galactosidase from *Penicillium janczewskii* (ACN38386.1), putative alpha-galactosidase from *Penicillium digitatum* (XP_014535717.1), alpha galactosidase from *Aspergillus flavus* (RAQ44733.1) and aldolase from *Penicillium expansum* (XP_016597363.1).

Property	Value				
	XP_002569027.1 <i>Penicillium rubens</i>	ACN38386.1 <i>Penicillium janczewskii</i>	XP_014535717.1 <i>Penicillium digitatum</i>	RAQ44733.1 <i>Aspergillus flavus</i>	XP_016597363.1 <i>Penicillium expansum</i>
Number of amino acids	722	748	765	751	747
Molecular weight	79.4 kDa	81.9 kDa	83.9 kDa	82.9 kDa	81.9 kDa
Theoretical pI	5.06	5.05	5.35	5.01	5.33
Total number of negatively charged residues (Asp + Glu)	86	90	86	87	85
Total number of positively charged residues (Arg + Lys)	59	58	64	58	62
Instability index	32.39	37.80	33.23	36.54	30.87
Aliphatic index	81.69	86.87	86.56	78.95	84.32

CONCLUSIONS

In this study, we characterized through *in silico* analyses the hypothetical protein from *P. rubens* (XP_002569027.1) and have attributed some important information to it, improving its status from hypothetical to a predicted function protein. Through interface-friendly bioinformatics tools, the HP XP_002569027.1 was characterized and information about its composition, secondary and tertiary structures, subcellular localization, similarity with other proteins were predicted. More importantly, we could assign a putative function to this protein. The results found here indicates that this HP is involved in glucose/galactose metabolism since it showed alpha galactosidase, aldolase and glycosidase domains. To confirm the functions here addressed to this hypothetical protein and prove its involvement in this metabolism, *in vivo* and *in vitro* studies should be performed. The results we obtained highlight the importance of bioinformatics as a tool for *in silico* studies, serving as a preliminary screening for *in vitro* analyses.

ACKNOWLEDGMENTS

This study was supported by CAPES (Conselho Nacional de Desenvolvimento Científico e Tecnológico), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) and FAPEMIG (Fundação de Amparo à Pesquisa do Estado de Minas Gerais). We wish to thank Cássio Siqueira Souza Cassiano for bioinformatics assistance.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Adinarayana KP, Sravani TS and Hareesh C (2011). The database of six hypothetical eukaryotic genes and proteins. *Bioinformation*. 6 (3): 128-130, doi: 10.6026/97320630006128.
- Aleksieva P, Tchorbanov B and Nacheva L (2010). High-Yield Production of Alpha-Galactosidase excreted from *Penicillium chrysogenum* and *Aspergillus Niger*. *Biotechnol. Biotechnol. Equip.* 24 (1): 1620-1623. doi.org/10.2478/V10133-010-0015-5.
- Almagro JJ, Tsigirgos KD, Sønderby CK, Petersen TN, et al. (2019). SignalP 5.0 Improves signal peptide predictions using neural deep networks. *Nat. Biotechnol.* 37: 420-423. doi: 10.1038/s41587-019-0036-z.
- Almagro JJ, Salvatore M, Emanuelsson O, Winther O, et al. (2019). Detecting sequence signals in targeting peptides using deep learning. *Life Sci. Alliance*. 2 (5); doi: 10.26508/lsa.201900429.
- Altschul SF, Gish W, Miller W, Myers EW, et al. (1990). Basic Local Alignment Search Tool. *J. Mol. Biol.* 215: 403-410. doi: 10.1016/S0022-2836(05)80360-2.
- Bayat A (2002). Science, medicine, and the future: Bioinformatics. *BMJ Clinical research* ed. 324 (7344): 1018-1022. doi: 10.1136/bmj.324.7344.1018.
- Bernstein FC, Koetzle TF, Williams GJB, Meyer EF, et al. (1977). The Protein Data Bank. *Eur. J. Biochem.* 80 (2): 319-324. doi: 10.1111/j.1432-1033.1977.tb11885.x.
- Bordoli L, Kiefer F, Arnold K, Benkert P, et al. (2009). Protein structure homology modeling using SWISS-MODEL workspace. *Nat. Protoc.* 4(1): 1-13, doi:10.1038/nprot.2008.197.
- Chang YC, Yang YC, Tien CP, Yang C, et al. (2018). Roles of Aldolase Family Genes in Human Cancers and Diseases. *Trends Endocrinol Metab.* 29 (8): 549-559. doi: 10.1016/j.tem.2018.05.003.
- Chinedu SN, Okafor UA, Thompson NE and Victoria IO (2008). Xylanase Production of *Aspergillus niger* and *Penicillium chrysogenum* from Ammonia Pretreated Cellulosic Waste. *Res. J. Microbiol.* 3: 246-253. 10.3923/jm.2008.246.253.
- Choo KH, Tan TW and Ranganathan S (2009). A comprehensive assessment of N-terminal signal peptides prediction methods. *BMC Bioinform.* 10: S2. doi.org/10.1186/1471-2105-10-S15-S2.
- Cole S, Brosch R, Parkhill J, T Garnier C, et al. (1998). Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence. *Nature*. 393 (6685): 537. doi: 10.1038/31159.
- Corpet F (1988). Multiple sequence alignment with hierarchical clustering. *Nucl. Acids Res.* 16 (22): 10881-10890. doi: 10.1093/nar/16.22.10881.
- Da Costa WLO, Araújo CL de A, Dias LM, Pereira LC de S, et al. (2018). Functional annotation of hypothetical proteins from the *Exiguobacterium antarcticum* strain B7 reveals proteins involved in adaptation to extreme environments, including high arsenic resistance. *PLoS One*. 13(6): e0198965. doi:10.1371/journal.pone.0198965.
- Dar GH, Kamili AN, Nazir R, Bandh SA, et al. (2015). Enhanced production of α -amylase by *Penicillium chrysogenum* in liquid culture by modifying the process parameters. *Microb. Pathog.* 88: 10-15. doi:10.1016/j.micpath.2015.07.016.
- De Simone G, Pasquabisceglie A, Proietto R, Polticelli F, et al. (2019). Contaminations in (meta)genome data: An open issue for the scientific community. *IUBMB Life*. doi:10.1002/iub.2216.
- Ferrer M, Plou FJ, Oscar MN, Reyes F, et al. (2000). Purification and properties of a lipase from *Penicillium chrysogenum* isolated from industrial wastes. *J. Chem. Technol.* 75: 569-576. doi: 10.1002/1097-4660(200007)75:7<569::AID-JCTB258>3.0.CO;2-S.
- Fiers W, Contreras R, Duerinck F, Haegeman G, et al. (1976). Complete nucleotide sequence of bacteriophage MS2 RNA: primary and secondary structure of the replicase gene. *Nature*. 260: 500-507. doi: 10.1038/260500a0.
- Garman SC (2007). Structure-function relationships in α -galactosidase A. *Acta Paediatr.* 96: 6-16. doi: 10.1111/j.1651-2227.2007.00198.x.
- Garnier J, Gibrat JF and Robson B. (1996). GOR Method for Predicting Protein Secondary Structure Amino Acid Sequence from. *Methods Enzymol, RF Doolittle Ed.* 266: 540-553. doi: 10.1016/s0076-6879(96)66034-0.
- Gasteiger E, Hoogland A, Gattiker DS, Wilkins MR, et al. (2005). Protein Identification and Analyses Tools on the ExPASy Server. John M. Walker (Ed) 1st edn: The Proteomics Protocols Handbook, Humana Press, Totowa NJ, pp. 571-607.
- Ge S and Guo YL (2020). Evolution of genes and genomes in the genomics era. *Sci China Life Sci.* 63. doi.org/10.1007/s11427-020-1672-0.
- Guruprasad K, Reddy, Reddy BV and Pandit MW (1991). Correlation between stability of a protein and its dipeptide composition: A novel approach for predicting *in vivo* stability of a protein from its primary sequence. *Protein Eng.* 4: 155-161. doi10.1093/protein/4.2.155.
- Guzman-chavex F, Zwahlen R, Bovenberg R and Driessen A (2018) Engineering of the filamentous fungus *Penicillium chrysogenum* as cell factory for natural products. *Front. Microbiol.* 9: 2768. doi:10.3389/fmicb.2018.02768.
- Haq IU, Mukhtar H and Umber H (2005). Production of Protease by *Penicillium chrysogenum* Through Optimization of Environmental Conditions. *J. Agri. Soc. Sci.* 2.

- Hawksworth DL and Lücking R (2017). Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiol Spectrum*. 5(4). doi:10.1128/microbiolspec.FUNK-0052-2016.
- Hesper B and Hogeweg P (1970). "Bioinformatica: een werkconcept. *Kameleon* 1 (6): 28-29. Dutch. Leiden: Leidse Biologen Club.
- Houbraken J, Frisvad JC and Samson RA (2011). Fleming's penicillin producing strain is not *Penicillium chrysogenum* but *P. rubens*. *IMA Fungus*. 2(1): 87-95. doi:10.5598/imafungus.2011.02.01.12.
- Ijaq J, Malik G, Kumar A, Das PS, et al. (2019). A model to predict the function of hypothetical proteins through a nine-point classification scoring schema. *BMC Bioinform*. 20: 14. <https://doi.org/10.1186/s12859-018-2554-y>.
- Jami MS, García-Estrada C, Barreiro C, Cuadrado AA, et al. (2010). The *Penicillium chrysogenum* Extracellular Proteome. Conversion from the Food-rotting Strain to a Versatile Cell Factory for White Biotechnology. *Mol. Cell. Proteomics*. 9 (12): 2729-2744. doi: 10.1074/mcp.M110.001412.
- Ji YY and Li YQ (2010). The role of secondary structure in protein structure selection. *Eur. Phys. J. E*. 32(1): 103-107. doi:10.1140/epje/i2010-10591-5.
- Jones P, Binns D, Chang HY, Fraser M, et al. (2014). InterProScan 5: genome-scale protein function classification. *Bioinformatics*. 30(9): 1236-1240. doi: 10.1093/bioinformatics/btu031.
- Kachurin AM, Golubev AM, Geisow MM, Veselkina OS, et al. (1995). Role of methionine in the active site of alpha-galactosidase from *Trichoderma reesei*. *Biochem J*. 308: 955-964. doi:10.1042/bj3080955.
- Kapp K, Schrempf S, Lemberg MK and Dobberstein B (2009). Post-Targeting Functions of Signal Peptides. In: *Madame Curie Bioscience Database* [Internet]. Landes Bioscience; Austin TX. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK6322/>. Accessed on 09/25/2019.
- Kim Y, Nandakumar MP and Marten MR (2007). Proteomics of filamentous fungi. *Trends in Biotechnol*. 25 (9): 395-400. doi: 10.1016/j.tibtech.2007.07.008.
- Kotzler MP, Hancock SM and Withers SG (2014). Glycosidases: Functions, Families and Folds. *eLS, John Wiley & Sons Ltd: Chichester*. doi: 10.1002/9780470015902.a0020548.pub2.
- Krogh A, Larsson B, Von Heijne G and Sonnhammer ELL (2001). Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes. *J. Mol. Biol*. 305 (3): 567-580. doi: 10.1006/jmbi.2000.4315x.
- Kumar V, Sharma N and Bhalla TC (2014). In Silico Analysis of β -Galactosidases Primary and Secondary Structure in relation to Temperature Adaptation. *J. Amino Acids*. doi:10.1155/2014/475839.
- Lander ES, Linton LM, Birren B, Nusbaum C, et al. (2001). Initial sequencing and analysis of the human genome. *Nature*. 409: 860-921. doi.org/10.1038/35057062.
- Liou B and Grabowski GA (2009). Participation of asparagine 370 and glutamine 235 in the catalysis by acid beta-glucosidase: the enzyme deficient in Gaucher disease. *Mol. Genet. Metab*. 97: 65-74. doi:10.1016/j.ymgme.2009.01.006.
- Lowe R, Shirley N, Bleackley M, Dolan, S, et al. (2017). Transcriptomics technologies. *PLoS Comput. Biol*. 13(5): doi.org/10.1371/journal.pcbi.1005457.
- Lubec G, Afjehi-Sadat L, Yang JW and John JP (2005) Searching for hypothetical proteins: theory and practice based upon original data and literature. *Prog Neurobiol*. 77(1-2): 90-127. doi:10.1016/j.pneurobio.2005.10.001.
- Mallick P and Kuster B (2010). Proteomics: a pragmatic perspective. *Nat. Biotechnol*. 28 (7): 695-709. doi: 10.1038/nbt.1658.
- Manzanares P, de Graaff LH and Visser J (1998). Characterization of galactosidases from *Aspergillus niger*: Purification of a Novel α -galactosidase Activity. *Enzyme Microb. Tech*. 22 (5): 383-390. doi: 10.1016/s0141-0229(97)00207-x.
- Margolles-Clark E, Tenkanen M, Soderlund H and Penttilä M (1996). Acetyl xylan esterase from *Trichoderma reesei* contains an active site serine and a cellulose binding domain. *Eur. J. Biochem*. 237: 553-560. doi:10.1111/j.1432-1033.1996.0553p.x.
- Meena M, Zehra A, Dubey MK, Aamir, et al. (2018). *Penicillium* Enzymes for the Food Industries. New and Future Developments in Microbial Biotechnology and Bioengineering, 1st edition. 167-186. doi:10.1016/b978-0-444-63501-3.00009-0.
- Mishra PK, Sonkar SC, SR Raj Chaudhry U, Saluja D, et al. (2013). Functional Analyses of hypothetical proteins of *Chlamydia trachomatis*: A Bioinformatics Based Approach for Prioritizing the Targets. *J. Comput. Sci. Syst. Biol*. 7: 010-014. doi: 10.4172/jcsb.1000132.
- Mitchell AL, Attwood TK, Babbitt PC, Blum M, et al. (2019). InterPro in 2019: improving coverage, classification and access to protein sequence annotations. *Nucleic Acids Res*. 47: 351- 360. doi.org/10.1093/nar/gky1100.
- Mohan R and Venugopal S (2012). Computational structural and functional analysis of hypothetical proteins of *Staphylococcus aureus*. *Bioinformation*. 8(15): 722-728. doi:10.6026/97320630008722.
- Moracci M, Capalbo L, Ciaramella M and Rossi M (1996). Identification of two glutamic acid residues essential for catalysis in the β -3-glycosidase from the thermoacidophilic archaeon *Sulfolobus solfataricus*. *Protein Eng*. 9: 1191-1195. doi.org/10.1093/protein/9.12.1191.
- Naveed M, Chaudhry Z, Ali Z, Amjad M, et al. (2018). Annotation and curation of hypothetical proteins: prioritizing targets for experimental study. *Adv. Life Sci*. 5 (3): 73-87.

- Papageorgiou L, Eleni P, Raftopoulou S, Mantaïou M, et al. (2018). Genomic big data hitting the storage bottleneck. *EMBNet J.* 24: e910.
- Pearson WR (2013). An Introduction to Sequence Similarity (“Homology”) Searching. *Curr Protoc Bioinformatics.* 42(1): 3.1.1-3.1.8. doi:10.1002/0471250953.bi0301s42.
- Peberdy JF (1994). Protein secretion in filamentous fungi - trying to understand the highly productive black box. *Trends Biotechnol.* 12 (2): 50-57. doi.org/10.1016/0167-7799(94)90100-7.
- Pohl C, Polli F, Schütze T, Viggiano A, et al. (2020). A *Penicillium rubens* platform strain for secondary metabolite production. *bioRxiv.* doi.org/10.1101/2020.04.05.026286. t.
- Poliakov E, Cooper DN, Stepchenkova EI and Rogozin IB (2015). Genetics in Genomic Era. *Genet. Res.* doi: 10.1155/2015/364960.
- Punt PJ, Van Biezen N, Conesa A, Albers A, et al. (2002). Filamentous fungi the cell factories for heterologous protein production. *Trends in Biotechnol.* 20 (5): 200-206. doi: 10.1016/s0167-7799(02)01933-9.
- Rentzsch R and Orengo CA. (2013). Protein function prediction using domain families. *BMC Bioinformatics.* 14 Suppl 3(Suppl 3): S5. doi:10.1186/1471-2105-14-S3-S5.
- Schmutz J, Wheeler J, Grimwood J, Dickson M, et al. (2004). Quality assessment of the human genome sequence. *Nature.* 429(6990): 365-368. doi: 10.1038/nature02390.
- Sivashankari S and Shanmughavel P (2006). Functional annotation of hypothetical proteins - A review. *Bioinformation.* 1(8): 335-338. doi:10.6026/97320630001335.
- Skolnick J, Fetrow JS and Kolinski A (2000). Structural genomics and its importance for gene function analysis. *Nat. Biotechnol.* 18(3): 283-287. doi:10.1038/73723.
- Srinivasa KG, Siddesh GM and Manisekhar SR (2020). Statistical Modelling and Machine Learning Principles for Bioinformatics Techniques, Tools, and Applications. Algorithms for Intelligent Systems. *Springer.* 3-9. doi:10.1007/978-981-15-2445-5.
- Tusnady GE and Simon I (2010). Topology Prediction of Helical Transmembrane Proteins: How Far Have We Reached? *Curr. Protein Pept. Sc.* 11(7): 550-561. doi:10.2174/138920310794109184.
- Uranga DC, Ghassemian M and Martinez-Hernandez A (2017). Novel proteins from proteomic analyses of the trunk fungus disease *Lasioidiplodia theobromae* (Botryosphaeriaceae). *Biochimie Open.* 4: 88-98. doi: 10.1016/j.biopen.2017.03.001.
- Van den Berg MA, Albang R, Albermann K, Badger JH, et al. (2008). Genome sequencing and analyses of the filamentous fungus *Penicillium chrysogenum*. *Nat Biotechnol.* 26 (10): 1161-1168. doi: 10.1038/nbt.1498.
- Waterhouse A, Bertoni M, Bienert S, Studer G, et al. (2018). SWISS-MODEL: homology modeling of protein structures and complexes. *Nucleic Acids Res.* 296-303. doi: 10.1093/nar/gky427.
- Weignerová L, Simerská P and Křen V (2009). α -Galactosidases and their applications in biotransformations. *Biocatal. Biotransformation.* 27(2): 79-89. doi:10.1080/10242420802583416.
- Yakimova BK, Tchorbanov BP and Stoineva IB (2017). Alpha-galactosidase and invertase from *Penicillium chrysogenum* sp.23: purification, characteristics and hydrolysis of raffinose. *Bulg. Chem. Special.* 101-106.
- Yang Z, Zeng X and Tsui SK (2019). Investigating function roles of hypothetical proteins encoded by the *Mycobacterium tuberculosis* H37Rv genome. *BMC Genomics.* 20: 394. doi.org/10.1186/s12864-019-5746-6.
- Yao Y, Yan S, Xu H, Han J, et al. (2014) Similarity/Dissimilarity analysis of protein sequences based on a new spectrum-like graphical representation. *Evol Bioinform Online.* 10: 87-96. doi:10.4137/EBO.S14713.
- Yegambaram K, Bulloch EM and Kingston RL (2013). Protein domain definition should allow for conditional disorder. *Protein Sci.* 22(11): 1502-1518. doi:10.1002/pro.2336.
- Yi G, Thon MR and Sze SH (2012). Supervised protein family classification and new family construction. *J. Comput. Biol.* 19(8): 957-967. doi:10.1089/cmb.2011.0044.