Essential Oil and Fatty Acid Composition of Endemic *Gypsophila laricina* Schreb. from Turkey

Hüseyin Servi, Betül Eren Keskin, Sezgin Çelik, Ümit Budak, Büşra Kababıyık

¹Department Of Pharmaceutical Botany, Faculty Of Pharmacy, Altınbas University, Istanbul, Turkey

²Department Of Molecular Biology And Genetics, Faculty Of Engineering And Natural Sciences, Uskudar University, Istanbul, Turkey

³Department Of Molecular Biology And Genetics, Faculty Of Arts & Sciences, Yıldız Technical University, Istanbul, Turkey

⁴Department Of Biology, Faculty Of Arts & Science, Bozok University, Yozgat, Turkey

INTRODUCTION: *Gypsophila* species have very high medicinal and commercial importance and also contains interesting natural substances. However, there is no report on the essential oil and fatty acid composition of any *Gypsophila* species. This prompted us to investigate the essential oil and fatty acid composition of *Gypsophila laricina* Schreb.

METHODS: Plant materials were collected during the flowering period. Essential oil composition of aerial parts of *Gypsophila laricina* Schreb. were analysed by gas chromatography and gas chromatography-mass spectrometry. And the fatty acid compositions were analysed by gas chromatography-mass spectrometry.

RESULTS: Sixty-six and ten compounds were identified in the essential oil and fatty acid of *G. laricina* Schreb. respectively. The major components of the essential oil were hexadecanoic acid (27.03%) and hentriacontane (12.63%). The main compounds of the fatty acid were (Z,Z)-9,12-octadecadienoic acid methyl ester (18: 2) 40.4%, (Z)-9-octadecenoic acid methyl ester (18: 1) 35.0 % and hexadecanoic acid methyl ester (16: 0) 13.0%.

DISCUSSION AND CONCLUSION: The result showed that the fatty acid composition are rich in polyunsaturated fatty acids. The essential oils of *G. laricina* Schreb. were dominated by fatty acid derivatives and n-alkanes. We believe the results obtained

from this research will stimulate further research on the chemistry of *Gypsophila* species.

Keywords: *Gypsophila laricina*, essential oil, fatty acid

INTRODUCTION

Caryophyllaceae family has about 85 genera and 2630 species in the world and distributed mainly in Mediterranean and Iran-Turan areas.¹ *Gypsophila* is the 3rd biggest genus in family of Caryophyllaceae to Turkey. *Gypsophila* species are annual, biennial or perennial herbaceous plants. Stem length of the plant is about 1 m and its flowering time is June and July.²

Some *Gypsophila* species are used in folk medicine as remedies for cough, cold, ailments of the upper respiratory tract³ and also used for medical treatment such as expectorant, diuretic, hepatitis, gastritis and bronchitis.⁴ The underground parts of *Gypsophila* genus have triterpenoid saponins as a main component. *Gypsophila* genus are used in industrial, medicinal and decorative application.⁵ The commercial *Merck* saponin which has been widely utilized as a standard for hemolytic test was obtained from the roots of several *Gypsophila* species.³ The genus were reported to have cytotoxic activity, α -glucosidase activity, immune-modulating effect and normalization of carcinogen induced cell proliferation.^{4,6} The saponins that get from *Gypsophila* genus are interest in terms of their applications in vaccines.⁷ Biological activities of the genus seems to be associated with triterpene saponins. Due to the various beneficial biological activities, *Gypsophila* was the focus of studies that described the phytochemistry of the genus extensively.

Previously, antioxidant and antibacterial activities of chloroform extracts of underground parts of *Gypsophila eriocalyx* and *Gypsophila sphaerocephala* var. *sphaerocephala* were investigated. The chloroform extracts of both species had high antioxidant properties but showed low antibacterial activity.⁸

Additionally, the toxic boron levels of some plant species (*Gypsophila sphaerocephala* var. *sphaerocephala*, *Gypsophila perfoliata*, *Puccinellia distans* subsp. *distans* and *Elymus elongates*) were reported. Among these plant species, *G. sphaerocephala* contained considerably higher boron concentrations in its above-ground parts compared to the roots and organs of the other species. This study

shows that *G. sphaerocephala* was not only able to grow on heavily boron contaminated soils, but was also able to accumulate extraordinarily high concentrations of boron.⁹

According to study from Iran, antimicrobial activity and chemical constituents of the essential oils from flower, leaf and stem of *Gypsophila bicolor* were investigated. The main components of the essential oil from flower were germacrene-D (21.2 %), *p*-cymene (20.6 %), bicyclogermacrene (17.6 %), γ -dodecadienolactone (11.7%) and terpinolene (9.4 %). The main components of the essential oil from leaves were germacrene-D (23.4 %), terpinolene (14.5 %), bicyclogermacrene (11.5 %), γ -dodecadienolactone (6.8 %), *p*-cymene (6.7 %) and *cis*- β -ocimene (6.3 %). The main components of the essential oil from stems were γ -dodecadienolactone (28.5 %), bicyclogermacrene (14.8 %), germacrene-D (12.6 %), *p*-cymene (12.5 %), terpinolene (11.6 %) and *trans*- β -ocimene (4.2 %). The essential oils had moderate effect on Gram-positive and Gram negative bacteria, but had significant effect on the fungi.¹⁰

Another study from Turkey, essential oil composition and fatty acid profile of *Gypsophila tuberculosa* and *Gypsophila eriocalyx* were reported. The main components of essential oils were determined hexadecanoic acid (25.3%) and hentriacontane (13.0%) for *G. tuberculosa* and octacosane (6.83%), eicosanal (6.19%), triacontane (5.03 %) and heneicosane (5.78%) for *G. eriocalyx*. The major compounds of fatty acids of *G. tuberculosa* and *G. eriocalyx* were (*Z*)-9-octadecenoic acid methyl ester (32.0%, 36.0%), (*Z,Z*)-9,12-octadecadienoic acid methyl ester (19.6%, 18.7%), and hexadecanoic acid methyl ester (17.7%, 25.2%), respectively.¹¹

As summarized above *Gypsophila* species have very high medicinal and commercial importance and also contains interesting natural substances. However, according to our literature survey we have not encountered any reports on the essential oil and fatty acid composition of *Gypsophila laricina* Schreb. This prompted us to investigate the essential oil and fatty acid composition of *Gypsophila laricina* Schreb. Here we report for the first time on the essential oil composition and fatty acid profile of *Gypsophila laricina* Schreb.

EXPERIMENTAL

Plant Materials

Plant materials were collected during the flowering period; *Gypsophila laricina* Schreb. was collected from 1740 m–1800 m height in Ucpinar, Sarkisla, Sivas, Turkey in July 2015 by Celik and Budak. Voucher specimen has been deposited in the Herbarium of Bozok University (Voucher no. Bozok HB 3302).

Fatty Acids Analyses

The aerial parts of the collected specimen, dried separately in shadow and ground with an electric mill (Retsch SM 100). Aerial part of the plant (400 g) was extracted with hexane for three days at room temperature. After filtration through filter paper, the extract was concentrated by rotary evaporator and 4 g crude hexane extract were obtained from the aerial parts of *Gypsophila laricina* Schreb. The crude extract was stored at 4°C. In this study we used hexane extract for fatty acid compositions. Methyl-ester derivatives of fatty acids (FAME), found in hexane extract was obtained by transesterification method.¹¹ In this method the dried extract was dissolved in 5 mL hexane and then extracted with 2 M methanolic KOH at room temperature. The mixture was shaken for 2 min and allowed to stand for 10 min. The upper phases were removed. *G. laricina* Schreb. afforded fixed oil from the hexane extract with 0.07% (v/w) yields. The fixed oils were analysed by GC-MS system.

Essential Oils Analyses

Aerial parts (200 g) of the dried plants subjected to hydrodistillation for 3 h, using a Clevenger-type apparatus to produce essential oils. Condenser of the Clevenger was attached to a microchiller that set to 4°C. *G. laricina* Schreb. afforded oils from the aerial parts with 0.01% (v/w) yields. The oils were recovered with 1 mL *n*-hexane and preserved in amber vials under -20°C until the day they were analysed.

Gas chromatography/mass spectrometry for fatty acids

The fatty acid compositions of the hexane extracts were investigated by means of Gas Chromatography-Mass Spectroscopy (GC-MS) system. The fatty acid methyl esters were analyzed using Agilent 5975C GC-MSD system with Innowax FSC polar column (30m x 0.25 mm, 0.25 µm). The inlet temperature was set at 250°C. Helium was the carrier gas at a constant flow rate of 1 ml/min. Split ratio was set to 50:1. The

oven temperature was programmed from 40°C to 210°C at rate of 5°C/min and kept constant at 210°C for 10 min. EI/MS was taken at 70 eV ionization energy. Mass range was from m/z 35-450 amu (atomic mass unit). Relative percentage amounts of the separated compounds were calculated from integration of the peaks in MS chromatograms. Identification of fatty acid components were carried out by comparison of their retention indices (RI) obtained by series of *n*-alkanes (C5 to C30) to the literature and with mass spectra comparison.¹³⁻¹⁹ Mass spectra comparison was done by computer matching with commercial Wiley 8th Ed./NIST 05 Mass Spectra library. Analysis was completed in 50 minutes.

Gas Chromatography/Mass Spectrometry for essential oils

The GC-MS analysis was performed with an Agilent 5975C GC-MSD system operating in EI mode. Essential oil samples were diluted 1/100 (v/v) with *n*-hexane. Injector and MS transfer line temperatures were set at 250°C. Innowax FSC column (60 m x 0.25 mm, 0.25 µm film thickness) and helium as carrier gas (1 mL/min) were used in both GC/MS analyses. Splitless injection was employed. Oven temperature was programmed to 60°C for 2 min. and raised to 220°C at rate of 4°C/min. Temperature kept constant at 220°C for 10 min. and then raised to 240°C at a rate of 1°C/min. Mass spectra were recorded at 70 eV with the mass range m/z 35 to 425.

Gas Chromatography for essential oils

The GC analyses were done with an Agilent 6890N GC system. FID detector temperature was set to 300°C and same operational conditions applied to a duplicate of the same column used in GC-MS analyses. Simultaneous auto injection was done to obtain the same retention times. Relative percentage amounts of the separated compounds were calculated from integration of the peaks in FID chromatograms. Identification of essential oil components were carried out by comparison of their relative retention indices (RRI) obtained by series of *n*-alkanes (C5 to C30) to the literature and with mass spectra comparison.²⁰⁻⁴⁰ Mass spectra comparison was done by computer matching with commercial Wiley 8th Ed./NIST 05 Mass Spectra library, Adams Essential Oil Mass Spectral Library and Pallisade 600K Complete Mass Spectra Library.

RESULT AND DISCUSSION

Fatty acid composition of *Gypsophila laricina* Schreb. were analysed by GC-MS. Ten compounds were identified in the fatty acid that represent 98.9% of the fatty acid. The extract consisted of six saturated fatty acids (21.8%) and four unsaturated fatty acids (77.2%). The major components of the fatty acid were (Z,Z)-9,12-octadecadienoic acid methyl ester (Linoleic acid) (18:2) 40.4%, (Z)-9-octadecenoic acid methyl ester (Oleic acid) (18:1) 35.0% and hexadecanoic acid methyl ester (Palmitic acid) (16:0) 13.0%. The fatty acid composition of *Gypsophila laricina* Schreb. are represented in Table 1.

Essential oil composition of *Gypsophila laricina* Schreb. were analysed by GC and GC-MS. The essential oils of aerial parts of *Gypsophila laricina* Schreb. afforded very low oils yields 0.03% (v/w) yield). Sixty-six compounds were identified in the essential oil of *G. laricina* Schreb. by GC system that represent 76.9% of the oil. The major components of the oil were hexadecanoic acid (27.02%) and hentriacontane (12.63%). The essential oil composition of *Gypsophila laricina* Schreb. are given in Table 2.

The essential oil composition of *Gypsophila laricina* showed similar chemical behaviour to *G. tuberosa*.¹¹ Both species had hexadecanoic acid and hentriacontane as major components in essential oils. But hexadecanoic acid contained at 4.64% levels in *G. eriocalyx* and in a nearly six amount in *G. tuberosa* and *G. laricina*. And also hentriacontane contained at very low amounts in *G. eriocalyx*.¹¹ Three *Gypsophila* species had linoleic acid, oleic acid and palmitic acid as main components in different percentage in fatty acids.

According to a study from Iran, *Gypsophila bicolor* was reported to contain germacrene⁵, *p*-cymene, bicyclogermacrene, γ -dodecadienolactone, terpinolene, *cis*- β -ocimene and *trans*- β -ocimene¹⁰ however these compounds were not detected in the oil of *Gypsophila laricina* Schreb. *Gypsophila laricina* Schreb. showed very different chemical behaviour from *G. bicolor*. These differences in the previous literature and present data could be related to different collection times, climatic and soil conditions, ecological factors, methods and instruments employed in analysis or different genotypes. There are very few reports on the essential oil or volatile composition of *Gypsophila* species therefore it is difficult to produce a comment on

the chemo-systematic position of this species according to current findings and the existing reports.

CONCLUSION

The essential oil composition and fatty acid profile of *G. laricina* Schreb. are investigated for the first time. The major components of the fatty acid were oleic acid, linoleic acid and palmitic acid. The unsaturated fatty acids content were higher than their saturated fatty acids content. The essential oils of *G. laricina* Schreb. were dominated by fatty acid derivatives and *n*-alkanes. Hexadecanoic acid and hentriacontane are found as major components in essential oil. The high hexadecanoic acid content might be explained about the collection time of plant materials at late flowering period. *Gypsophila laricina* exhibited important differences *Gypsophila bicolor* and *Gypsophila eriocalyx* highlighting the existence of different main chemical constituents. Thus, the results of this study certainly contributed to the taxonomy of the genus *Gypsophila* via essential oil chemistry. We believe the results obtained from this research will stimulate further research on the chemistry of *Gypsophila* species.

ACKNOWLEDGEMENTS

We are indebted to YTÜBAP 2015-01-07-KAP05 and TUBITAK (Project no. TBAG-111T820) for the financial support.

REFERENCE

1. Mabberley D.J. Mabberley's plant-book: a portable dictionary of plants, their classifications, and uses (3th ed). Cambridge; Cambridge University Press; 2002;1-300.
2. Davis PH, Mill RR, Tan K and Edmondson JR. Flora of Turkey and the East Aegean Islands (2nd ed). Edinburgh; University Press; 1982;2;149-180.
3. Elbandy M, Miyamoto T. & Lacaille-Dubois MA. Sulfated lupane triterpene derivatives and a flavone C-glycoside from *Gypsophila repens*. Chem Pharm Bull. 2007;55:808-811.
4. Yücekutlu AN. & Bildacı I. Determination of Plant Saponins and Some of *Gypsophila* Species: A review of the literature. Hacettepe J Biol Chem. 2008;36:129-135.

5. Krasteva IN. Popov IS. Balabanova VI. Nikolov SD. & Pencheva IP. Phytochemical study of *Gypsophila trichotoma* wend (Caryophyllaceae). Quim Nova. 2008;31:1125-1126
6. Xie LX. Zhang HC. Wang HY. Wang Y. Wang FL. & Sun JY. Two new triterpenoids from *Gypsophila oldhamiana*. Nat Prod Res. 2016;30:1068-1074.
7. Gevrenova R. Stancheva T. Voynikov Y. Laurain-Mattar D. & Henry M. Root in vitro cultures of six *Gypsophila* species and their saponin contents. Enzyme Microb Tech. 2010;47:97-104.
8. Çona IG. *Gypsophila eriocalyx* Boiss. ve *G. sphaerocephala* Fenzl ex tchihat var. *sphaerocephala* taksonlarının antioksidan ve antibakteriyel aktivitelerinin belirlenmesi. Master thesis. Pamukkale University. 2012.
9. Babaoglu M. Gezgin S. Topal A. Sade B. & Dural H. *Gypsophila sphaerocephala* Fenzl ex Tchihat.: a boron hyperaccumulator plant species that may phytoremediate soils with toxic B levels. Turk J Bot. 2014;28:273-278.
10. Shafagha A. Shafaghatlonbar M. Antimicrobial activity and chemical constituents of the essential oils from flower, leaf and stem of *Gypsophila bicolor* from Iran. Nat Prod Commun. 2011;6:215-217.
11. Servi H. Keskin BE. Celik S. Budak U. Kızıltas FC. and Bektas EB. Essential oil composition and fatty acid profile of two endemic *Gypsophila* species from Turkey. Am J Essent Oil Nat Prod. 2017;5:16-20
12. Paquot C. Standard Methods for the Analysis of Oils, Fats and Derivatives (6th ed). Oxford;Pergamon Press, 2011;16:1-188.
13. Eyol PC. Sarikahya NB. Kankoc CC. Gokce A. Demirci F. Kirmizigul S. & Goren N. Fatty Acid Composition and Biological Activities of *Tanacetum zahlbruckneri* (Náb.) Giesb. Growing in Turkey. Rec Nat Prod. 2017;11:401-405.
14. Kirmizigul S. Sarikahya NB. Sümbül H. Arda N. Essential fatty acid components and antioxidant activities of eight *Cephalaria* species from southwestern Anatolia. Pure Appl Chem. 2007;79:2297-2304.
15. Maltas E. Vural CM. Yildiz S. Antioxidant activity and fatty acid composition of *Ginkgo biloba* from Turkey. Pure Appl Chem. 2011;35:803-818.
16. Laribi B. Rouki K. Mougou A. Marzouk B. Fatty acid and essential oil composition of three Tunisian caraway (*Carum carvi* L.) seed ecotypes. J Sci Food Agr. 2010;90:391-396.
17. Denis C. Wielgosz-Collin G. Bretch A. Ruiz N. Rabesaotra V. Boury-Esnault N. Kornprobst JM. Barnathan G. New 17-Methyl-13-Octadecenoic and 3,16-Docosadienoic Acids from the Sponge *Polymastia penicillus*. Lipids. 2009; 44:655-663.
18. Tsevegsuren N. Fujimoto K. Christie WW. Endo Y. Occurrence of a Novel *cis,cis,cis*-Octadeca-3,9,12-trienoic (*Z,Z,Z*-octadeca-3,9,12-trienoic) Acid in *Chrysanthemum* (*Tanacetum*) *zawadskii* Herb. (Compositae) Seed Oil. Lipids. 2003;38:573-578.

19. Nickavar B. Mojab F. Javidnia K. Amoli MA. Chemical Composition of the Fixed and Volatile Oils of *Nigella sativa* L. from Iran. Z Naturforsch C. 2003;58:629-631.
20. Polatoğlu K. Şen A. Bulut G. Bitiş L. Gören N. Essential Oil Composition of *Centaurea stenolepis* Kerner. from Turkey. J Essent Oil Bear Pl. 2014;17:1268-1278.
21. Polatoğlu K. Arsal S. Demirci B. Başer KH. DPPH Scavenging, PRAP Activities and Essential Oil Composition of Edible *Lathyrus ochrus* L. (Cyprian Vetch, Luvana) from Cyprus. J Oleo Sci. 2015;64:309-314.
22. Polatoğlu K. Demirci F. Demirci B. Gören N. Başer KHC. Antibacterial Activity and the Variation of *Tanacetum parthenium* (L.) Schultz Bip. Essential Oils from Turkey. J Oleo Sci. 2010;59:177-184.
23. Polatoğlu K. Demirci B. Gören N. Başer KHC. Essential oil composition of *Tanacetum kotschyi* from Turkey. Chem Nat Compd. 2011;47:297-299.
24. Polatoğlu K. Gören N. Karakoç ÖC. Phytotoxic, DPPH scavenging, insecticidal activities and essential oil composition of *Achillea vermicularis*, *A. teretifolia* and proposed chemotypes of *A. biebersteinii* (Asteraceae). Ind Crop Prod. 2013;51:35-45.
25. Polatoğlu K. Karakoç ÖC. Yücel YY. Demirci B. Gören N. Başer KH. Composition, insecticidal activity and other biological activities of *Tanacetum abrotanifolium* Druce essential oil. Ind Crop Prod. 2015;71:7-14.
26. Polatoğlu K. Demirci B. Demirci F. Gören N. Başer KH. Biological activity and essential oil composition of two new *Tanacetum chiliophyllum* (Fisch. & Mey.) Schultz Bip. var. *chiliophyllum* chemotypes from Turkey. Ind Crop Prod. 2012; 39:97-105.
27. Sura BE. Demirci B. Demirci F. Karaalp C. Başer KH. Composition of the essential oils of *Centaurea aphrodisea*, *C. polyclada*, *C. athoa*, *C. hyalolepis* and *C. iberica*. J Essent Oil Res. 2013;25:79-84.
28. Karamenderes C. Demirci B. Başer KH. Composition of Essential Oils of Ten *Centaurea* L. taxa from Turkey. J Essent Oil Res. 2008;20:342-349.
29. Demirci B. Başer KH. Yıldız B. Bahçecioğlu Z. Composition of the essential oils of six endemic *Salvia* spp. from Turkey. Flavour Frag J. 2003;18:116-121.
30. Moronkola DO. Ogunwande IA. Başer KHC. Ozek T. Ozek G. Essential Oil Composition of *Gmelina arborea* Roxb., Verbenaceae from Nigeria. J Essent Oil Res. 2009;21:264-266.
31. Bassolet IHN. Ouattara AS. Nebie R. Ouattara CAT. Kabore ZI. Traore SA. Chemical composition and antibacterial activities of the essential oil of *Lippia chevalieri* and *Lippia multiflora* from Burkina Faso. Phytochemistry, 2003;62: 209-212.
32. Kose YB. Altintas A. Tugay O. Uysal T. Demirci B. Ertugrul K. Baser KHC. Composition of the Essential Oils of *Centaurea sericeae* Wagenitz and *Centaurea ensiformis* P.H. Davis from Turkey. Asian J Chem. 2010;22:7159-7163.

33. Kirimer N. Tabanca N. Özek T. Tümen G. Baser KHC. Essential oils of Annual *Sideritis* Species Growing in Turkey. Pharm Biol. 2000;38:106-111.
34. Saidana D. Mahjoub MA. Boussaada O. Chriaa J. Cheraif I. Daami M. Mighri Z. Helal AN. Chemical composition and antimicrobial activity of volatile compounds of *Tamarix boveana* (Tamaricaceae). Microbiol Res. 2008;163:445-455.
35. Bardakcı H. Demirci B. Yesilada E. Kirmizipekmez H. Baser KHC. Chemical Composition of the Essential Oil of the Subterranean Parts of *Valeriana alliariifolia*. Rec Nat Prod. 2012;6:89-92.
36. Tabanca N. Demirci B. Ozek T. Kirimer N. Baser KHC. Bedir E. Kocaoglu M. Wedge DE. Gas chromatographic–mass spectrometric analysis of essential oils from *Pimpinella* species gathered from Central and Northern Turkey. J Chromatogr A, 2006;1117:194-205.
37. Polatoglu K. Demirci F. Demirci B. Gören N. Baser KHC. Antimicrobial activity and Essential oil composition of a new *T. argyrophyllum* (C.Koch) Tuzel. var. *argyrophyllum* chemotype. J Oleo Sci. 2010;59:307-313.
38. Dregus M. Engel KH. Volatile constituents of uncooked rhubarb (*Rheum rhabarbarum* L.) stalks. J Agr Food Chem. 2003;51:6530-6536.
39. Miyazawa M. Nagai S. & Oshima T. Volatile Components of the Straw of *Oryza sativa* L. J Oleo Sci. 2008;57:139-143.
40. Viegas MC. & Bassoli DG. Utilização do índice de retenção linear para caracterização de compostos voláteis em café solúvel utilizando GC-MS e coluna HP-Innowax. Quím Nova. 2007;30:2031-2034.

Table 1. The fatty acid composition of *Gypsophila laricina* Schreb.

RI ¹	Compound	Mean(%) ²	Id. Met. ³
1299	Dodecanoic Acid ME (Lauric acid)	0.3	RI, MS
1499	Tetradecanoic Acid ME (Myristic acid)	1.2	RI, MS
1678	(Z)-9-Hexadecenoic Acid ME* (Palmitoleic acid)	0.6	RI, MS
1699	Hexadecanoic Acid ME (Palmitic acid)	13.2	RI, MS
1867	(Z,Z)-9,12-Octadecadienoic Acid ME* (Linoleic acid)	40.4	RI, MS
1873	(Z)-9-Octadecenoic Acid ME* (Oleic acid)	35.0	RI, MS
1899	Octadecanoic Acid ME (Stearic acid)	2.3	RI, MS
1984	(Z)-11-Eicosenoic Acid ME (Gonioic acid)	1.2	RI, MS
1999	Eicosanoic Acid ME (Arachidic acid)	3.4	RI, MS
2299	Docosanoic Acid ME (Behenic acid)	1.5	RI, MS
	Total saturated acid	21.8	
	Total unsaturated acid	77.2	
	Total	98.9	
	Unsaturated/saturated	3.6	

* Fatty acids with *cis* (Z) configuration, ME: Methyl ester,

¹RI: Retention indices,

²The results of the analysis,

³Identification method: RI: identification based on the retention times (RI) of genuine compounds on the HP Innowax column and the literature data; MS: identification based on MS comparison with the database or the literature data.

Table 2. The essential oil composition of *Gypsophila laricina* Schreb.

No	RRI ¹	RRI Lit. ²	Compound	Mean (%) ³	Id. Met. ⁴	Lit. ⁵
1	1233	1244	2-pentyl furan	0.27	RI, MS	20
2	1397	1399	Nonanal	0.29	RI, MS	20
3	1400	1400	Tetradecane	0.16	RI, MS, Ac	
4	1442	1443	Dimethyl-tetradecane	0.03	RI, MS	27
5	1499	1505	Dihydroedulan II	0.15	RI, MS	27
6	1502	1500	Pentadecane	0.11	RI, MS, Ac	
7	1504	1505	Decanal	0.17	RI, MS	28
8	1510	1516	Theaspirane B	0.7	RI, MS	28
9	1525	1532	Camphor	0.04	RI, MS	22
10	1529	1535	Dihydroedulan I	0.14	RI, MS	28
11	1543	1545	(E)-2-nonenal	0.12	RI, MS	28
12	1549	1553	Theaspirane A	0.64	RI, MS	27
13	1558	154	1-Tetradecene	0.08	RI, MS	28
14	1602	1600	Hexadecane	0.29	RI, MS, Ac	
15	1622	1638	β -cyclocitral	0.13	RI, MS	28
16	1635	1644	Thujopsene	0.04	RI, MS	32
17	1652	1655	(E)-2-decanal	0.25	RI, MS	28
18	1660	1664	Nonanol	0.1	RI, MS	28
19	1693	1685	6,10-dimethyl-2-undecanone	0.1	RI, MS	39
20	1702	1700	Heptadecane	0.28	RI, MS, Ac	

21	1717	1722	Dodecanal	0.29	RI, MS	28
22	1761	1763	Naphthalene	0.32	RI, MS	28
23	1775	1779	(<i>E,Z</i>)-2,4-Decadienal	0.13	RI, MS	28
24	1804	1779	Octadecane	0.21	RI, MS, Ac	

Table 2. Continued

No	RRI ¹	RRI Lit. ²	Compound	Mean (%) ³	Id. Met. ⁴	Lit. ⁵
25	1824	1827	(<i>E,E</i>)-2,4-decadienal	0.4	RI, MS	28
26	1831	1823	(<i>E</i>)- α -Damascenone	0.2	RI, MS	20
27	1836	1838	(<i>E</i>)- β -Damascenone	0.36	RI, MS	28
28	1865	1864	(<i>E</i>)-Geranyl acetone	1.2	RI, MS	28
29	1879	1871	Undecanol	0.17	RI, MS	33
30	1886	1864	<i>p</i> -Cymene-8-ol	0.6	RI, MS	28
31	1931	1933	Tetradecanal	0.8	RI, MS	28
32	1953	1958	(<i>E</i>)- β -Ionone	1.03	RI, MS	28
33	1968	1973	Dodecanol	0.63	RI, MS	28
34	2002	2000	Eicosane	0.29	RI, MS, Ac	
35	2005	2007	Caryophyllene oxide	0.29	RI, MS	23
36	2037	2033	Pentadecanal	0.26	RI, MS	21
37	2043	2050	(<i>E</i>)-Nerolidol	0.05	RI, MS	24
38	2051	2056	13-Tetradecanolide	0.35	RI, MS	37
39	2135	2131	Hexahydro farnesyl acetone	1.65	RI, MS	21
40	2138	2142	Spathulenol	0.05	RI, MS	20
41	2145	2136	Hexadecanal	0.3	RI, MS	27
42	2170	2192	Nonanoic acid	0.5	RI, MS	22
43	2276	2282	Decanoic acid	1.03	RI, MS	20
44	2304	2300	Tricosane	0.55	RI, MS, Ac	

45	2315	2315	2,4-bis(<i>tert</i> -butyl)phenol	0.35	RI, MS	40
46	2354	2353	Octadecanal	0.28	RI, MS	36
47	2382	2384	Farnesyl acetone	1.41	RI, MS	20
48	2407	2400	Tetracosane	0.31	RI, MS, Ac	

Table 2. Continued

No	RRI ¹	RRI Lit. ²	Compound	Mean (%) ³	Id. Met. ⁴	Lit. ⁵
49	2448	2471	Nonadecanal	0.2	RI, MS	30
50	2488	2492	Dodecanoic acid	3.51	RI, MS	20
51	2508	2500	Pentacosane	1.4	RI, MS, Ac	
52	2585	2582	Eicosanal	2.07	RI, MS	30
53	2590	2617	Tridecanoic acid	0.23	RI, MS	28
54	2606	2600	Hexacosane	0.3	RI, MS, Ac	
55	2615	2614	Phytol	1.6	RI, MS	20
56	2671	2676	Heneicosanal	1.97	RI, MS	30
57	2701	2704	Tetradecanoic acid	4.7	RI, MS	21
58	2708	2700	Hexacosane	0.7	RI, MS, Ac	
59	2775	2783	Docosanol	0.31	RI, MS	30
60	2795	2800	Octacosane	0.25	RI, MS, Ac	
61	2803	2809	Pentadecanoic acid	1.4	RI, MS	20
62	2833	2857	Palmito-γ-lactone	0.21	RI, MS	37
63	2921	2931	Hexadecanoic acid	27.03	RI, MS	25
64	2982	2990	Docosanal	0.22	RI, MS	30
65	3108	3100	Hentriacontane	12.63	RI, MS, Ac	
			Total	76.0		

In addition to the above data, diisobutyl phthalate is a common plasticizer contaminant and it was detected as a considerable component as 2.15 percentage for *G. laricina* Schreb.

¹RRI(FID): Relative retention time indices calculated against *n*-alkanes (C5-C30) in FID chromatograms

²RRI Lit.: Relative retention time given in the literature for the compound in similar columns and analysis conditions.

³The result of the analysis in FID chromatograms.

⁴Identification method: RI: identification based on the relative retention times (RRI) of genuine compounds on the HP Innnowax column and the literature data; MS: identification based on MS comparison with the database or the literature data, Ac: Identification is done according to RRI and MS values of the authentic compounds.

Lit.⁵: Literature

Uncorrected proof