Flavonoid Glycosides From *Heracleum pastinaca* Fenzl.

Perihan Gürbüz

Department Of Pharmacognosy, Erciyes University Faculty Of Pharmacy, Kayseri, Turkey

Abstract

INTRODUCTION: The objective is the study to isolate and characterize the secondary metabolites of *Heracleum pastinaca* which has not been previously investigated.

METHODS: Conventional chromatographic procedures were carried out for isolation of the compounds. The structures of the compounds were elucidated by extensive 1D and 2D NMR spectroscopic analysis in combination with MS experiments and comparing with relevant literature data.

RESULTS: The first phytochemical investigation on the whole parts of *Heracleum pastinaca* Fenzl. led to the isolation and identification of seven known flavonoid glycosides; isoquercetin (1), rutin (2), afzelin (3), astragalin (4), isorhamnetin 3-O-glucoside (5), nicotiflorine (6) and narcissoside (7).

DISCUSSION AND CONCLUSION: This is the first report on the isolation of these flavonoid glycosides from *H. pastinaca* and compounds 3, 5, 6 and 7 from genus *Heracleum*.

Keywords: *Heracleum*, Apiaceae, isorhamnetin, flavonoid glycosides, chemotaxonomy

INTRODUCTION

The genus *Heracleum* which is known as “hogweed,” is one of the largest genera of Apiaceae containing more than 120 species widely distributed in Central Europe and Asia as well as 17 species with 41% endemism in flora of Turkey. ¹⁻²

Heracleum species have been traditionally used as spice and food additive as well as in the treatment of inflammation, flatulence, stomachache, epilepsy and psoriasis. They also act as carminative, antiseptic, antimicrobial, analgesic and anticonvulsant agents. ³ Some *Heracleum* species are used traditionally for different purposes i.e. *H. crenatifolium* as vegetable and condiment ⁴; *H. trachyloma* against asthma and bronchitis ⁵; *H. spondylium* L. subsp. *ternatum* as galactagogue ⁶; *H. persicum* and *H. platytaenium* for gastritis, epilepsy and as sedative ⁷ in Turkey.

There are plenty of phytochemical studies on *Heracleum* species mainly focused on furanocoumarins⁸⁻⁹ and furanocoumarin glycosides¹⁰⁻¹¹ together with alkaloids¹², polyacetylenes¹³ and flavonoids.¹⁴⁻¹⁶ Essential oil compounds of the genus were also studied.¹⁷⁻¹⁹ Although the phytochemical studies and bioactivity studies are mostly conducted with the coumarin compounds of the genus, we would like to examine the flavonoid content of *H. pastinaca* Fenzl. (Figure 1) which is a rare endemic tiny plant, mainly distributed in the inner and the southwest region of Anatolia.²⁰

Conventional chromatographic purification procedures were carried out to isolate the compounds of the *H. pastinaca*. The structures of the compounds were elucidated by extensive 1D and 2D NMR and ESI-MS experiments confirmed by the relevant literature data. Chemotaxonomic significance of the compounds was discussed.

EXPERIMENTAL

General

NMR (400 MHz for ¹H NMR, 100 MHz for ¹³C NMR, both use TMS as internal standard) were measured on a Bruker AM 400 spectrometer and MS spectra on a LC/MS Shimadzu 8040 instrument. Kieselgel 60 (Merck, 0.063-0.200 mm) was used for open column chromatography (CC). Sephadex LH-20 (SP LH-20) (General Electrics Healthcare) was used for Gel Permeation Chromatography (GPC). LiChroprep C₁₈ (Merck, 40-63 µm) was used for Medium Pressure Liquid Chromatography (MPLC) (Buchi Pump Module: C-601, UV-Photometer: C-640, control Unit: C-620, Fr. Collector: C-660). TLC analyses were carried out on pre-coated Kieselgel 60 F₂₅₄ aluminum plates (Merck). Compounds were detected by UV fluorescence and spraying 1% vanillin/H₂SO₄, followed by heating at 100 °C for 1-2 min.

Plant material

Whole parts (aerial parts & roots) of *H. pastinaca* were collected from Maden- Kızıltepe region (Niğde-Ulukışla), at about 2600 m altitude of calcareous rock clefts on August 2017. A voucher specimen was deposited at the Herbarium of Hacettepe University, Faculty of Pharmacy under the code HUEF-17015.

Extraction and isolation

The dried and powdered whole parts of *H. pastinaca* (90 g) was extracted with MeOH (500 mL x 4) at 37 °C. After the evaporation of the solvent (yield 18%), the crude MeOH extract (17 g) was first dissolved in water than partitioned between *n*-hexane and *n*-BuOH respectively. *n*-BuOH (4.5 g) extract was first submitted to column chromatography on

Sephadex LH-20 (2,5 x 60 cm) and eluted with MeOH. Four main fractions (Fr. 1 (1.7 g); Fr. 2 (2.2 g), Fr. 3 (332.6 mg), Fr. 4 (130 mg) were obtained. Fr. 3 (332.6 mg) was submitted to reverse phase column (1.5 cm x 15 cm) and eluted with gradient H₂O:MeOH solvent system (10%→ 50%; 10 mL/min; 4-5 mbar) with the MPLC system coupled with a fraction collector to give four sub fractions (Fr.3 a-d).

Fr. 3d gave compound **3** (4 mg). Further purification of Fr. 3a (116 mg) with reverse phase column (1.5 cm x 15 cm) and eluted with gradient H₂O:MeOH solvent system (20%→ 30%; 10 mL/min; 4-5 mbar) yielded two sub fractions. These two fractions were submitted to TLC plate (20x20 cm) separately and eluted with 70:30:3 (CHCl₃: MeOH: H₂O) to yield compounds **1** (22 mg) and compound **2** (32 mg) respectively. Fr. 3c (63 mg) was submitted to two different TLC plates (20x20 cm) and eluted with 70:30:3 (CHCl₃: MeOH: H₂O). After elution, the bands that belong to the compounds were detected under UV₂₅₄ light and scraped to obtain compounds **4** and **5** (24 mg) and compounds **6** and **7** (16 mg) respectively as mixtures.

Structure elucidation

The structures of the compounds (Figure 2) were elucidated by 1D and 2D NMR experiments. The positions of the sugar units are confirmed by 2D HMBC experiments. Together with ESI-MS data and comparison with relevant literature the compounds are elucidated as follows; isoquercetin (**1**)²¹, rutin (**2**)²¹⁻²², afzelin (**3**)²³, astragalin (**4**)^{21, 24}, isorhamnetin 3-*O*- β -glucopyranoside (**5**)²⁵, kaempferol 3-*O*-rutinoside (**6**)²¹⁻²² and isorhamnetin 3-*O*-rutinoside (**7**).²²

Quercetin 3-*O*- β -glucopyranoside (Isoquercetin) (**1**)

Yellow powder; Negative ESI/MS *m/z*: 463 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 7.72 (d, *J*=2.1 Hz, 1H, H-2'), 7.58 (dd, *J*=8.5, 2.1 Hz, 1H, H-6'), 6.86 (d, *J*=8.5 Hz, 1H, H-5'), 6.24 (d, *J*=2.0 Hz, 1H, H-8), 6.08 (d, *J*=2.0 Hz, 1H, H-6), 5.11 (d, *J*=7.6 Hz, 1H, H-1''), 3.71 (dd, *J*=11.8, 2.3 Hz, 1H, H-6a''), 3.59 (dd, *J*=12.5, 4.6 Hz, 1H, H-6b''), 3.56–3.17 (m, 4H, remaining sugar signals).

Quercetin 3-*O*- α -rhamnopyranosyl (1→6)- β -glucopyranoside (Rutin) (**2**)

Yellow powder; Negative ESI/MS *m/z*: 609 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 7.68 (brs, 1H, H-2'), 7.64 (brd, *J*=8.0 Hz, 1H, H-6'), 6.87 (d, *J*=8.2 Hz, 1H, H-5'), 6.30 (brs, 1H, H-

8), 6.13 (brs, 1H, H-6), 5.03 (d, $J=7.6$ Hz, 1H, H-1''), 4.53 (brs, 1H, H-1'''), 3.81 (brd, $J=10.5$ Hz, 1H, H-6a''), 3.70-3.20 (m, 9H, remaining sugar signals), 1.15 (d, $J=6.2$ Hz, 3H).

Kaempferol 3-*O*- α -rhamnopyranoside (Afzelin) (3)

Yellow powder; Negative ESI/MS m/z : 431 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 7.78 (d, $J=8.9$ Hz, 2H, H-2', 6'), 6.95 (d, $J=8.8$ Hz, 2H, H-3', 5'), 6.39 (d, $J=2.1$ Hz, 1H, H-8), 6.21 (d, $J=2.1$ Hz, 1H, H-6), 5.38 (d, $J=1.6$ Hz, 1H, H-1''), 4.23 (dd, $J=3.3, 1.6$ Hz, 1H, H-2''), 3.76 – 3.68 (m, 1H, H-3''), 3.53 – 3.40 (m, 2H, H-4'', 5''), 0.93 (d, $J=5.7$ Hz, 3H, H-6'').

Kaempferol 3-*O*- β -glucopyranoside (Astragalin) (4)

Yellow powder; Negative ESI/MS m/z : 447 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.04 (d, $J=8.8$ Hz, 2H, H-2', 6'), 6.89 (d, $J=8.2$ Hz, 2H, H-3', 5'), 6.24 (d, $J=1.9$ Hz, 1H, H-8), 6.09 (d, $J=1.9$ Hz, 1H, H-6), 5.27 (d, $J=7.3$ Hz, 1H, H-1''), 3.76 – 3.57 (m, 2H, H-6''), 3.56 – 3.16 (m, 4H, remaining sugar signals).

Isorhamnetin 3-*O*- β -glucopyranoside (5)

Yellow powder; Negative ESI/MS m/z : 477 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 7.91 (d, $J=1.9$ Hz, 1H, H-2'), 7.59 (dd, $J=8.5, 1.9$ Hz, 1H, H-6'), 6.89 (d, $J=8.5$ Hz, 1H, H-5'), 6.24 (d, $J=1.9$ Hz, 1H, H-8), 6.09 (d, $J=1.9$ Hz, 1H, H-6), 5.10 (d, $J=7.4$ Hz, 1H, H-1''), 3.94 (s, 3H, OCH₃), 3.76–3.57 (m, 2H, H-6''), 3.56–3.16 (m, 4H, remaining sugar signals).

Kaempferol 3-*O*-rutinoside (Nicotiflorin) (6)

Yellow powder; Negative ESI/MS m/z : 593 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.06 (d, $J=8.8$ Hz, 2H, H-2', 6'), 6.89 (d, $J=8.0$ Hz, 2H, H-3', 5'), 6.28 (brs, 1H, H-8), 6.12 (d, $J=1.8$ Hz, 1H, H-6), 5.14 (d, $J=7.3$ Hz, 1H, H-1''), 4.51 (brs, 1H, H-1'''), 3.86–3.62 (m, 2H, H-6''), 3.61–3.22 (m, 8H, remaining sugar signals), 1.15 (d, 6.2 Hz, 3H, H-6''').

Isorhamnetin 3-*O*-rutinoside (Narcissoside) (7)

Yellow powder; Negative ESI/MS m/z : 623 [M-H]⁻; ¹H NMR (400 MHz, MeOH-*d*₄) δ 7.95 (d, $J=1.8$ Hz, 1H, H-2'), 7.62 (dd, $J=8.5, 1.8$ Hz, 1H, H-6'), 6.89 (d, $J=8.0$ Hz, 1H, H-5'), 6.28 (brs, 1H, H-8), 6.12 (d, $J=1.8$ Hz, 1H, H-6), 5.02 (d, $J=7.3$ Hz, 1H, H-1''), 4.52 (brs, 1H, H-1'''), 3.95 (s, 3H, OCH₃), 3.86–3.62 (m, 2H, H-6''), 3.61–3.22 (m, 8H, remaining sugar signals), 1.12 (d, 6.2 Hz, 3H, H-6''').

RESULTS AND DISCUSSION

The present work reports for the first time the characterization of seven flavonoid glycosides **1-7** from the whole parts of *H. pastinaca*. To best of our knowledge, this is the first report of compounds **3**, **5**, **6** and **7** from the genus *Heracleum* while others were reported from different *Heracleum* species. i.e. isoquercetin (**1**) from *H. napalense*²⁶ and *H. mollendorffii*¹⁵; astragalin from *H. mollendorffii*¹⁵ and rutin from *H. sphondylium*²⁷⁻²⁸ before. The presence of flavonoids in higher plants has been associated with various environmental conditions such as; high-light/UV-stress, cold stress, nutritional deficiencies and pathogen protection etc.²⁹⁻³¹

The habitat of the samples were about 2600 m altitudes where the plants were exposed to a high UV radiation. This fact should effect the production of different type and quantities of flavonoids of the plant. The phytochemical investigation of the *Heracleum* species are mostly focused on the linear and angular type furanocoumarins and different biological activities of the genus such as, insecticidal, antibacterial, antiviral and antifungal were may be attributed to the these coumarin type compounds.³ There are limited phytochemical studies about the isolation of the flavonoids from *Heracleum* species. Few flavonoids i.e. kaempferol, quercetin, isorhamnetin¹⁶ rutin²⁸, astragalin¹⁵, flavantaside and epirutin³² were reported from different *Heracleum* species. In this study, the isolated and elucidated flavonols are mainly kaempferol, quercetin and isorhamnetin glycosides. Flavonoids possesses many important biological activities such as antimicrobial³³, antioxidant³⁴, antiviral³⁵ etc. The presence of those valuable flavonoids in *Heracleum* species definitely enriches the chemical diversity and provides evidence for the chemotaxonomic studies of *Heracleum* species and the family Apiaceae as well.

CONCLUSION

The first phytochemical study of *Heracleum pastinaca* led to the isolation and structure identification of seven flavonoid glycosides. The structure of isolated compounds were elucidated by 1D and 2D NMR analyses, together with ESI-MS data and comparison with relevant literature data; isoquercetin (**1**)²¹, rutin (**2**)²¹⁻²², afzelin (**3**)²³, astragalin (**4**)^{21, 24}, isorhamnetin 3-*O*- β -glucopyranoside (**5**)²⁵, nicotiflorin (**6**)²¹⁻²² and narcissoside (**7**).²² Notably this is the first report of these flavonol glycosides from *H. pastinaca* and compounds **3**, **5**, **6** and **7** from the genus *Heracleum*. In conclusion, when considering the relationship between the bioactivities and the chemistry of *Heracleum* species; it should be a possible fact that; flavonoids can also play an important role to contribute the bioactivity and traditional uses of the *Heracleum* species.

ACKNOWLEDGEMENTS

The author is indebted to Dr. Şengül Dilem Doğan (Erciyes University) for helping structure elucidation and Dr. Ahmet Savran for identification of the plant material and for providing the high-resolution photo of the plant.

Conflict of Interest: No conflict of interest was declared by the authors.

Uncorrected proof

REFERENCES

1. The Plant List (2013). Version 1.1. Published on the Internet; <http://www.theplantlist.org/>. (accessed 1st January).
2. <http://www.bizimbitkiler.org/Heracleum>. (accessed 1st January).
3. Bahadori MB, Dinparast L, Zengin G. The genus *Heracleum*: a comprehensive review on its phytochemistry, pharmacology, and ethnobotanical values as a useful herb. *Compr Rev Food Sci Food Saf*. 2016;15:1018-1039.
4. Özgökçe F, Özçelik H. Ethnobotanical aspects of some taxa in East Anatolia, Turkey. *Econ Bot*. 2004;58:697.
5. Polat R, Cakilcioglu U, Satıl F. Traditional uses of medicinal plants in Solhan (Bingöl-Turkey). *J Ethnopharmacol*. 2013;148:951-963.
6. Kültür Ş. Medicinal plants used in Kırklareli province (Turkey). *J Ethnopharmacol*. 2007;111:341-364.
7. Yıldırım B, Terzioglu Ö, Özgökçe F, Türközü D. Ethnobotanical and pharmacological uses of some plants in the districts of Karpuzalan and Adıgüzel (Van-Turkey). *J Anim Vet Adv*. 2008;7:873-878.
8. Razdan TK, Kachroo V, Harkar S, Koul GL. Furanocoumarins from *Heracleum canescens*. *Phytochemistry*. 1982;21:923-927.
9. Steck W. Leaf furanocoumarins of *Heracleum lanatum*. *Phytochemistry*. 1970;9:1145-1146.
10. Fischer FC, Jasperse PH, Karlsen J, Svendsen AB. Ein neues furanocumaringlykosid aus *Heracleum mantegazzianum*. *Phytochemistry*. 1974;13:2334-2335.
11. Mi J, Peng Y, Zhang H, Wang X, Huo Y, Wang Z, Gao Y, Zhang H. A new benzofuran derivative glycoside and a new coumarin glycoside from roots of *Heracleum dissectum* Ledeb. *Med Chem Res*. 2017;1-6.
12. Gupta BD, Banerjee SK, Handa KL. Alkaloids and coumarins of *Heracleum wallichii*. *Phytochemistry*. 1976;15:576.
13. Nakano Y, Matsunaga H, Saita T, MORI M, KATANO M, OKABE H. Antiproliferative Constituents in Umbelliferae Plants II.: Screening for Polyacetylenes in Some Umbelliferae Plants, and Isolation of Panaxynol and Falcarindiol from the Root of *Heracleum moellendorffii*. *Biol Pharm Bull*. 1998;21:257-261.
14. Ghodsi B. Flavonoids of three *Heracleum* species: *H. persicum* L., *H. sphondylium* L. and *H. montanum* Schl. *Bull Trav Soc Pharm Lyon*. 1976;20:3-8.

15. Park H-J, Nugroho A, Jung B-R, Won Y-H, Jung Y-J, Kim W-B, Choi J-S. Isolation and Quantitative Analysis of Flavonoids with Peroxynitritescavenging Effect from the Young Leaves of *Heracleum moellendorffii*. Korean J Plant Resour. 2010;23:393-398.
16. Siwon J, Karlsen J. The isolation and identification of flavonoid aglycones from *Heracleum mantegazzianum* Somm. et Lev. Medd Nor Farm Selsk. 1976;38:11-12.
17. Skalicka-Woźniak K, Grzegorzczak A, Świątek Ł, Walasek M, Widelski J, Rajtar B, Polz-Dacewicz M, Malm A, Elansary HO. Biological activity and safety profile of the essential oil from fruits of *Heracleum mantegazzianum* Sommier & Levier (Apiaceae). Food Chem Toxicol. 2017;109:820-826.
18. Firuzi O, Asadollahi M, Gholami M, Javidnia K. Composition and biological activities of essential oils from four *Heracleum* species. Food Chem. 2010;122:117-122.
19. Iscan G, Ozek T, Ozek G, Duran A, Baser K. Essential oils of three species of *Heracleum*. Anticandidal activity. Chem Nat Comp. 2004;40:544-547.
20. Babaç MT. Possibility of an information system on plants of South-West Asia with particular reference to the Turkish plants data service (TUBIVES). Turk J Botany. 2004;28:119-127.
21. Han J-T, Bang M-H, Chun O-K, Kim D-O, Lee C-Y, Baek N-I. Flavonol glycosides from the aerial parts of *Aceriphyllum rossii* and their antioxidant activities. Arch Pharm Res. 2004;27:390-395.
22. Chaurasia N, Wichtl M. Flavonolglykoside aus *Urtica dioica* 1, 2. Planta Med. 1987;53:432-434.
23. Lee SY, So Y-J, Shin MS, Cho JY, Lee J. Antibacterial effects of afzelin isolated from *Cornus macrophylla* on *Pseudomonas aeruginosa*, a leading cause of illness in immunocompromised individuals. Molecules. 2014;19:3173-3180.
24. Wei Y, Xie Q, Fisher D, Sutherland IA. Separation of patuletin-3-O-glucoside, astragalin, quercetin, kaempferol and isorhamnetin from *Flaveria bidentis* (L.) Kuntze by elution-pump-out high-performance counter-current chromatography. J Chromatog A. 2011;1218:6206-6211.
25. Lee YS, Lee HS, Shin KH, Kim B-K, Lee S. Constituents of the halophyte *Salicornia herbacea*. Arch Pharm Res. 2004;27:1034-1036.
26. Dash S, Nath LK, Bhise S. Antioxidant and antimicrobial activities of *Heracleum nepalense* D Don root. Trop J Pharm Res. 2005;4:341-347.

27. Benedec D, Hanganu D, Filip L, Oniga I, Tiperciuc B, Olah N-K, Gheldiu A-M, Raita O, Vlase L. Chemical, antioxidant and antibacterial studies of Romanian *Heracleum sphondylium*. *Farmacia*. 2017;65:252-256.
28. Harborne J, Williams CA. Flavonoid patterns in the fruits of the Umbelliferae. *Phytochemistry*. 1972;11:1741-1750.
29. Dixon RA, Paiva NL. Stress-induced phenylpropanoid metabolism. *Plant Cell*. 1995;7:1085.
30. Kusano M, Tohge T, Fukushima A, Kobayashi M, Hayashi N, Otsuki H, Kondou Y, Goto H, Kawashima M, Matsuda F. Metabolomics reveals comprehensive reprogramming involving two independent metabolic responses of *Arabidopsis* to UV-B light. *Plant J*. 2011;67:354-369.
31. Roberts MR, Paul ND. Seduced by the dark side: integrating molecular and ecological perspectives on the influence of light on plant defence against pests and pathogens. *New Phytologist*. 2006;170:677-699.
32. Komissarenko N, Satsyperova I. Flavonoids and coumarins of leaves of *Heracleum antasiaticum* Manden. *Rastit Resur*. 1974.
33. Cushnie TT, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents*. 2005;26:343-356.
34. Burda S, Oleszek W. Antioxidant and antiradical activities of flavonoids. *J Agr Food Chem*. 2001;49:2774-2779.
35. Kaul TN, Middleton E, Ogra PL. Antiviral effect of flavonoids on human viruses. *J Med Vir*. 1985;15:71-79.

Figure 1

	R ₁	R ₂
1	Glucose	OH
2	Rutinose	OH
3	Rhamnose	H
4	Glucose	H
5	Glucose	OCH ₃
6	Rutinose	H
7	Rutinose	OCH ₃

Figure 2. Isolated flavonoids from *H. pastinaca*