Evaluation of Depression and Anxiety of Patients with Multiple Sclerosis During the COVID-19 Pandemic: A Comparison with the General Population

COVID-19 Döneminde Multipl Sklerozlu Hastaların Anksiyete ve Depresyon Düzeylerinin Değerlendirilmesi: Genel Popülasyon ile Karşılaştırılması

© Gizem Yağmur Yalçın¹, © Cansu Polat Dünya¹, © Zeliha Tülek¹, © Murat Kürtüncü², © Mefkure Eraksoy²

¹Florence Nightingale Faculty of Nursing; Istanbul University-Cerrahpasa, Istanbul, Turkey

²Istanbul University, Istanbul Faculty of Medicine, Department of Neurology, Istanbul, Turkey

Abstract

Objective: The coronavirus disease-2019 (COVID-19) pandemic can increase anxiety and depression in patients with multiple sclerosis (PwMS). This study aimed to evaluate the predictors of anxiety and depression in PwMS during the COVID-19 pandemic and compare it with the general population (GP).

Materials and Methods: The study was conducted in an MS clinic in Istanbul University Istanbul Faculty of Medicine. Data were collected online using a form including sociodemographics and questions on COVID-19-related knowledge and the hospital anxiety and depression scale.

Results: This study included 800 participants (GP: 421, MS: 379). Anxiety (42.3% vs. 32.2%, $p=0.002$) and depression (53.9% vs. 39.6%, $p<0.001$) were more common, COVID-19-related knowledge was lower (9.1 ± 1.5 vs. 9.9 ± 1.2 , $p<0.001$), and rate of COVID-19 infection was higher (15.4% vs. 6.9%, $p<0.001$) in the GP than PwMS. Among other factors, COVID-19-related knowledge was a common predictor of anxiety and depression in both GP and MS groups [odds ratio (OR): 0.8, 95% confidence interval (CI) 0.7-0.9, $p=0.007$ and OR: 0.7, 95% CI 0.7-0.9, $p=0.020$, for anxiety; OR: 0.7, 95% CI 0.7-0.9, $p=0.001$ and OR: 0.8, 95% CI 0.7-1.0, $p=0.023$ for depression, respectively].

Conclusion: This study revealed that PwMS have less anxiety and depression than the GP. Our study shows the importance of COVID-19-related knowledge and the need for psychological support during the pandemic.

Keywords: Multiple sclerosis, COVID-19, anxiety, depression, COVID-19-related knowledge

Öz

Amaç: Koronavirüs hastalığı-2019 (COVID-19) pandemisi multipl skleroz (MS) hastalarında anksiyete ve depresyonu artırabilir. Bu çalışmanın amacı COVID-19 pandemisi sürecinde MS hastalarında anksiyete ve depresyonu değerlendirmek, yordayıcılarını belirlemek ve genel popülasyon (GP) ile karşılaştırmaktır.

Gereç ve Yöntem: Çalışma İstanbul Üniversitesi İstanbul Tıp Fakültesi MS kliniğinde yürütüldü. Veriler, sosyodemografik özellikleri, COVID-19 bilgi düzeyini ve hastane anksiyete ve depresyon ölçeğine ilişkin soruları içeren bir form ile online olarak toplandı.

Bulgular: Çalışmaya 800 katılımcı (GP: 421, MS: 379) dahil edildi. Anksiyete (GP: %42,3 vs. MS: %32,2, $p=0.002$) ve depresyon (GP: %53,9 vs. MS: %39,6, $p<0,001$) MS hastalarında genel popülasyona kıyasla daha düşük bulundu. COVID-19 bilgisi ise daha yüksekti (GP: $9,1\pm1,5$ 'e vs. MS: $9,9\pm1,2$, $p<0,001$). COVID-19 enfeksiyonu geçirme oranı daha düşüktü (GP: %15,4'e vs. %6,9, $p<0,001$). Diğer faktörlerin yanı sıra, COVID-19 hakkında bilgi düzeyi hem GP hem de MS gruplarında anksiyete ve depresyonun ortak bir yordayıcısıdır [anksiyete için odds oranı (OR): 0,8, %95 güven aralığı (GA): 0,7-0,9, $p=0,007$ ve OR: 0,7, %95 GA: 0,7-0,9, $p=0,020$, depresyon için; OR: 0,7, %95 GA: 0,7-0,9, $p=0,001$ ve OR: 0,8, %95 GA: 0,7-1,0, $p=0,023$].

Sonuç: Çalışmamız MS hastalarının GP'ye göre daha az anksiyete ve depresyona sahip olduğunu ortaya koydu. Bu çalışma pandemi sürecinde COVID-19'a ilişkin bilgi düzeyinin önemini ve psikolojik destek ihtiyacının gerekliliğini göstermektedir.

Anahtar Kelimeler: Multipl skleroz, COVID-19, anksiyete, depresyon, COVID-19 bilgi düzeyi

Address for Correspondence/Yazışma Adresi: Gizem Yağmur Yalçın Nurse, PhD Student, Florence Nightingale Faculty of Nursing; Istanbul University-Cerrahpasa, Istanbul, Turkey Phone: +90 530 230 38 22 E-mail: gvizviz@gmail.com ORCID: orcid.org/0000-0001-8398-9181

Received/Geliş Tarihi: 06.10.2021 **Accepted/Kabul Tarihi:** 10.12.2021

©Copyright 2021 by Turkish Neurological Society
Turkish Journal of Neurology published by Galenos Publishing House.

Introduction

Coronavirus disease-2019 (COVID-19) affected and killed millions of people in a short time after its emergence in China and the World Health Organization declaration as a pandemic (1). Turkey is among the most affected countries by COVID-19, with 6,987,494 cases and 62,745 deaths as of September 2021 (2). In our society, where the epidemic had got out of control, public health practices, such as individual measures, quarantine, lockdown, and remote work, became inevitable. The epidemic threatened society in terms of personal health, loss of close relatives, social and occupational life interruption, and economic burden. This threat had negative psychological impacts (fear, stress, insomnia, anxiety, and depression). These psychological impacts have been reported higher among populations with chronic diseases and high-risk groups (3).

Like other individuals with a chronic disease, patients with multiple sclerosis (PwMS) are known to be at risk for anxiety and depression (4). Additionally, the unknown course of immunosuppressed patients with COVID-19, isolation from daily life, and lack of social support due to the lockdown worsened the anxiety and depression of patients (4,5). The COVID-19 pandemic has become a severe stressor in PwMS due to insufficient information at the initial and misinformation in the later stages. Increasing the awareness and elimination of the misinformation may decrease the risks of patients' anxiety and depression.

This study aimed to assess the predictors of anxiety and depression during the COVID-19 pandemic and the effect of COVID-19 awareness on the psychological state of patients compared to the general population (GP).

Material and Methods

Research Design, Sample, and Setting

This case-control study was performed in PwMS who were followed up in the outpatient MS clinic at the Istanbul University, Istanbul Faculty of Medicine. Our study included 379 patients who are older than 18 years without a relapse in the last month. The control group consisted of 421 healthy individuals.

Data Collection

The data were collected using an online survey (Google forms) between October 2020 and February 2021. Participants were asked to complete the questionnaire, which took approximately 15 min, via e-mail or other communication applications. A snowball sampling strategy was used to access the healthy controls.

Data Collection Tools

An online questionnaire and hospital anxiety and depression scale (HADS) were employed. The questionnaire included sociodemographic and clinical characteristics (duration of diagnosis, drugs used, etc.), questions regarding COVID-19 infection status, and lifestyle changes (medication use, occupation, daily physical activities, sleep, nutrition, etc.). Participants were also assessed for COVID-19-related knowledge (transmission route, protection, etc.). This part of the questionnaire was prepared in line with the COVID-19 Guidelines of the Ministry of Health. Each correct answer was evaluated as one point, and each incorrect or blank answer was evaluated as zero. The total score of the test was calculated by summing the item scores.

The HADS was developed by Zigmond and Snaith (6) and adapted to Turkish by Aydemir et al. (7) This is a 14-item self-assessment four-points Likert scale that evaluates anxiety and depression. The scores range between 0 and 21. The cut-off score is 10 for anxiety and 7 for depression in the Turkish population.

Ethical Aspect of the Study

The ethical principle compliance was evaluated by the ethics committee of the institution where the study was conducted, and the Ethics Committee approval was obtained (Istanbul University, Istanbul Faculty of Medicine, Department of Neurology, Istanbul, Turkey, file number: 2020/1095, no: 29624016-050.99- 1215). Consent to participate was obtained using online forms.

Statistical Analysis

The Statistical Package for the Social Sciences (v 21.0, IBM) software was used for statistical data analysis. Data were analyzed for normality distribution using the Kolmogorov-Smirnov test and expressed as mean ($\pm$ standard deviation) for continuous variables or frequencies (n, %) for categorical variables. Independent samples t-test and One-Way analysis of variance test were used to compare continuous variables between the groups and chi-square test to compare the categorical variables. The Pearson correlation analysis was used to correlate continuous variables and multivariate logistic regression analysis to determine the associated factors with anxiety and depression. The analysis of covariance (ANCOVA) was used to adjust for group differences based on covariates.

Results

A total of 800 individuals (MS: 379, controls: 421) participated in the study. The mean age of patients was 35.8 ± 9.9 [minimum-maximum (min-max): 18-67] years, 67.5% (n=256) were female, 50.9% (n=193) were married, and 70.4% (n=267) had a university degree. The mean age of the controls was 28.6 ± 8.5 (min-max: 16-65) years, 37.8% (n=159) were female, 40.4% (n=170) were married, and 62% (n=261) had a university degree (Table 1).

The mean disease duration was 8.2 ± 6.6 (min-max: 1-34) years. During the COVID-19 pandemic, 21.4% (n=81) of the patients reported an attack, 47.5% (n=180) were admitted to the hospital due to MS, 13.7% (n=52) reported worsening in motor function, and 4.7% (n=18) reported starting antidepressants. Interestingly, 13.7% (n=52) of the patients thought to discontinue their disease-modifying therapies (DMT) and 12.9% (n=49) discontinued their DMT due to fear of getting COVID-19.

More individuals reported changes in their working conditions in the MS group compared to the GP (46.6% vs. 24.9%, $p < 0.001$). The rate of starting smoking during the pandemic was higher in the controls compared to patients (5.9% vs. 2.9%, $p = 0.039$). The number of COVID-19 infected individuals in the GP was higher than in the PwMS (15.4% vs. 6.9%, $p = 0.039$). The rate of COVID-19 deaths among relatives was higher in the GP than in PwMS (24.7% vs. 11.9%, $p < 0.001$). In both groups, during the COVID-19 pandemic, at least half of the individuals reported changes in their sleeping and eating habits, physical activity, and body weight. The most affected habit was physical activity in both groups. Furthermore, compared to PwMS, insomnia ($p < 0.001$), eating habits ($p < 0.001$), weight gain ($p < 0.001$), and decreased physical activity ($p = 0.015$) were more frequent in the GP (Table 2).

Table 1. Sociodemographic characteristics of participants (n=800)

Characteristics	Control		PwMS		χ^2	p
	n	%	n	%		
Age (Mean $\pm$ SD), (min-max)	28.6 $\pm$ 8.5	(16-65)	35.8 $\pm$ 9.9	(18-67)	11.1	<0.001*
Gender						
Female	159	37.8	256	67.5	70.8	<0.001
Male	262	62.2	123	32.5		
Education						
Primary/secondary education	32	7.6	32	8.4	9.0	0.011
High school	128	30.4	80	21.1		
University	261	62.0	267	70.4		
Income						
Less than expenses	143	34.0	86	23.1	29.4	<0.001
Equal to expenses	218	51.8	262	70.6		
More than expenses	60	14.3	31	8.4		
Marital status						
Married	170	40.4	193	50.9	41.8	<0.001
Single	246	58.4	153	40.4		
Divorced/widow	5	1.2	33	8.8		
Children						
Yes	130	30.9	167	44.1	14.8	<0.001
No	291	69.1	212	55.9		
Smoking						
Yes	154	36.6	100	26.5	9.4	0.002
No	267	63.4	279	73.5		
Antidepressant use						
Yes	15	3.6	84	22.2	63.6	<0.001
No	406	96.4	295	77.8		

*Independent samples t-test, SD: Standard deviation, PwMS: Patients with multiple sclerosis, min: Minimum, max: Maximum

The mean score of COVID-19-related knowledge was higher in patients than in the GP (9.9 $\pm$ 1.2 vs. 9.1 $\pm$ 1.5, p <0.001). Patients had better knowledge, especially on questions regarding transmission routes, whereas their knowledge on protective measures was comparable (Table 3).

Interestingly, the rate of anxiety and depression, when some confounders (age, gender, education, marital status, antidepressant use, COVID-19-related knowledge, and COVID-19 infection status) were adjusted using the ANCOVA, were higher in the GP compared to the patients (42.3% vs. 32.2% for anxiety, p =0.002; 53.9% vs. 39.6% for depression, p <0.001) (Figure 1). In line with this observation, anxiety and depression scores were higher in the controls than PwMS (8.5 $\pm$ 4.5 vs. 7.7 $\pm$ 4.5 for anxiety, p =0.005; 7.7 $\pm$ 4.0 vs. 6.4 $\pm$ 3.9 for depression, p <0.001). Additionally, when correlations among anxiety, depression, and COVID-19-related knowledge were assessed, a significant positive correlation was found between the anxiety and depression scores (r =0.63, p <0.001) and weak negative correlations between anxiety, depression, and knowledge scores (Figure 2). The anxiety and depression scores of participants are shown in Table 4.

The relationship between the patients' clinical features, anxiety, and depression scores revealed no significant correlations between the disease duration, anxiety (r =0.81, p =0.099), and depression

(r =0.01; p =0.781). As expected, the anxiety scores of individuals who started using antidepressants during the pandemic period were higher than those who did not use antidepressants (10.2 $\pm$ 5.5 vs. 7.5 $\pm$ 4.4, p =0.013). No significant difference was found in anxiety and depression scores between the patients according to hospital admission status due to worsening MS (7.4 $\pm$ 4.5 vs. 7.9 $\pm$ 4.5, p =0.271 and 6.6 $\pm$ 3.9 vs. 6.3 $\pm$ 3.9, p =0.377, respectively). Anxiety and depression scores of patients with relapse were higher than those on remission (9.1 $\pm$ 4.6 vs. 7.3 $\pm$ 4.4, p =0.028 and 8.1 $\pm$ 3.9 vs. 6.0 $\pm$ 3.8, p <0.001, respectively). The anxiety and depression scores did not differ according to the state of planning to discontinue the DMT (8.7 $\pm$ 4.2 vs. 7.5 $\pm$ 4.5, p =0.065 and 6.4 $\pm$ 3.3 vs. 6.5 $\pm$ 4.0, p =0.855, respectively).

A multivariate logistic regression analysis was performed to determine the predictive factors of anxiety and depression. The model included age, gender, education, marital status, child status, antidepressant use, COVID-19 infection status, loss of relative due to COVID-19, and COVID-19-related knowledge as covariates. The predictors of anxiety include gender (being female) [odds ratio (OR): 1.7, 95% confidence interval (CI): 1.1-2.7, p =0.015] and knowledge about COVID-19 in the GP (OR: 0.8, 95% CI: 0.7-0.9, p =0.007), and they were age (OR: 0.9 95% CI: 0.9-1.0, p =0.010), education (OR: 0.3, 95% CI: 0.1-0.9, p =0.025), antidepressant

Table 2. Lifestyle changes due to the COVID-19 pandemic

Characteristics	Control		PwMS		χ^2	p
	n	%	n	%		
COVID-19 diagnosis						
Yes	65	15.4	26	6.9	14.5	<0.001*
No	356	84.6	353	93.1		
Place of COVID-19 treatment						
Home	46	10.9	25	6.6	20.2	<0.001*
Hospitalized	19	4.5	1	0.3		
Death of close relatives due to COVID-19						
Yes	104	24.7	45	11.9	21.6	<0.001*
No	317	75.3	334	88.1		
Concern about COVID-19						
None	71	16.9	35	9.2	83.7	<0.001*
Mild	99	23.5	82	21.6		
Moderate	199	47.3	112	29.6		
High	52	12.4	150	39.6		
Following the COVID-19 chart						
Never	30	7.1	15	4.0	17.7	<0.001*
Sometimes	191	45.4	129	34.0		
Always	200	47.5	235	62.0		
Change in working conditions						
Yes	69	24.9	173	46.6	31.9	<0.001*
No	208	75.1	198	53.4		
Start smoking						
Yes	25	5.9	11	2.9	4.3	0.039
No	396	94.1	368	97.1		
Change in sleep						
None	124	29.5	183	48.3	39.7	<0.001*
Medium disruption	204	48.5	160	42.2		
High disruption	93	22.1	36	9.5		
Change in eating habits						
None	158	37.5	203	53.6	21.8	<0.001*
Medium disruption	185	43.9	132	34.8		
High disruption	78	18.5	44	11.6		
Change in body weight						
Decreased	59	14.0	83	21.9	19.1	<0.001*
Not changed	164	39.0	172	45.4		
Increased	198	47.0	124	32.7		
Change in physical activity						
Decreased	284	67.5	231	60.9	8.4	0.015
Not changed	113	26.8	106	28.0		
Increased	24	5.7	42	11.1		
*The significance was evaluated as p<0.003 with the Bonferroni correction. COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis						

*The significance was evaluated as $p < 0.003$ with the Bonferroni correction. COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis

use (OR: 2.2, 95% CI: 1.3-3.8, $p = 0.004$), and COVID-19-related knowledge (OR: 0.7, 95% CI: 0.7-0.9, $p = 0.020$) in PwMS (Table 5).

The predictors of depression include loss of relatives due to COVID-19 (OR: 1.8, 95% CI 0.9-1.0, $p = 0.010$) and COVID-19-related knowledge (OR: 1.7, 95% CI 0.7-0.9, $p = 0.001$) in the GP, whereas education (higher education) (OR: 0.3, 95% CI: 0.1-0.7, $p = 0.005$), marital status (being divorced or widowed) (OR: 2.7,

95% CI: 1.1-7.1, $p = 0.033$), antidepressant use (OR: 2.8, 95% CI: 1.7-4.8, $p < 0.001$), and knowledge on COVID-19 (OR: 0.8, 95% CI 0.7-1.0, $p = 0.023$) (Table 6) in the PwMS.

Discussion

The COVID-19 pandemic continues to be a burden on public health worldwide. The insufficient known effective treatment at the beginning of the pandemic, the anxiety of getting the

Table 3. COVID-19-related knowledge of patients with MS and the general population

Characteristics	Control		PwMS		χ^2	p
	n	%	n	%		
COVID-19 is transmitted by the coughing or sneezing of the infected person						
*Yes	335	79.6	319	84.2	2.8	0.093
No	86	20.4	60	15.8		
COVID-19 is transmitted through contact with the mouth, nose, and eyes						
*Yes	266	63.2	365	96.3	131.3	<0.001
No	155	36.8	14	3.7		
COVID-19 is transmitted by the contact of the hands with the mouth, nose, and eyes after touching the surface where the droplets adhere						
*Yes	373	88.6	352	92.9	4.3	0.038
No	48	11.4	27	7.1		
The mask is necessary to protect against COVID-19						
*Yes	419	99.5	376	99.2	0.3	0.906
No	2	0.5	3	0.8		
Hand washing is necessary to protect against COVID-19						
*Yes	396	94.1	361	95.3	0.5	0.457
No	25	5.9	18	4.7		
Social distancing is required to protect from COVID-19						
*Yes	421	100	379	10	-	1.00
No	-	-	-	-		
COVID-19 is transmitted from domestic animals						
Yes	142	33.7	90	23.7	9.6	0.002
*No	279	63.3	289	76.3		
Washing the nose with salty water prevents COVID-19						
Yes	84	20.0	82	21.7	0.4	0.545
*No	337	80.0	297	78.3		
Using alcohol prevents COVID-19						
Yes	67	15.9	21	5.5	21.9	<0.001
*No	354	84.1	358	94.5		
COVID-19 is transmitted by mosquitoes						
Yes	124	29.5	59	15.6	21.8	<0.001
*No	297	70.5	320	84.4		
The flu vaccine protects from COVID-19						
Yes	68	16.2	62	16.4	0.006	0.937
*No	353	83.8	317	83.6		
No. of correct answers mean ± SD (min-max)	9.10±1.50	(4-11)	9.85±1.21	(4-11)	7.813	<0.001**
*Correct answer option, **Independent samples t-test, COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis, SD: Standard deviation, min: Minimum, max: Maximum						

*Correct answer option, **Independent samples t-test, COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis, SD: Standard deviation, min: Minimum, max: Maximum

virus, and the uncertainties of the disease caused panic in society. Additionally, social and economic issues arose due to inevitable measures, such as social distancing and lockdown, that further affected the public mental health. This study was carried out during the second wave of the pandemic in Turkey to compare the anxiety and depression levels of PwMS with the GP and evaluate some related factors, including COVID-19-related knowledge.

Studies conducted during the COVID-19 period draw attention to the detrimental psychological effects of the pandemic (8,9,10). A study conducted in China at the beginning of the epidemic revealed that 54% of the society was psychologically affected, 17% experienced moderate-severe depression, and 29% had moderate-severe anxiety symptoms, whereas, in Turkey, 24% complained of depression, and 45% of anxiety (9,11). The pandemic caused

Figure 1. Comparison of the means of anxiety and depression scale scores between patients with MS and controls
MS: Multiple sclerosis

Figure 2. Scatterplot of anxiety and depression scores with COVID-19 knowledge scores
COVID-19: Coronavirus disease-2019, MS: Multiple sclerosis, GP: General population

additional unfavorable psychological effects in individuals with chronic diseases, including MS (4,8). Since the psychological status may alter the immune system, knowledge of the emotional changes of PwMS during the pandemic is critical (12).

Our study revealed that PwMS had increased levels of anxiety (32%) and depression (40%) compared to the pre-pandemic period (13). Additionally, we observed that anxiety and depression are related to the sociodemographic and clinical characteristics of patients. Similar to the literature, our study revealed that young age (14), low educational level (15), low income (16), being divorced/widowed (17), smoking (15), motor disability (16), frequent relapses (18), and antidepressant use (19) were all associated with increased levels of anxiety and depression. The regression models of patients revealed that young age, low educational level, and antidepressant use are predictors of anxiety, whereas low educational level, being divorced/widowed, and antidepressant use for depression. A study (20) revealed that 51% of patients discontinued their treatments because of anxiety, and this rate was 13% in our study. Another study reported that 26% of PwMS stopped their DMT and 13% decreased their dose without the permission of their health care providers (21).

Our study revealed that 42% of the general Turkish population had anxiety and 54% had depression. A previous Turkish study, conducted at the onset of the pandemic, reported that 45% of the GP had anxiety and 24% had depression (9). Another

study reported mild to moderate depression in 48% of patients (22). A meta-analysis reported that the prevalence of anxiety was 32%, and the prevalence of depression was 34% during the COVID-19 period (23). During the pandemic, various studies reported that some sociodemographic characteristics might affect anxiety and depression (11,22). Consistent with previous studies, our study revealed that female gender (9), unemployment (24), low educational level (11), and low income (25) were associated with anxiety and depression. As expected, many studies reported that having COVID-19 and the death of a close relative due to COVID-19 caused worsening in anxiety and depression levels (11,26). Likewise, our study also showed that the female gender was a predictor of anxiety, and the death of a close relative was a predictor of depression.

Contrary to the literature, the anxiety and depression levels of the GP were higher than PwMS when sociodemographic characteristics were adjusted using ANCOVA (21,27) due to acquired adaptability to unexpected conditions in chronic patients and active use of social media. The use of social media has been reported as an effective method in coping with depression in PwMS (28). Additionally, PwMS usually had easier access to their health care providers (i.e., physicians and nurses) during the pandemic, which might positively affect their anxiety and depression. Moreover, the GP's higher levels of anxiety and depression might have been caused by the high rate of unemployment, being single, antidepressant use, COVID-19 infection, and death of close relatives in the GP. It is worth mentioning that the majority of the GP was composed of young male individuals and a significant proportion was unemployed. Considering the gender roles of males in our patriarchal society, we can postulate that the endangered employment status of males might have been perceived as a higher stress factor that cause increased levels of anxiety and depression in the GP.

Another important feature that affects anxiety and depression is COVID-19-related knowledge. As the understanding increases, anxiety and depression are expected to decrease. The COVID-19-related knowledge was discovered as a predictor of anxiety and depression in all individuals. As in the literature, the knowledge level of the GP was lower than PwMS in our study (29). Hence, this low-level knowledge may explain the high levels of anxiety and depression in the GP.

The literature reported that the prevalence of COVID-19 was similar in MS and the GP (30). However, our study revealed more COVID-19 cases in the control than in the MS group. Interestingly, the fear of being infected by COVID-19 was lower in PwMS. PwMS have a lower perceived risk of COVID-19 due to their high level protection and isolation compared to controls.

Study Limitations

Our study has some limitations. First, the groups differ in terms of sociodemographic characteristics. Second, this study was conducted online, thus most participants were young individuals who actively used the technology. Individuals who did not have internet access were excluded from the study. Third, our study shows a momentary status in the constantly changing COVID-19 pandemic. Our study might have underestimated the COVID-19 prevalence because the participation rate of participants with active COVID-19 infection might have been low.

Table 4. Comparison of anxiety and depression scores of participants according to sociodemographic features and COVID-19 related changes in lifestyle

Characteristics	Control		PwMS	
	Anxiety	Depression	Anxiety	Depression
Age	-0.056	-0.058	-0.102	0.035
(r, p)	0.249	0.234	0.047	0.500
Gender				
Male	8.2±4.0	7.5±3.8	7.3±4.0	6.3±4.0
Female	9.0±4.2	7.8±4.2	7.8±4.6	6.4±3.8
p	0.049	0.388	0.290	0.825
Education				
Primary/secondary education	8.0±3.9	6.5±3.3	10.0±4.4	8.9±4.4
High school	8.2±4.0	7.9±3.9	8.2±4.7	7.5±4.2
University	8.7±4.2	7.6±4.1	7.1±4.3	5.8±3.5
p	0.435	0.209	0.001	<0.001
Income				
Less than expenses	9.6±4.59	8.7±4.2	10.0±4.4	8.5±4.2
Equal to expenses	7.8±3.82	6.8±3.8	6.9±4.3	5.9±3.5
More than expenses	8.3±3.6	8.0±3.7	6.9±4.1	4.6±3.2
p	0.001	<0.001	<0.001	<0.001
Marital status				
Married	8.3±3.7	7.2±3.9	7.5±4.3	6.3±3.9
Single	8.6±4.4	7.9±4.1	7.5±4.5	6.1±3.5
Divorced/widow	8.6±2.9	8.8±3.3	8.5±4.9	8.1±4.5
p	0.738	0.196	0.500	0.027
Smoking				
Yes	8.7±4.0	8.1±3.8	8.5±4.7	7.3±4.2
No	8.3±4.2	7.3±4.1	7.3±4.3	6.0±3.7
p	0.333	0.045	0.024	0.004
Antidepressant use				
Yes	9.5±2.9	8.7±3.9	9.1±5.1	7.9±4.5
No	8.4±4.1	7.6±4.0	7.2±4.2	6.0±3.5
p	0.340	0.295	0.003	<0.001
COVID-19 diagnosis				
Yes	8.2±4.68	7.0±4.0	9.5±5.4	6.5±4.1
No	7.2±4.33	6.0±3.7	7.5±4.4	6.4±3.8
p	0.037	0.014	0.028	0.85
Concerns about COVID-19				
None	6.8±4.4	6.8±4.4	7.1±4.47	5.7 ± 3.4
Mild level	7.6±3.5	6.9±3.5	7.4±4.64	6.4 ± 4.0
Moderate level	8.9±4.0	7.7±4.0	7.6±4.35	6.2 ± 3.6
High level	10.7±4.0	9.6±3.8	7.9±4.57	6.7 ± 4.1
p	<0.001	<0.001	0.749	0.533
Change in physical activity				
Worse	9.2 ± 4.1	8.0 ± 4.1	8.1±4.6	7.1±3.9
None	7.1 ± 3.9	6.5 ± 3.6	6.9±4.5	5.7±3.8
Better	7.4 ± 4.0	7.9 ± 3.8	6.9±3.9	4.6±2.6
p	<0.001	0.002	0.052	<0.001
Change in sleep				
None	6.3 ± 3.6	5.8 ± 3.7	5.4 ± 3.7	5.1±3.4
Medium disruption	8.8 ± 3.7	7.8 ± 3.7	9.5 ± 4.0	7.4±3.6
High disruption	10.7 ± 4.3	9.5 ± 4.0	10.3 ± 4.9	8.2±4.9
p	<0.001	<0.001	<0.001	<0.001

Table 4. continued				
	Control		PwMS	
Change in eating habits				
None	6.7±3.7	6.2±3.8	5.9±3.8	5.2±3.6
Medium disruption	9.0±3.9	7.9±3.6	9.3±4.2	7.4±3.3
High disruption	10.9±3.8	9.7±4.0	10.4±4.6	8.7±4.6
p	<0.001	<0.001	<0.001	<0.001
Change in body weight				
Decreased	9.1±3.9	8.4±4.3	7.8±4.4	6.9±4.1
Not changed	7.3±3.8	6.7±3.6	6.8±4.3	5.6±3.6
Increased	9.3±4.2	8.1±4.1	8.5±4.5	7.2±3.9
p	<0.001	0.001	0.005	0.001
COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis				

COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis

Table 5. Multivariate logistic regression analysis to determine the predictors of anxiety						
Variables	Control			PwMS		
	OR	95% CI (lower-upper)	p	OR	95% CI (lower-upper)	p
Age	0.9	1.0-1.0	0.823	0.9	0.9-0.9	0.010
Gender (female)	1.7	1.1-2.6	0.015	1.0	0.6-1.7	0.833
Education						
Primary school	-	-	0.200	-	-	0.009
High school	0.8	0.3-1.9	0.627	0.3	0.1-0.8	0.025
University	1.2	0.5-2.7	0.581	0.2	0.1-0.6	0.002
Marital status						
Married	-	-	0.800	-	-	0.220
Single	0.8	0.1-6.4	0.907	1.8	0.7-4.8	0.229
Divorced/widow	0.7	0.4-1.6	0.504	0.8	0.4-1.7	0.600
Having children	1.3	0.6-3.0	0.400	1.4	0.7-2.9	0.301
Antidepressant use	1.0	0.3-3.1	0.949	2.2	1.3-3.8	0.004
COVID-19 diagnosis	1.6	0.9-2.9	0.059	2.9	0.9-5.4	0.085
Loss of relatives due to COVID-19	1.1	0.7-1.8	0.524	1.6	0.8-3.2	0.160
Knowledge of COVID-19	0.8	0.7-0.9	0.007	0.7	0.6-1.0	0.020

COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis, OR: Odds ratio, CI: Confidence interval

Table 6. Multivariate logistic regression analysis to determine the predictors of depression						
Variables	Control			PwMS		
	OR	95% CI (lower-upper)	p	OR	95% CI (lower-upper)	p
Age	1.0	0.9-1.0	0.780	0.9	0.9-1.0	0.269
Gender (female)	1.4	0.9-2.1	0.109	0.7	0.4-1.1	0.200
Education						
Primary school	-	-	0.549	-	-	0.015
High school	1.1	0.4-2.6	0.742	0.4	0.1-1.0	0.071
University	1.3	0.6-3.0	0.395	0.3	0.1-0.7	0.005
Marital status						
Single	-	-	0.601	-	-	0.103
Married	1.0	0.5-2.0	0.919	1.3	0.6-2.5	0.387
Divorced/widow	0.3	0.1-2.1	0.332	2.7	1.1-7.1	0.033
Having children	0.7	0.3-1.5	0.423	0.8	0.4-1.5	0.504
Antidepressant use	1.6	0.5-5.0	0.406	2.8	1.6-4.8	<0.001
COVID-19 diagnosis	1.1	0.6-1.9	0.724	0.7	0.3-1.9	0.565
Loss of relatives due to COVID-19	1.8	1.1-2.9	0.010	1.3	0.7-2.6	0.346
Knowledge of COVID-19	0.7	0.7-0.9	0.001	0.8	0.7-1.0	0.023

COVID-19: Coronavirus disease-2019, PwMS: Patients with multiple sclerosis, OR: Odds ratio, CI: Confidence interval

Conclusion

In conclusion, our study revealed that the COVID-19 pandemic had a significant effect on the anxiety and depression levels of the GP and PwMS. Additionally, the GP had higher levels of anxiety and depression compared to PwMS. Anxiety and depression levels were also determined to be associated with various sociodemographic and clinical characteristics of patients. The female gender was related to anxiety, the death of a close relative due to COVID-19 to depression, and COVID-19-related knowledge to both anxiety and depression in the GP. In PwMS, age was related to anxiety, being single to depression, and low education level, antidepressant use, and COVID-19-related knowledge to both anxiety and depression. Our study emphasizes the importance of psychological support to the GP and PwMS during the pandemic.

Ethics

Ethics Committee Approval: The ethical principle compliance was evaluated by the ethics committee of the institution where the study was conducted, and the Ethics Committee approval was obtained (Istanbul University, Istanbul Faculty of Medicine, Department of Neurology, Istanbul, Turkey, file number: 2020/1095, no: 29624016-050.99- 1215).

Informed Consent: Consent to participate was obtained using online forms.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Concept: G.Y.Y., C.P.D., Z.T., M.K., M.E., Design: G.Y.Y., C.P.D., Z.T., M.K. Data Collection or Processing: G.Y.Y., C.P.D., Analysis or Interpretation: G.Y.Y., C.P.D., Z.T., Literature Search: G.Y.Y., C.P.D., Writing: G.Y.Y., C.P.D.

Conflict of Interest: The authors have not declared any conflict of interest related to this article.

Financial Disclosure: No financial support was received from any institution or person for our study.

References

- World health organization. Coronavirus disease pandemic. Access date 10.06.2021 Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
- Health TRM of platform of covid-19 information. Access date 10.06.2021 Available from: <https://covid19.saglik.gov.tr/>
- Kendzerska T, Zhu DT, Gershon AS, et al. The Effects of the Health System Response to the COVID-19 Pandemic on Chronic Disease Management: A Narrative Review. *Risk Manag Healthc Policy* 2021;14:575-584.
- Capuano R, Altieri M, Bisecco A, et al. Psychological consequences of COVID-19 pandemic in Italian MS patients: signs of resilience? *J Neurol* 2021;268:743-750.
- Mantero V, Abate L, Balgera R, et al. Assessing the susceptibility to acute respiratory illness COVID-19-related in a cohort of multiple sclerosis patients. *Mult Scler Relat Disord* 2020;46:102453.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-370.
- Aydemir Ö, Güvenir T, Kuey L, Kültür S. Validity and reliability of Turkish version of hospital anxiety and depression scale. *Turk Psikiyatri Derg* 1997;8:280-287.
- Chiaravallotti ND, Amato MP, Brichetto G, et al. The emotional impact of the COVID-19 pandemic on individuals with progressive multiple sclerosis. *J Neurol* 2021;268:1598-1607.
- Özdin S, Bayrak Özdin Ş. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *Int J Soc Psychiatry* 2020;66:504-511.
- Wang Y, Di Y, Ye J, Wei W. Study on the public psychological states and its related factors during the outbreak of coronavirus disease 2019 (COVID-19) in some regions of China. *Psychol Health Med* 2021;26:13-22.
- Nie XD, Wang Q, Wang MN, et al. Anxiety and depression and its correlates in patients with coronavirus disease 2019 in Wuhan. *Int J Psychiatry Clin Pract* 2021;25:109-114.
- Mohr DC, Hart SL, Julian L, Tasch ES. Screening for depression among patients with multiple sclerosis: two questions may be enough. *Mult Scler* 2007;13:215-219.
- Boeschoten RE, Braamse AMJ, Beekman ATF, et al. Prevalence of depression and anxiety in Multiple Sclerosis: A systematic review and meta-analysis. *J Neurol Sci* 2017;372:331-341.
- Alschuler KN, Roberts MK, Herring TE, Ehde DM. Distress and risk perception in people living with multiple sclerosis during the early phase of the COVID-19 pandemic. *Mult Scler Relat Disord* 2021;47:102618.
- Pham T, Jetté N, Bulloch AGM, et al. The prevalence of anxiety and associated factors in persons with multiple sclerosis. *Mult Scler Relat Disord* 2018;19:35-39.
- McKay KA, Tremlett H, Fisk JD, et al. Psychiatric comorbidity is associated with disability progression in multiple sclerosis. *Neurology* 2018;9:e1316-e1323.
- Landfeldt E, Castelo-Branco A, Svedbom A, et al. The long-term impact of multiple sclerosis on the risk of divorce. *Mult Scler Relat Disord* 2018;24:145-150.
- Fisher PL, Salmon P, Heffer-Rahn P, et al. Predictors of emotional distress in people with multiple sclerosis: A systematic review of prospective studies. *J Affect Disord* 2020;276:752-764.
- Motolese F, Rossi M, Albergo G, et al. The Psychological Impact of COVID-19 Pandemic on People With Multiple Sclerosis. *Front Neurol* 2020;11:580507.
- Alnajashi H, Jabbad R. Behavioral practices of patients with multiple sclerosis during Covid-19 pandemic. *PLoS One* 2020 22;15:e0241103.
- Shaygannejad V, Afshari-Safavi A, Hatef B. Assessment of mental health, knowledge, and attitude of patients with multiple sclerosis and neuromyelitis optica spectrum disorder in response to 2019 novel coronavirus. *Neurol Sci* 2021;42:2891-2901.
- Ustun G. Determining depression and related factors in a society affected by COVID-19 pandemic. *Int J Soc Psychiatry* 2021;67:54-63.
- Salari N, Hosseini-Far A, Jalali R, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Global Health* 2020;16:57.
- Shah SMA, Mohammad D, Qureshi MFH, Abbas MZ, Aleem S. Prevalence, Psychological Responses and Associated Correlates of Depression, Anxiety and Stress in a Global Population, During the Coronavirus Disease (COVID-19) Pandemic. *Community Ment Health J* 2021;57:101-110.
- Smith L, Jacob L, Yakkundi A, et al. Correlates of symptoms of anxiety and depression and mental wellbeing associated with COVID-19: a cross-sectional study of UK-based respondents. *Psychiatry Res* 2020;291:113138.
- Mazza MG, De Lorenzo R, Conte C, et al. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain Behav Immun* 2020;89:594-600.
- Costabile T, Carotenuto A, Lavorgna L, et al. COVID-19 pandemic and mental distress in multiple sclerosis: Implications for clinical management. *Eur J Neurol* 2021;28:3375-3383.
- Farpour HR, Hoveidaei AH, Habibi L, Moosavi M, Farpour S. The impact of social media use on depression in multiple sclerosis patients. *Acta Neurol Belg* 2020;120:1405-1409.
- Sahraian MA, Azimi A, Navardi S, Ala S, Naser Moghadasi A. Evaluation of the rate of COVID-19 infection, hospitalization and death among Iranian patients with multiple sclerosis. *Mult Scler Relat Disord* 2020;46:102472.
- Berger JR, Brandstadter R, Bar-Or A. COVID-19 and MS disease-modifying therapies. *Neurol Neuroimmunol Neuroinflamm* 2020;7:e761.