Objectives

To compare the 6-month clinical performance of a “universal” adhesive in non-carious Class V lesions using four different adhesive strategies.

Materials and Methods

21 patients participated in this study, in which 70 Class V restorations were placed. The restorations were randomly assigned into four experimental groups according to different adhesive strategies of Scotchbond Universal Adhesive (SBU, 3M ESPE): A. 3-step etch-and-rinse: 34% phosphoric acid (PA, Scotchbond Universal Etchant, 3M ESPE) and application of SBU followed by one coat of the non-solvated bonding resin Scotchbond Multi-Purpose Adhesive (SBMPA, 3M ESPE); B. 2-step etch-and-rinse: 34% PA followed by SBU; C. 2-step self-etch: SBU followed by one coat of SBMPA; D. 1-step self-etch: SBU alone. All restorations were evaluated at baseline and after 6 months by two blind observers using the USPHS criteria. Statistical analysis was performed with the non-parametrical tests Kruskal-Wallis, Mann Whitney U and Wilcoxon ($p < 0.05$).

Results

Only one restoration from the group 1-step/SBU was lost at six months. Marginal adaptation was the only criterion for which statistically worse scores were measured after 6 months ($p < 0.01$). Significantly more bravo scores were detected when SBU was used following a self-etch strategy. The restorations performed with SBU as 1-step self-etch adhesive exhibited a significantly deterioration of the marginal adaptation after 6 months.

Conclusions

Restorations performed with SBU under a self-etch strategy showed worse marginal adaptation after 6 months of clinical use compared to those with SBU under an etch-and-rinse strategy. The addition of a non-solvated hydrophobic coating (SBMPA) did not influence the clinical performance.

- Oral Presentation 31

TITLE: Endodontic treatment for avoiding an inferior alveolar nerve paresthesia

AUTHORS: Gómez Álvarez G, Gómez Martín C, Del Valle Aleixandre B, Zorita García M, Mena Álvarez J.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S15.

* doi:10.4317/jced.17643815

<http://dx.doi.org/10.4317/jced.17643815>

Introduction

In some cases, the proximity of the inferior alveolar nerve to the lower molars roots causes that when we face chronic apical periodontitis with pulpal origin, it might exist an inferior alveolar nerve affectation because of the invasion of the mandibular canal by the injury. An accurate diagnostic of the situation of that injury will be the key in the plan of treatment we should make.

Case report

A 69 year old male patient comes to consulting room asking for a routine consultation. With a first visualization of his ortopantomography, we can see the tooth number 4.7 affected of a chronic apical periodontitis which overlaps with the canal of the alveolar inferior nerve. This overlapping is then better appreciated using a periapical radiography. The patient is completely asymptomatic.

It is decided to do a Cone Beam Computed Tomography (CBCT) to see the injury's location; it is localized in a lingual position of the mandibular canal, starting to invade it and breaking the lingual plate.

Then we proceed to remove the metal crown the patient is wearing in that tooth, and after an evaluation of the remaining tooth under the metal crown, we proceed with the root canal treatment.

When we face these kind of injuries that are overlapping the alveolar inferior nerve canal, we should be careful and make an appropriate diagnostic, with the aim of getting success of the treatment; a quick action in these casual findings will avoid major injuries, being very important the following in the future of these kind of injuries.

Conclusions

The use of Cone Beam Computed Tomography (CBCT) is essential when we find this kind of overlappings, allowing three dimensional visualization. Root canal treatment is able to avoid higher pathologies, such as paresthesias, obtaining bone regeneration in the area and thus improving patient's health.

- Oral Presentation 32

TITLE: Application of Bioinformatics in the Mount/Hume classificacition of caries and his relationship with Orthopantomography

AUTHORS: Hernando Dumaraog B, De Paz JF, Corchado JM, García E, Aliaga I, Campo L, Vera V.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S15.

* doi:10.4317/jced.17643816

<http://dx.doi.org/10.4317/jced.17643816>

Objectives

The incorporation of computational techniques and artificial intelligence in the area of biomedicine has made remarkable progress in the prevention and detection of diseases. Decision trees are a prediction model used in the field of artificial intelligence that provides a human expert information, a caries classification in this study, made by the system, generating rules that will support decision making.

Dental caries is one of the most prevalent infectious diseases in patients and the diagnosis is made by dentists using clinical and radiological examination.

The aim of this study is to analyze the relationship between the Mount/Hume classification of caries and Orthopantomography x-ray.

Materials and Methods

The study was realized on fifty patients who attended to the triage of the School of Dentistry in the Complutense University of Madrid and performed by a faculty of the department of Conservative Dentistry. After clinical examination and radiological study, all the findings were uploaded in patient's chart, the data collected is sent to the University of Salamanca to be analyzed by Bioinformatics.

Results

Three decision trees were generated by the algorithm J48. Decision tree number two, associated the variables size and location of the lesion using the Mount/Hume classification and radiographic variable.

Conclusions

Decision trees are a simple tool that allows us to visualize and analyze the relationship between the Mount/Hume classification of caries and extraoral radiographic study.

Decision trees used in this study highlight the need for the use of the panoramic radiograph for classifying caries sizes 2, 3 and 4 and location 2 of Mount/Hume and not in sizes 0 and 1 that require more specific diagnostic tests.

- Oral Presentation 33

TITLE: Evaluation of the shades in the space ciel-ch of amaris composites

AUTHORS: Guzmán Pina S, Funes Gil I, Fernández Sánchez G, Chiva García F.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S16.

* doi:10.4317/jced.17643817

<http://dx.doi.org/10.4317/jced.17643817>

Objectives

To evaluate color change parameters in space Ciel-ch of Amaris(VOCO®) dentin composite resin when adding its enamel colors

Materials and Methods

45 discs, 2mm thick and 6 mm in diameter were created of Amaris dentin shade, and divided in 5 groups: group1-shadeO1, group2-shadeO2, group3-shadeO3, group4-shadeO4 and group5-shadeO5. In each group (n=9) we added to the dentin shade disks, 1mm of Amaris enamel shade (TL, TN, TD), obtaining 3 discs per group of each enamel shade. The color parameters (L, c, h) were determined with the composite-resin light-cured, by a spectrophotometer EasyShade(Vita®), before and after adding the enamel shade. The results were analyzed by comparing the variances (ANOVA) with statistic package SPSS-v.15

Results

In groups 4 and 5, all enamel shades increased significantly its lightness (L) ($p < 0.001$). In group 3 there were no significant differences ($p = 0.08$), in group 2 there was only a significant increase in the shade TL ($p = 0.03$), and in group 1 there was an increase with the shade TL but a decrease with the shade TN and TD ($p = 0.02$). Regarding chroma (c), all 5 groups decreased significantly when placing the 3 enamel shades. As for hue (h) there was not significant decrease in groups 2 ($p = 0.05$) and 5 ($p = 0.17$). In groups 3 and 5, only the enamel shade TL presented significant differences ($p = 0.001$). In group 1, all 3 enamel shades presented a significant difference ($p = 0.002$). The correlation with Vita® shades was A3.5 for 2mm of dentin shades O1, O2 and O3, and A4 for shades O4 and O5. For 1mm of enamel shades, its correlation was A2 for TN, B1 for TL and A1 for TD.

Conclusions

The placing of 1mm thickness of enamel shade, decreased the chroma of the dentin shades. The shade TL increased the lightness but the shade TD did not decrease it.

- Oral Presentation 34

TITLE: Endodontic failure due to forgotten duct : a series of cases

AUTHORS: García Marcos JI, Santos Cubero J, Alonso Ezpeleta LO, Mena Alvarez J.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S16.

* doi:10.4317/jced.17643818

<http://dx.doi.org/10.4317/jced.17643818>