- Oral Presentation 56

TITLE: Micro-invasive treatment of orthodontic white-spot lesions

AUTHORS: Peixoto A, Carmo J, Sanches C, Manso AG.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S27.

* doi:10.4317/jced.17643840

<http://dx.doi.org/10.4317/jced.17643840>

Introduction

Icon® is a resin infiltrant used for micro-invasive treatment of smooth surface and proximal caries lesions. It's used to treat lesions up to first third of dentin (D1). Resin infiltration aims to penetrate lesion using low viscosity resin with high penetration properties, in a single visit, with no drilling. The system consists in Icon-Etch (Hydrochloric acid, pyrogenic silicic acid, surface-active substances), Icon-Dry (99% ethanol), Icon-Infiltrant (Methacrylate-based resin matrix, initiators, additives), Approximal-Tips, Luer-Lock-Tip and dental wedges. The purpose of these cases is to demonstrate the use of this technique in orthodontic white-spots.

Case report

Two female patients were selected from CESEM's University Clinic, Caparica-Portugal. Both presented white-spot lesions due to orthodontic appliance. Case 1- 13 years old, orthodontic removal appointment 14 days before Icon treatment, tooth chosen 11. Case 2- 21 years old, orthodontic removal appointment 7 years ago, tooth chosen 44. The teeth were cleaned, rubber dam placed and Resin infiltrant applied according to manufacturer's instructions. The patient was instructed to floss and brush with fluoridated toothpaste. The lesions were photographed before, immediately after the resin infiltrant application and 2 weeks later. White-spot lesions were filled with resin infiltrant and looked similar to sound enamel, immediately. Porous enamel has lower refraction than sound enamel, but when infiltrated, refraction increases, improving, dramatically, the aesthetic. Other studies refer that infiltration combined with remineralization shows better results.

Conclusions

Clinical management of white spots, with restorations, fluor therapy, microabrasion, is challenging and aesthetic outcome is not always predictable, however in these cases we demonstrated successful use of Icon® for masking orthodontic white-spots. The infiltrant is simple and conservative. Further research is needed to evaluate long-term stability, efficacy and microscopic changes.

- Oral Presentation 57

TITLE: Multidisciplinary treatment of endo-periodontal lesions

AUTHORS: Peña Alcázar M, Zubizarreta Macho A, Rico Romano C, Sierra Armas L.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S27.

* doi:10.4317/jced.17643841

<http://dx.doi.org/10.4317/jced.17643841>

Introduction

The tooth and supporting tissues should not be considered as separate entities but as a biological unit, the result of the close relationship between the two. These structures have many roads that allow two-way exchange of nutrients and harmful substances. This anatomical link explains the simultaneous presence of periodontal and endodontic conditions of nature. Endo-periodontal lesions are described as inflammatory nature of those injuries that simultaneously affect the dental pulp tissue and tooth support. The therapeutic objective of the combined lesions lies in the etiological treatment of these pathological processes.

Case report

The case of a 53-year-old referred to the Master in Clinical Endodontics and Microsurgery Periapical University of Alfonso X El Sabio, presenting grade II mobility on tooth 3.4 is described. The results of the clinical and radiological evidence showed concurrent endo-periodontal lesion in tooth 3.4. Canal treatment of the affected tooth followed by periodontal debridement surgery that allowed the crowded bacterial deposits on the root surface was performed. The multidisciplinary treatment of this tooth was decisive for the survival of the affected tooth and allow tissue repair of the affected tissues.

Conclusions

Endo-periodontal lesions are a challenge for the clinician. It should be emphasized the importance of a correct diagnosis, and the prognosis of these lesions depends on immediate treatment plan.

- Oral Presentation 58

TITLE: Influence of clinical usage of ProTaper S1 instrument in cyclic fatigue resistance

AUTHORS: Pérez-Higueras JJ, Arias A, de la Macorra JC.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S27.

* doi:10.4317/jced.17643842

<http://dx.doi.org/10.4317/jced.17643842>

Objectives

To compare the influence of the amount of in vivo shaped root canals (RC) in static cyclic fatigue (CF) resistance in ProTaper Universal (PTU) S1 instruments at different points of curvature.

Materials and Methods

A total of 80 files PTU S1 were divided in four groups of 20 instruments each: control group 0 (unused files), group 1 (files used to shape up to 10 RC), group 2 (files used in 11-15 RC) and group 3 (files used in more than 15 RC). The same endodontist with more than 15 years of experience in rotary canal instrumentation shaped all RC using always the same protocol (PTU S1 was used to working length after the achievement of a reproducible glide path with a #15 hand file. 5,25% NaOCl solution was used for irrigation). CF resistance was tested in stainless-steel curved canals (60°, r=3mm). All instruments were rotated at 300rpm until fracture at 5mm and at 10 mm from the tip, making a total of 8 subgroups (0-5, 0-10, 1-5, 1-10, 2-5, 2-10, 3-5, 3-10). Time to fracture was recorded. Mean half-life, Beta and Eta were calculated for each group and were compared with Weibull analysis.

Results

PTU S1 instruments will last significantly longer at 5 than at 10 mm with a 100% probability independently of the number of shaped RC, however there were not significant differences in CF resistance among the four groups (unused instruments or files used to shape different number of RC), neither at 5 nor at 10 mm.

Conclusions

PTU S1 was significantly more resistant at 5 than at 10 mm from the tip. The increase in the number of RC shaped with the same instrument did not decrease CF resistance of the instrument.

- Oral Presentation 59

TITLE: In vitro comparison of the antimicrobial effect of Tri-antibiotic paste components, amoxicillin and clavulanic acid against *Fusobacterium nucleatum* and *Actinomyces naeslundii*

AUTHORS: Prieto González L, González Nieto A, Martínez Pinar A, Basel Sawas Jiménez M, Rubio Flores D, García Barbero E.

SOURCE: J Clin Exp Dent. 2014 1;6 (Supplement1):S28.

* doi:10.4317/jced.17643843

<http://dx.doi.org/10.4317/jced.17643843>

Objectives

To compare the effectiveness of Tri-antibiotic Paste components (Minocycline / metronidazole / ciprofloxacin), amoxicillin and clavulanic acid against two endodontic pathogens, *Fusobacterium nucleatum* and *Actinomyces naeslundii*.

Materials and Methods

It was determined the minimum inhibitory concentration (MIC) of minocycline, metronidazole, ciprofloxacin, amoxicillin and clavulanic acid by microdilution assay in 96-well Mueller- Hinton agar plates according to EUCAST. We seeded 50 µl of standard bacterial suspension (corresponding to 0.5 of the McFarland scale), with 50 µl of antibiotic solution in a concentration gradient. Under spectrophotometry, we calculated the optical density (OD 600) to determine the lowest concentration (µg / ml) that inhibit the visible growth of the microorganisms after overnight incubation (MIC). Statistical analysis was carried out by one way ANOVA and post- hoc Duncan test at a level of significance of p<0,05.

Results

For *Fusobacterium Nucleatum*, no significant differences were found for amoxicillin, metronidazole and minocycline. Nevertheless, ciprofloxacin resulted in higher MIC value than the others, indicating its lower antibacterial activity.

For *Actinomyces naeslundii* colonies, no differences were found between amoxicillin and minocycline. Ciprofloxacin and metronidazole showed higher MIC values in a significant manner.

for both bacteria, the values obtained from the clavulanic acid can be translated in terms of bacterial resistance.

Conclusions

No significant differences were found for inhibiting bacterial colonies with amoxicillin or Tri-antibiotic paste components, needing higher concentrations to inhibit the growth of ciprofloxacin with *Fusobacterium nucleatum*, and metronidazole and ciprofloxacin with *Actinomyces naeslundii*.

By itself, clavulanic acid is not able to inhibit the growth of both bacteria studied.

In view of the results of this study, it seems appropriate to analyse the effectiveness of Tri-antibiotic paste components combined with amoxicillin and the combination of amoxicillin with clavulanic acid.