- Oral Presentation 41

TITLE: Comparison between graft block autologous and distraction osteogenesis as treatment for remodeling of atrophic jaw

AUTHORS: Malvis Bugallo M, Oliván Molina S, Ingelmo Navarro P, Loughney González A, Fernández Domínguez M.

Master en Cirugía Bucal e Implantología. Universidad San Pablo CEU. Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S17.

* doi:10.4317/medoral.17643931
http://dx.doi.org/10.4317/medoral.17643931

Objectives: Bone distraction and grafting in block are the most suitable techniques in cases of large bone destruction. Then both techniques will be compared and an analysis based on the literature will be.

Material and Methods: Search for articles, in the last 10 years, published in Medline (PUBMED) and EMBASE; key words: distraction osteogenesis, graft-block.

Results: Distraction osteogenesis, offers lower infection rate, always soft tissues are won and there is a donor area morbidity. Its difficulty lies in directing the vector of distraction, although there is also a risk of necrosis of the absent-minded fragment, and its success depends on the behavior and care of the patient.

Chin is a widely used donor area because it presents an acceptable bone, mainly cortical quantity, good surgical access. The main drawback is the morbidity from the donor area, which depends on the dimension of retrieved bone and surgical management. The most important aspect is to obtain a primary closure of the receiving area.

Conclusions: To reset the bone volume in an atrophic maxilla, should be individualized each case, studying the form and severity of the defect. Horizontal defects in the technique of choice are bone grafts. While in vertical defects, alveolar distraction is a predictable method, particularly in the former sector. There are cases in which there is loss of vertical and horizontal, where you can combine both techniques.

- Oral Presentation 42

TITLE: Resolution of clinical cases in Dentistry on Patients with Special Needs: technique of Aronson's Puzzle

AUTHORS: Mariño Pérez R; Gándara Lorenzo D; Piñeiro Abalo S; Suárez Cunqueiro MM; Tomás Carmona I.

Department of Stomatology, School of Medicine and Dentistry. University of Santiago de Compostela, Spain.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S17.

* doi:10.4317/medoral.17643932
http://dx.doi.org/10.4317/medoral.17643932

Background: To assess the student's perception and the effectiveness of a Cooperative Learning technique known as Aronson's Puzzle for solving clinical cases.

Methods: During the academic years 2012/13 and 2013/14, the technique of Aronson's Puzzle was applied in the subject "Dentistry on Patients with Special Needs" belonging to the fifth degree course in Dentistry. A total of 109 students were enrolled in the study. We compared the final qualifications in Puzzle Group (55 students) and in Non-Puzzle Group (54 students). The perception of the students was assessed using a questionnaire with 13 questions.

Results: The academic performance in the subject was expressed with global mean (Standard Deviation, SD) exam score for the Puzzle Group 6.45 (SD= 1.49) and for the Non-Puzzle Group 6.13 (SD= 1.50). There was a significant decrease in the Puzzle Group in the number of students who did not show up for exams (0% vs 13% in the Non-Puzzle Group, $p < 0.006$). The mean value for all items in the questionnaire was 3.80 with the highest response score 4.30 for the item "This type of activity has allowed me to see the complexity of the search for the complete solution of a case". Internal consistency (Cronbach's alpha) for questionnaire was 0.90.

Conclusion: Our results suggest that Puzzle Technique of Aronson contributes to a better understanding of the complexity in solving clinical cases. Further investigations should be conducted to analyze the influence of this Cooperative Learning technique in the improvement of student's academic performance in the Clinical Dentistry.

- Oral Presentation 43

TITLE: Study of carotid artery calcifications on panoramic radiographs

AUTHORS: Martí Ainsa A., Viñals Iglesias H.

Facultat d'Odontologia. Universitat de Barcelona.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S17.

* doi:10.4317/medoral.17643933
http://dx.doi.org/10.4317/medoral.17643933

Objective: The aim of this study is to determine the prevalence of the carotid artery calcifications (CAC) detected by panoramic radiographs, and to associate the CAC with other diseases like type II diabetes, dyslipidemia, hypertension, obstructive sleep apnea and menopause.

Material and Methods: A total of 48 males and 48 females were chosen at random. Each one had available a panoramic radiography and at least one of the diseases mentioned above. They were classified in four age groups (50-59, 60-69, 70-79, ≥ 80) with 12 individuals in each group. Panoramic radiographs were visually analyzed and the presence of images compatibles with a calcified atheroma plaque (CAP) was detected. We consider a CAP as an irregular radiopaque mass, circular or heterogeneous, bilateral or unilateral which is located at posterior and inferior to the mandibular angle, approximately between C3 and C4 vertebrae, which can appear in the upper or lower border of the hyoid bone.

Results: Our results show:

- The prevalence of CAC of the total panoramic radiographies analyzed.
- The prevalence of the patients with the considered diseases and the presence of CAC on panoramic radiography.
- The description of CAC presence/absence according to age and sex.

Conclusions: The early detection of CAC on panoramic radiographs – diagnostic tool routinely used in dentistry – would mean to refer the patient to a specialist for further examination. It contributes to improve the prevention of cardiovascular diseases and cerebrovascular events.

- Oral Presentation 44

TITLE: Which factors condition treatment choice and outcomes of implant-based rehabilitation of edentulous geriatric patients?

AUTHORS: Martín Ares M, Buesa Báez JM, Serrano Torrecilla M, Merchán Morales S, Barona Dorado C.

Máster de Cirugía Bucal e Implantología. Hospital Virgen de la Paloma.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S18.

* doi:10.4317/medoral.17643934
http://dx.doi.org/10.4317/medoral.17643934

Objectives: To identify factors influencing the outcomes of implant-based rehabilitation of geriatric patients evaluated by a satisfaction survey of subjective patient opinion and by professional evaluation of clinical and radiological parameters.

Materials and Methods: This transversal retrospective study included 150 completely edentulous geriatric patients with implant-based rehabilitations. A 19-item questionnaire was designed to assess patient satisfaction; clinical outcomes were evaluated by registering clinical and periodontal parameters. Age, sex, patient habits and restoration type were registered to determine whether these factors influenced the clinical results of implant-based rehabilitation.

Results: Significant differences were found in patient satisfaction conditioned by age, sex, and prosthetic type. Patient habits influenced long-term clinical results significantly.

Conclusions: For the geriatric patient, choice of prosthetic type must be based on clinical and anatomical parameters, as well as age, sex and the patient's habits. Patient satisfaction with the outcome will be conditioned by adequate treatment selection and planning.

- Oral Presentation 45

TITLE: Granular cell tumor in the geriatric patient. Clinical, histological and immunohistochemical study

AUTHORS: Martínez Balayo P, Pérez López D, Lucas González R, Peñamaría Mallón MA, Otero Rey E, Blanco Carrión A.

Máster de Odontología Práctica Diaria. Universidad de Santiago de Compostela.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S18.

* doi:10.4317/medoral.17643935
http://dx.doi.org/10.4317/medoral.17643935

Objective: Analysing the clinical and histological characteristics of the granular cell tumor and remarking the part played by the dentist in the diagnosis and treatment of this neoplasia.

Material and Method: Four cases of tumors of granular cells of oral location are presented and a bibliographical revision is made through a search in Pubmed-Medline.

Results: Four cases of Abrikossoff's tumor located on labial mucosa (1) and tongue (3) are presented. Clinically observed in all of them an exophytic and white-yellowing lesion. To get the diagnostic, a biopsy and a histological and immunohistochemical study was performed. This study tested positive for specific tumor markers: Vimentin and S100 Protein

Conclusions: Histologic features and positivity of specific markers in the immunohistochemical study defines the definitive diagnosis of this tumor.

- Oral Presentation 46

TITLE: Platelet rich plasma and its indication as an adjuvant treatment in hard and soft tissue regeneration

AUTHORS: Martínez Boreas C, Pérez Corral I, De Antonio Torres C, Loughney González A, Fernández Domínguez M.

Master en Cirugía Bucal e Implantología. Universidad San Pablo CEU. Madrid.