7,733mm ($\pm 1,8619$), opening 45,422mm ($\pm 6,1572$). The condylar path values have been 46,6925° ($\pm 7,53386$) for the right side and 49,2299° ($\pm 7,97528$) for the left one. Significant statistically differences have been found between the left and the right lateralities, being bigger the left one ($p < 0.000$) and between the right and the left condylar path, being higher the left one ($p = 0.009$).

Conclusion: In the current study we find out that in the population studied, the lateral movement is bigger and the condylar path is higher on the left side.

- Oral Presentation 57

TITLE: Épulis fissuratum. A case report

AUTHORS: Puente Fernández S, De la Calle Cañas C, Martínez Rodríguez N, Fernández Cáliz F, Belarra Arenas C.

Máster en Cirugía Bucal e Implantología. Hospital Virgen de la Paloma.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S23.

* doi:10.4317/medoral.17643947
http://dx.doi.org/10.4317/medoral.17643947

Introduction: Inflammatory fibrous hiperplasia or epulis fissuratum) is the most common lesion of soft tissue in the oral cavity. It is more frequent in females. It is an adaptive overgrowth located over the soft tissues of the vestibular sulcus or alveolar ridges caused by low intensity chronic irritation from poorly adapted prostheses. This lesion progresses to the formation of asymptomatic fibrous tissue.

Objective: The aim to this case is to describe the clinical characteristics and to establish the treatment of this lesion.

Case Report: A 68-years-old woman was referred to go to the Oral Surgery Hospital Virgen de la Paloma. She reported a tissue growth into the oral cavity located over left buccal mucosa. No relevant family and personal history. The intraoral examination revealed a pedunculated lesion 3x4cm, whitish, appeared with a fibrous consistency and ulcerated surface. After informed consent, we proceeded to anesthesia in the region with an infiltrative local technical and subsequent cleavage by scalpel. Histopathological analysis showed dense connective tissue with few fibroblasts and blood vessels and this result confirmed epulis fissuratum. The patient returned one year later, without any signs of recurrence.

Conclusion: It is characteristic of epulis fissuratum the presence of a lesion in folds enlarged or elongated in the seat area of the prosthesis with an ulcerated or not surface and a color ranging from white to erythematous. Treatment involves surgical removal of the lesion and elimination of the causing factors always with appropriate prosthetic reconstruction (the denture must be remade or substantially adjusted).

- Oral Presentation 58

TITLE: Oral pathology in the Inflammatory Bowel diseases.

AUTHORS: Rio Rouco P. Gallas Torreira M. Barreiro De Acosta M.

Faculty of Medicine and Dentistry. University of Santiago de Compostela (USC).

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S23.

* doi:10.4317/medoral.17643948
http://dx.doi.org/10.4317/medoral.17643948

Objectives: To analyze the predominance of different oral pathologies in patients with Inflammatory Bowel diseases (IBDs), studying the distribution in patients with Crohn disease (CD) and ulcerative colitis (UC).

Material and Methods: A cross-sectional prevalence study was performed. The population of this study are patients with IBD including biological treatment protocols (Remicade® and Humira®) that which they perform a basic clinical oral examination and a survey designed to detect oral pathology.

Results: Oral lesions affect 73,53 % of patients, 50 % of patients present aphtae. gingival bleeding (49 %), gingivitis (38,23 %), periodontitis (26,47%) and the pain of Temporo-Mandibular Joint (TMJ) (26,47 %). The lesion with the lowest prevalence are the oral/dental abscesses, (11,76 %). Exist an heterogeneous distribution between the patients with CD or with UC, the prevalences are predominant for the gingivitis (32,35 %), the periodontitis (17,65 %), the gingival bleeding (38,8 %) and ATM pain (16,18%) in the CD, while in the UC more abscesses are presented (7,35 %). But only exist significantly statistically differences for gingival bleeding ($p = 0,028$).

Conclusions: Oral diseases in IBD patients, treated with Remicade® and Humira® is high. Our results indicate the existence of oral pathology, although patients are being controlled with biological treatments and in periods of remission of their disease.

- Oral Presentation 59

TITLE: Perception of the “educational climate” in students from the university of Santiago de Compostela: medicine vs dentistry

AUTHORS: Romero González C, Ruiz Casallerrey D, Arce Vázquez VM, Casares de Cal MA, Palés Argullos J, Tomás Carmona I.

Oral Sciences Research Group. School of Medicine and Dentistry, University of Santiago de Compostela, Spain.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S23.

* doi:10.4317/medoral.17643949
<http://dx.doi.org/10.4317/medoral.17643949>

Aims: To analyze and compare the “Educational Climate” (EC) of medical and dental students from the University of Santiago de Compostela (USC), using the DREEM questionnaire (“Dundee Ready Education Environment Measure”).

Material and Methods: The study group consisted of 671 medical students and 110 dental students, who responded to the DREEM questionnaire during the 2014-2015 academic year. This questionnaire has 50 items that are grouped into 5 domains: “Learning”, “Teachers”, “Academic”, “Atmosphere” and “Social”. Taking into account the maximum scores for the different domains and for EC, the data are converted into percentages of the global scale or of their respective subscale. Broadly, a higher score means a perception that is more positive than negative in relation to the educational aspect being examined.

Results: The EC score in Medicine and Dentistry were 110,7 (55.0%) and 116,5 (58.2%) respectively. The scores obtained in both careers in the different domains were: 47.3% and 52.5% in “Learning”, 51.6% and 55.9% in “Teachers”, 60.2% and 61.9% in “Academic”, 56.2% and 60.2% in “Atmosphere” and 59.1% and 62.5% in “Social”. In 17 (34%) and 12 items (24%) respectively, an average <2 was detected.

Conclusions: USC students felt that their EC was more positive than negative, because the different domains were considered “positive and acceptable”. Although the perception of the EC in Dentistry was significantly better, in both university degrees the students identified a considerable number of problematic educational aspects.

- Oral Presentation 60

TITLE: The use of the CBCT in the geriatric patient to control large apical lesions. Apropos of a case

AUTHORS: Ruiz Piñón M, Martin Cruces J, Castelo Baz P.

Master of Endodontics. University of Santiago de Compostela.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S24.

* doi:10.4317/medoral.17643950
<http://dx.doi.org/10.4317/medoral.17643950>

Introduction: Geriatric patients have a healing and bone regeneration lower than young patients. The use of CBCT technology allows us to test the healing process in these patients with a low dose of radiation.

Case Report: 69 year old male patient referred to our

office for endodontic teeth 11 and 13 and later perform a surgery at the apical region of a 12 chronic apical periodontitis and presented no previous endodontic treatment.

After talking with the patient and the dentist who referred, we decided to perform endodontic treatment in order to try to reduce the size of the apical lesion because we supposed an endodontic origin of the injury (was more than 2 cm) and affected maxillary sinus, nasal cavity with bone destruction.

After performing a root canal using ProTaper and obtured with Thermafil, radiographs and CBCT system control is performed annually, 2.3 4 and 6 below.

Conclusions: The lesion was reduced almost entirely, although healing has been slow due in part to the patient's age. The CBCT is an excellent method to evaluate and control the size reduction in this type of apical lesions.

- Oral Presentation 61

TITLE: Oral health and quality of life at the senior citizen's social clubs for people aged 65 and over in Valencia. Preliminary study

AUTHORS: Sáez Prado B, Haya Fernández MC, Sanz García MT.

Universidad Cardenal Herrera CEU, Valencia.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S24.

* doi:10.4317/medoral.17643951
<http://dx.doi.org/10.4317/medoral.17643951>

Aim: The aim of this survey was to investigate if there is a relationship between the oral health status and the oral and general health related quality of life of elderly people aged 65 years or more in Valencia, Spain, through a preliminary study.

Methods: We designed a cross-sectional oral health survey plus an oral examination in an elderly population. Subjects: 48 adults (22 men and 26 women). Age: 65 years of age and over. Setting: Randomly selected senior citizen's social clubs. We assessed the Oral Health Impact Profile (OHIP-14) and a general health-related quality of life instrument: the EuroQol-5d with a visual analogue scale (VAS). We also inquired sociodemographic and oral health questions.

Results: The mean additive score of the OHIP-14 was 10.48, the mean value of the EuroQol-5d was 0.56 and of the VAS, 70.33. We found no relationship between Euro-QoL5D and OHIP-14. Neither there were relation between these instruments and variables as decayed teeth or absences. On the other hand the OHIP-14 was consistently and significantly correlated with the oral