Hygiene (brushing, toothpaste) and the dental status (being or not edentulous).

Conclusions: The oral health has a high impact on quality of life. The oral quality of life does not seem to be related to the general quality of life. The oral hygiene and getting toothless influence negatively on the quality of life of elderly population.

- Oral Presentation 62

TITLE: Complications in prosthetic rehabilitations on implants: a retrospective observational clinical study

AUTHORS: Sanz Sánchez A, Linares Bódalo S, Buesa Báez JM, Fernández Cáliz F, Santos Marino J. *Master en Cirugía Bucal e Implantología. Hospital Virgen de la Paloma.*

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S25.

* doi:10.4317/medoral.17643952
http://dx.doi.org/10.4317/medoral.17643952

Objectives: analyze the complications in prosthesis on implants in patients rehabilitated at the Oral surgery and implantology master from Hospital Virgen de la Paloma, as well as to establish the percentage of these depending on the type of prosthesis and the time elapsed since its placement.

Material and Methods: An observational retrospective clinical study was carried out on a total of 284 stories, in which the following variables were evaluated: epidemiological data, elapsed time, type of prosthesis and complication. The results were submitted to descriptive and inferential statistical treatment using the SPSS 21.0 system for Mac OC X.

Results: 284 stories were reviewed among which, 64 patients were found with complications in their prosthetic rehabilitation on implants. The average age was 69, 2±11, 19 years old. The most used prosthesis was fixed (65.6%). The main complication (40.6%) was related to prosthetic attachments, however, no statistically significant difference was found between the types of prostheses, and complications. The average elapsed time from the placement of the prosthesis until the complication appears was 21,41 ± 34,38 months.

Conclusions: Complications on implants rehabilitations frequently affect prosthetic attachments. Therefore, it is necessary the maintenance of prosthetic rehabilitations every 6-12 months to avoid complications and to assess the condition of the periimplant tissue.

- Oral Presentation 63

TITLE: Ulcerative oral lesion on a leukoplakia. A case report

AUTHORS: Schemel Suárez M, Dávila Monzón C, López López J, Jané Salas E.

Facultad de Odontología, Universidad de Barcelona.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S25.

* doi:10.4317/medoral.17643953
http://dx.doi.org/10.4317/medoral.17643953

Introduction: Oral ulcers are very frequent and have a diverse etiology (infectious factors, trauma, neoplasia, multiple syndromes, among others). The most common cause of oral ulcers is trauma, and the treatment in this cases is the removal of the traumatic agent (sharp teeth, inadequate dental prosthesis, etc.), but when a lesion does not resolve in a 2 week period after the removal of the traumatic agent, the diagnosis will be confirmed with a biopsy, because it could be a process that needs a more complex treatment.

Case report: A 65 year old woman with a medical history of OSCC in the upper left maxilla, on a lesion years before diagnose as an oral lichen planus, attends the dental clinic for evaluation. Intraoral examination revealed an ulcer on a non-removable white plaque in the left lateral border of the tongue. Because of the patient's medical history, a biopsy and histological study was performed. The result rules out a malignant lesion but indicates the presence of viral inclusions.

Conclusions: A periodic dental evaluation is very important, especially in patients that are on chronic treatment with topical or systemic corticosteroids since the immunosuppressive action of this drug can produce oral lesions that sometimes can mimic malignant pathologies.

- Oral Presentation 64

TITLE: Oral Cancer: Risk factors in elderly. An update

AUTHORS: Schiavo Di Flaviano V, Martín C, Somacarrera ML.

Facultad de Ciencias Biomédicas. Departamento de Odontología. Universidad Europea de Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S25.

* doi:10.4317/medoral.17643954
http://dx.doi.org/10.4317/medoral.17643954

Objective: Oral cancer is a prevalent disease in the elderly. It is estimated to claim about 1117 lives a year in

Spain, where more than 70% are patients over 60 years old. The aim of this review is to identify the different risk factors involved in the development of this disease in the elderly in order to reduce their prevalence and improve morbidity and mortality.

Methods: Using the keywords “oral cavity cancer in elderly”, “oral cavity cancer” and “oral cavity cancer risk factors” a literature search of articles published in the last 10 years was performed.

Results: Cancer is a multifactorial disease which can be influenced by a variety of factors. Genetic alterations can make the patient more susceptible to carcinogens. Interactions can occur between various factors, such as diet, alcohol, tobacco, viral infections such as HPV and family history of cancer.

Conclusion: Although alcohol and tobacco are still major risk factors that must be addressed to prevent the development of cancer, advanced age has a great influence on individual susceptibility to carcinogens, so it should be considered as another risk factor of great importance. With this we must bear in mind the need and importance of our elders receive a meticulous reviews of the oral cavity often. We must not forget that the best treatment is prevention.

- Oral Presentation 65

TITLE: Differential diagnosis of soft tissues lesions. A case report

AUTHORS: Serrano Torrecilla M, Merchán Morales S, Cortés-Bretón Brinkmann J, Moncada Escoria R, Martínez Rodríguez N.

Máster en Cirugía Bucal e Implantología del Hospital Virgen de la Paloma.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S26.

* doi:10.4317/medoral.17643955
http://dx.doi.org/10.4317/medoral.17643955

Introduction: The term “epulis” was used for the first time by Virchoff in 1864 and it comes from the Greek words “epi”(on) and “oulon”(gingiva). Clinically, it is defined as a granulomatous chronic inflammatory tumor on the gingiva and related to the periosteum. The granulomatous epulis or pyogenic granuloma is presented as a benign tumor of epithelial origin, with vascular content and fast growth, which etiology is attributable to local irritants such as dental calculus, trauma, presence of root fragments or foreign bodies, and which definitive diagnosis is obtained through its histopathological study.

Case presentation: 70-years-old female patient, who came to the dental office presenting pain on the lower right side. Clinically, slight inflammation was observed in the right mandibular canine area, with negative pro-

bing and positive vitality. After making the canine root treatment, a sessile gingival lesion appeared, first with reddish and fibrous appearance and evolving towards a softer consistency with no reddish aspect, which was caused by small teleangiectasias. The total excision of the tumor was performed under local anesthesia, as well as the histopathological analysis, obtaining Pyogenic Epulis as definitive diagnosis.

Conclusions: The treatment of these lesions starts by performing a detailed clinical history and examination of the patient, being essential the histological study of all excised oral tissues in order to be able to establish its definitive diagnosis.

- Oral Presentation 66

TITLE: Denosumab: renovation and updating

AUTHORS: Sirvent Iniesta JR, González Navarro B, Arranz Obispo C, Mari Roig A, Jané Salas E, López López J.

Postgraduate Student of Oral Medicine, Oral Surgery and Implantology. Dental School. University of Barcelona, Catalonia, Spain.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S26.

* doi:10.4317/medoral.17643956
http://dx.doi.org/10.4317/medoral.17643956

Introducción: Denosumab is a fully human monoclonal antibody against the receptor activator of nuclear factor kappa-B ligand (RANKL), is an antiresorptive agent that reduces osteoclastogenesis, inhibiting osteoclast activity, reducing bone resorption and increasing bone density. Is used in the treatment of osteoporosis and bone metastases. This drug have reported associated cases of osteonecrosis of jaw (ONM).

Objective: This review is to perform a relation between treatment with denosumab and the incidence of osteonecrosis and to obtain information on dosage, first event apparition, and treatment approaches for patients with osteonecrosis related to denosumab.

Materials and Methods: A search was made at databases Pubmed, Medline and Cochrane with the keywords: denosumab, elderly, osteonecrosis, osteoporosis, bone metastases and anti-RANKL.

Results: We found a total of 139 articles. They read the summary of which is relevant to this review found 15 of them for the last five years were 4 literature reviews, 1 meta-analysis, 1 randomized placebo-controlled trial, 1 lifetime cohort Markov model and 8 cases report.

Discussion: Treatment with denosumab has a number of adverse effects including ARONJ. Although the literature cites a low incidence arising from denosumab use, when it does occur, it supposes serious functional and masticatory disorders, with a major impact on patient quality of life.