Conclusions: The attitude and tracing is still unclear, combined detection of adverse effects, antibiotic treatment for the treatment of infection combined with debridement and bone resection.

- Oral Presentation 67

TITLE: Proliferative verrucous leukoplakia. Medical case

AUTHORS: Tafache Torres R, María Colom G, Rodríguez de Rivera E, Jané Salas E, López López J.

Postgraduate student of Oral Medicine, Oral Surgery and Implantology. Dental School. University of Barcelona, l'Hospitalet de Llobregat, Barcelona, Spain.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S27.

* doi:10.4317/medoral.17643957
http://dx.doi.org/10.4317/medoral.17643957

Introduction: OPVL is a disease of unknown etiology that begins as a slow growing hyperkeratosis, which tends to spread and become multifocal and eventually develops, similar to warts, exophytic erythroplastic areas or patches with aggressive biological behavior due to its high probability of recurrence and a high rate of malignant transformation. It is more common in older women who have suffered for years leukoplakia lesions. Diagnosis is based on clinical and histopathological study of injuries. This is a very dangerous disease because of its high rate of recurrence and dangerous development.

Case: 67 years old woman, smoker of 40 cigarettes a day for over 40 years with no medical history of interest. The patient has whitish lesions, asymptomatics, of rough appearance on both jugal mucosa, hard palate, floor of the mouth, ventral surface of the tongue and tonsillar pillars since 4 years ago. We indicate to eliminate smoking habits completely, without any changes in the lesions after two weeks, it was decided to biopsy lesions of the ventral surface of the tongue and floor of mouth. The pathology revealed Stomatitis interface with severe dysplasia.

Conclusions: OPVL is a very dangerous disease because of its high percentage of malignant, difficult and controversial treatment monitoring, and questionable effectiveness.

- Oral Presentation 68

TITLE: Peripheral giant cells granuloma. Two case reports

AUTHORS: Teijeiro López P, Navarro Sangrà C, Jané Salas E, López López J.

Postgraduate student of Oral Medicine, Oral Surgery and Implantology. Dental School. University of Barcelona, Catalonia, Spain.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S27.

* doi:10.4317/medoral.17643958
http://dx.doi.org/10.4317/medoral.17643958

Introduction: Peripheral Giant Cells Granuloma (PGCG) is the giant cell lesion most common in the maxillary region. It originates in the connective tissue of the periosteum and periodontal membrane. Its etiology is still unknown, although it is usually related to irritation or aggressive factors, and diseases such as hyperparathyroidism (Brown tumour). The treatment consists of complete removal of the lesion and meticulous curettage, considering that relapse is usual.

Case reports:

Case 1: Women, 88 years old and medical history of DM1, HBP, dyslipidemia, arthralgia and gait disturbance. The patient was referred to Master in Oral Medicine, Surgery and Implantology (University of Barcelona, Catalonia, Spain). She presented a single, asymptomatic, exophytic lesion measuring approximately 1.5x1 cm in diameter, situated next to right central incisor. The patient reveals one year of evolution and the lesion was related to a prosthodontic restoration.

Case 2: Man, 53 years old without relevant medical history. The patient was referred to the same department as the previous patient. He reported a solitary and exophytic lesion situated in hard palate between 1.6, 1.7 and 1.8. Evolution time of approximately one year, about 3x2 cm in diameter and was probably related to endodontic treatment.

Conclusions: Granulomatous lesions situated in soft tissues require differential diagnosis with other entities. The treatment of choice is excisional biopsy which helps to confirm the final diagnosis.

- Oral Presentation 69

TITLE: Most common oral pathology in the elderly. Study in patients from the Polyclinic of the European University of Madrid

AUTHORS: Turégano Alarcón BV; Moquillaza González LE; Martín C, Somacarrera Pérez ML.

Departamento de Odontología. Facultad de Ciencias Biomédicas. Universidad Europea de Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S27.

* doi:10.4317/medoral.17643959
http://dx.doi.org/10.4317/medoral.17643959

Aim. To make a cross – sectional and descriptive study of the most common oral diseases in the elderly. Compare the results with the current literature and other studies conducted in the last years.

Methods. We took a sample of 100 patients over 65 years old from the Polyclinic of the European University of Madrid. The oral symptoms were reported, in most of cases consisting of xerostomia associated to drugs, denture ulcers, gingival bleeding associated with anticoagulant therapy and tooth mobility among others. We also studied risk factors, systemic diseases and prescribed medication as well as oral hygiene and used or worn prosthetic elements. As far as oral pathology, we performed an extra and intraoral exploration. We searched for ulcerative lesions, lingual, viral, bacterial, fungal and autoimmune pathology among others, associated to poor hygiene, caries, gingival recession and periodontal disease.

Results. Traumatic ulcer associated to prosthesis in poor condition was the most common oral lesions in the study group. The most common symptom was xerostomia associated with the consumption of drugs.

Conclusion. Increased life expectancy and declining birth rates are responsible for the aging of population in Spain and in other developed countries. The dentist and other health professionals must recognize this social change and focus actions to the prevention and detection of oral lesions and symptoms caused by systemic disorders.

- Oral Presentation 70

TITLE: Short implants for atrophic maxillary restoration. A systematic review

AUTHORS: Veiga López B, Cabezas Mojón J, Guenín Gutiérrez CV, Loughney González A, Fernández Domínguez M.

Master en Cirugía Bucal e Implantología. Universidad San Pablo CEU. Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S28.

* doi:10.4317/medoral.17643960
http://dx.doi.org/10.4317/medoral.17643960

Objective: To assess studies concerning short implants (<10 mm) as an alternative to standard implants (≥10 mm) in the treatment of atrophic jaws. These cases always present longitudinal bone resorption.

Material and Methods: A Pubmed-Medline research of 20 articles published in the last 10 years was undertaken. We have used the following key words: "short implant", "partial edentulism", "survival implant", "implant length", "surface topography".

Results: Short implant's success/failure rate and marginal bone have been evaluated in comparison of standard implants and both reach similar outcomes. Authors agree with the fact that treatment's prognosis improves when short implants are installed on the mandible instead of the maxilla.

tead of the maxilla. Implant's surface treatment also has influence considering that roughed surface implants get better results than mechanized surface ones.

Conclusion: Short implants (<10 mm) are a reliable and predictable treatment alternative in the prosthetic rehabilitation of atrophic jaws. In posterior areas of the maxilla and mandible, height alveolar bone availability is usually reduced. This mean in a reduction of treatment time, costs and patient's morbidity in comparison with bone augmentation surgical techniques.

- Oral Presentation 71

TITLE: Oral candidiasis diagnosis. Smear vs. culture

AUTHORS: Vilar Villanueva M; Yáñez Busto A; Durán Somacarrera J; Blanco Carrión A.

Master in Oral Medicine, Oral Surgery and Implantology. Faculty of Medicine and Dentistry of Santiago de Compostela University.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 Junio 1;20(Supplement1):S28.

* doi:10.4317/medoral.17643961
http://dx.doi.org/10.4317/medoral.1764396

Objectives: To compare culture and smear for the oral candidiasis diagnosis.

Material and Methods: 80 patients from the Master in Oral Medicine, Oral Surgery and Implantology from the University of Santiago de Compostela, who had signs, symptoms and /or predisposing factors commonly related to oral candidiasis, were selected. Clinical and pharmacological data, systemic diseases, denture use and Whole Saliva Test (WST) were evaluated. Additionally, culture and smear were taken from all patients. The culture was carried out using a cotton swab and a Sabourau Agar Petri dish. The smear was obtained using a cement spatula and a microscope slide. Finally, both tests were sent to the Pathology Department at the University Hospital of Santiago de Compostela to be analyzed.

Results: Within the 80 patients studied, 50% of them obtained positive culture and 20% positive smears. 100% of the patients who obtained positive smears also had positive cultures. 30% of the cases showed positive cultures and negative smears, which means colonization but not infection.

Conclusions: According to our data, culture cannot be taken as a reliable diagnostic test due to its high percentage of false positives. We consider smear as the technique of choice for oral candidiasis diagnosis.