Materials and Methods: A search is realized in the databases of PudMed, MedLine and Cochrane by means of the key words: “Peri-implantitis“, “Infection peri-implantitis”, “Factors of risk”, “Treatment peri-implantitis”. More excellent publications index-linked between the year 2010 and 2015 were included.

Results: The 14 most influential articles were selected. The evidence review that there are many possible risk factors that may relate to the peri-implant disease : poor oral hygiene , smoking , history of periodontitis , diabetes , genetic factors, etc. Probing depth , bleeding and oozing should be regularly assessed for the diagnosis of peri-implantitis ; and radiographs to assess levels of peri-implantary bone.

Conclusions: It is convenient to reach consensus on diagnostic criteria and definition of peri order to obtain meaningful conclusions about prevalence and severity, and this make an effective protocol for action against perimplantitis, however prospective studies are required.

- Poster 11

TITLE: Diagnosis and treatment of autoimmune blistering diseases

AUTHORS: Gual Vaqués P, Roselló Camps A, Egido Moreno S, Marí Roig A, López López J, Jané Sala E. *Máster Medicina, Cirugía e Implantología Oral. Universidad de Barcelona.*

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S34.

* doi:10.4317/medoral.17643974
http://dx.doi.org/10.4317/medoral.17643974

Introduction: Autoimmune blistering diseases comprise a heterogeneous group of diseases that are characterized by autoantibodies directed against adhesion molecules of the skin and adjacent mucous membranes – including lichen planus, lupus erythematosus, pemphigoid, pemphigus vulgaris, exudative erythema multiforme, herpetiform dermatitis and linear IgA disease.

Aim: A review of the clinical manifestations of these diseases, with an analysis of the diagnosis and treatment modalities most commonly employed in such patients.

Materials and Methods: An electronic literature search was conducted by two reviewers (PG and AR) in several databases, including MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials and Cochrane Oral Health Group Trials Register databases for articles written in English from January 2010 up to April 2015.

Results: The diagnosis of autoimmune bullous diseases is by clinical suspicion of injuries, and confirmed by

biopsy and determination of antibodies in the blood or tissues. Immunoprecipitation studies are currently considered one of the most definitive.

The absence of long-term controlled studies to assess the efficacy of treatments has led to the use of a wide variety of drugs, being topical or systemic corticosteroids the most commonly used.

Conclusions: Knowledge of these disorders is important in elderly, particularly because the population is aging, and as the immune system evolves with senescence, immunobullous diseases increase in incidence.

- Poster 12

TITLE: Clinical characteristics of burning mouth syndrome (BMS) in elderly patients. Comparative study with younger patients of a total population of 183 patients of an Oral Medicine clinics

AUTHORS: Henriques I, Cardoso I, Trancoso P, Azul AM.

CIMO- Department of Oral Surgery and Medicine – Lisbon- Portugal.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S34.

* doi:10.4317/medoral.17643975
http://dx.doi.org/10.4317/medoral.17643975

Aims: To evaluate the characteristics of BMS in an elderly patients population (≥ 65 y) and compare with younger patients of the same Oral Medicine clinic and the international bibliography.

Material and Methods: Retrospective study by evaluation of patients' clinical records.

Results: 183 files of patients with BMS were evaluated. Of these 96 elderly (52.5%) (male = 14.6% , female= 85.4%) and 87 younger patients (47.5%) (male=14.9% , female= 85.1%)

Time between the beginning of the symptoms and final diagnosis was up to 6 months in 33.3%, up to 2 years in 41.7% and more than 2 years in 18.8% of the elderly group. With the younger patients percentages were 41.8%, 39.0% and 13.8% respectively. Of the elderly group 23% had psychiatric treatment (14.4% medicated with psycho drugs) and the same occurred in 15% of the younger patients (11.5% medicated with psycho drugs), without proper diagnosis. Also all this period 30% of the elderly patients were already medicated for their symptoms mainly with antifungal drugs (48.2%) as well as submitted to several useless diagnostic tests. At the younger group percentages are 24 % y 52.4% respectively

Conclusions: Our patients spent much time in specialties' consultations without a proper diagnosis, but

submitted to several tests and treatments with none efficacy and with a high economic and social cost until the definitive diagnosis of BMS. It is mandatory that, in Portugal, medical general practitioners and other medical specialties recognize BMS, for a correct diagnosis with no useless delay.

- Poster 13

TITLE: Influence of oral bisphosphonates in patients with dental implants

AUTHORS: Jalil-Abumalham D, Hurtado Celotti D, Buesa Báñez JM, Rubio Alonso L, Martínez-González JM.

Master en Cirugía Bucal e Implantología. Hospital Virgen de la Paloma.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S35.

* doi:10.4317/medoral.17643976
http://dx.doi.org/10.4317/medoral.17643976

Introduction: Bisphosphonates (BPs) are a class of agents used to treat osteoporosis and malignant bone metastases, among others. These drugs have been associated with a serious adverse reaction known as bisphosphonate-related osteonecrosis of the jaw (BRONJ) despite the benefits of this drug. The prevalence of BRONJ in patients receiving long term oral BPs was reported at 0, 0007 % per 100.000 patients (< 1 case per 100.000 exposed). The aim of this work it's to explain the association between the use of oral bisphosphonate therapy and dental implant failure.

Case presentation: This report documents a case of BRONJ after two dental implant placement in a osteoporotic patient (female/66 years) treated with ibandronic acid taken orally once a month for eight years. After clinical examination and an appropriate radiographic study, peri-implant radiolucency is observed which shows the failure of osseointegration of dental implants.

Conclusion: Despite the low risk of BRONJ occurrence after implant surgery in oral bisphosphonates users, the fate of dental implants may be affected to the type of drug used and the time of administration as well as the role of potential infection at site of surgery and hard tissue management.

- Poster 14

TITLE: Oral manifestations of syphilis. Systematic review

AUTHORS: León Blanco A, Arranz Opispo C, Roselló Llabrés X, Jané Salas E, López López J, Marí Roig A.

Máster Medicina, Cirugía e Implantología Oral, Universidad de Barcelona, Cataluña, España.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S35.

* doi:10.4317/medoral.17643977
http://dx.doi.org/10.4317/medoral.17643977

Objective: To review the epidemiology and oral manifestations of syphilis, highlighting the importance of knowing the dentist, who will diagnose and refer the patient to a specialist to apply the appropriate treatment.

Materials and Methods: An electronic search was performed in Pubmed data base. Using the terms "oral manifestations", "Syphilis". Duplicates are removed, take those published in the last five years, we discard those not included epidemiological data, statements in oral cavity and obtained a total of 10 items.

Results: In recent years, there have been reporting outbreaks in the United States, Canada, Russia, China, Central and Eastern Europe. This resurgence of the disease could be due to the high rate of promiscuity, lack of knowledge of the disease, the booming sex industry, decreased use of condoms and the sense of security that today Sexually Transmitted Diseases (STDs) are cured. It is well known that STDs are more common in the younger population, although the risk of STDs has increased in adults possibly due to the arrival of drugs for erectile dysfunction.

Conclusions: Because oral lesions are highly contagious, it is vital to make the correct diagnosis quickly, to initiate appropriate therapy and interrupt the chain of infection. The demonstration in oral cavity should be considered in the differential diagnosis of unusual oral ulcers and white patches that affect any patient.

- Poster 15

TITLE: Nasal floor elevation in the rehabilitation of atrophic maxilla with dental implants

AUTHORS: Lois Franco A, Oliván Molina S, Adarve Pérez A, Loughney González A, Fernández Domínguez M.

Master en Cirugía Bucal e Implantología. Universidad San Pablo-CEU. Madrid.

SOURCE: Med Oral Patol Oral Cir Bucal. 2015 June 1;20(Supplement1):S35.