“Efficacy of aloe vera dressing in management of decubitus ulcer”

Mr Rajveer Singh¹ Dr.T.Bhattacharjee²

1 Phd Scholar, Nims University Rajasthan Jaipur

2 PhD Guide, Nims Nursing College, Nims University

Corresponding Email: creativeraj_singh@yahoo.com

Mail Guide; dr.bhattacharjee@gmail.com

ABSTRACT

Introduction: Pressure ulcer is a continuing major health problem worldwide. The development of pressure ulcer wound is more prone to those patients who are confined to bed for long periods. Pressure ulcers represent a major burden of sickness and reduced quality of life for patients and their caregivers. The aim of the study was to assess efficacy of aloe vera dressing in management of decubitus ulcer among bedridden patients.

Material & Methods: Pre-experimental, two group pre-test post-test research design ($O_1 \times O_2$) was used. Total 200 patients of decubitus ulcer wound were taken by convenience sampling technique. Result: There was a significant difference between pre test and post test score of decubitus ulcer wound healing.

Conclusion: The result from the study reveals that aloe vera gel is effective in decubitus ulcer wound healing.

Keywords: Decubitus ulcer, Aloe vera Gel, Bed ridden patients

Email: INTRODUCTION:

Pressure ulcer also known as bedsores which usually occur over a bony prominence due to long-term pressure, or pressure along with friction. The most common sites for developing pressure ulcer are the sacrum, coccyx, heels, and hips, and least common sites include the elbows, knees, ankles, back of shoulders, or the back of the cranium. Pressure ulcers occur due to pressure which completely or partially obstructs blood flow to the soft tissue. Pressure ulcers most commonly develop in person who is not moving about or on chronic bed rest.

NEED FOR STUDY:

Aloe Vera contains almost 75 known compounds which includes 20 minerals, 20 amino acids, water and vitamins.

Early wound healing is very important for patients suffering from the pressure ulcers. When pressure ulcer occurs, the length of hospital stay and overall cost of treatment increase, and treatment of pressure ulcers become more costly, than prevention, when an

ulcer develops, the increased cost of nursing care alone is estimated as 50%.

STATEMENT OF PROBLEM:

An Experimental study on “Efficacy of aloe vera dressing in management of decubitus ulcer among bedridden patients admitted in selected hospitals at Jaipur”

OBJECTIVES:

- To assess the decubitus ulcer among the bedridden patients in experimental and control group.
- To determine the effectiveness of aloe vera gel in management of decubitus ulcer among bedridden patients
- To compare management of decubitus ulcer among the bedridden patients in the control and experimental group
- To find out the association between management of decubitus ulcer with selected demographic variables

HYPOTHESIS:

H1: The Aloe Vera Gel dressing will have a significant effect in healing the pressure ulcers.

H2: There will be significant association between healing of pressure ulcers and selected socio-demographic variables.

MATERIAL AND METHODS:

In this study, the quantitative research approach was used, the investigation aims at evaluating the efficacy of aloe vera dressing in management of decubitus ulcer among bedridden patients.

Research Design:-The research design was a pre-experimental one group pre-test and post-test design.

Population:-The population was comprised of decubitus ulcer patients admitted in selected hospitals.

Sample size:-200 bedridden patients diagnosed with decubitus ulcer patients from selected hospital Jaipur.

Sampling technique:-Non-random, convenient sampling technique

Criteria for selection of sample:-

Inclusion criteria

- Age group above 30 years
- Subjects of both sex
- Bedridden patients with pressure ulcers.

Exclusion criteria

- Subjects who are suffering from chronic illnesses like diabetes mellitus, tuberculosis, anemia, vascular diseases, septicemia and mental diseases.
- Subjects who had been previously treated with other topical medicine or antibiotics.

Development and Description of Tool:-Data collection tools are the procedures or instruments used by the researcher to observe or measure key variables in the research problem.

Validation of Tool:-

Standardised tool e.i. Bete Jensen Scale for pressure ulcer healing assessment was used in this study.

Reliability of the Tool:-

Reliability denotes degree of consistency. The reliability of the tool was assessed by using inter rater method and its correlation coefficient r -value is 0.84. This correlation coefficient is high and it is good tool for assessing effectiveness of Aloe vera gel dressing in healing of pressure ulcers among subjects

Data collection:-

A formal written permission was obtained from the head of the departments of Medical, Surgical and

Orthopedics wards of Saket Hospital, and selected hospitals Jaipur. By using Non-random, convenient sampling technique three or four subjects were selected on every day. Subjects selected for Pilot study were excluded. The subjects for the study were selected based upon the of Bates Jensen wound assessment tool. The study purpose and method were explained to all subjects and informed written consent was obtained. Confidentiality was assured to all the subjects. The information regarding demographic profile and pre wound assessment related information were collected from 200 pressure ulcer subjects and observing health records. 100 subjects were included for both experimental and control group.

RESULTS

Frequency and percentage distribution of socio demographic variables in experimental and control group. Regarding the age, majority of the subjects 42% were between 31-40 years in experimental group and majority of subjects 43% were between 41-50 years in Control group.

sex majority of subjects 63% were males in Experimental group and

majority of subjects Considering the sex 43% were females in Control group. In considering the education, majority of subjects 48% were having education of senior secondary level in Experimental group and majority of subjects 44% were having matric level education in Control group.

In occupation, majority of subjects 72% were skilled worker in Experimental group and most of the subjects 71% were skilled worker in Control group. As for the income, majority of subjects 47% get a monthly income of Rs.10000-20000 in Experimental group and most of the subjects 41% get a monthly income of Rs.10000-2000 in Control group. Considering the area of living, majority of subjects 77% belonged to Urban area in Experimental group whereas most of the subjects 60% belonged to Urban area in Control group. Regarding length of hospital stay, majority of subjects 49% were stayed in hospital for more than 21-30 days in Experimental group and most of the subjects 40% were stayed in hospital for 11-21 days in Control group. Experiment subjects, 75% were having none/others group of diseases whereas in control group 88% were having none/others group of diseases. Regarding body built majority of subjects 57% were obese in experimental group and most of the subjects 46% were moderate in control group.

N=200

Demographic variables		Group			
		Experiment		Control	
		N	%	N	%
Age in years	21-30	10	10	9	9
	31-40	42	42	39	39
	41-50	29	29	43	43
	Above 50	19	19	9	9
Sex	Male	63	63	57	57
	Female	37	37	43	43
Education	Illiterate	2	2	5	5
	Primary	7	7	5	5
	Matric	27	27	44	44
	S. Secondary	48	48	38	38
Occupation	Skilled worker	72	72	71	71
	Unskilled worker	28	28	29	29
Family income Rs. Per month	Less than 5000	3	3	7	7
	5,000-10,000	11	11	33	33
	10,000- 20,000	47	47	41	41
	More than 20000	39	39	19	19
Area of living	Rural	77	77	60	60
	Urban	23	23	40	40
Length of hospital stay	10 days	1	1	15	15
	11- 20 days	33	33	40	40
	21-30 days	49	49	29	29
	More than 30 days	17	17	16	16
Associated diseases	Diabetes	18	18	7	7
	Hypertension	3	3	4	4
	Anemia	4	4	1	1
	Fracture/ others	75	75	88	88
Body Built	Thin	15	15	0	0
	Moderate	46	46	43	43
	Obese	39	39	57	57

Table shows the frequency and percentage distribution of samples according to their selected socio demographic variables in experimental and control group.

To assess the decubitus ulcer among the bedridden patients in experimental and control group.

In pre test distribution of bedridden patients in experimental group were having the mean of 47.62 with SD of 3.06 and in control group the mean of bedridden patients were 48.56 with SD of 2.42.

In post -test distribution of bedridden patients in experimental group were having the mean of 8.06 with SD of 3.54 and in control group the mean of bedridden patients were 16.88 with SD of 1.98.

N=200

	Pre Test	
	MEAN	SD
Experimental group	47.62	3.06
Control Group	48.56	2.42

Table Frequency and percentage distribution of bedridden patients in experimental and control group as per pre-test status of decubitus ulcer

N=200

	Post Test	
	MEAN	SD
Experimental group	8.06	3.54
Control Group	16.88	1.98

Frequency and percentage distribution of bedridden patients in experimental and control group as per pre-test status of decubitus ulcer.

Comparison of management of decubitus ulcer among the bedridden patients in the control and experimental group.

In experimental group, mean post-test score (8.06) with SD (3.54) was lower than mean pre-test score (47.62) with SD (3.06) which shows aloe Vera dressing application is effective on wound healing.

N=200

	CONTROL GROUP		EXPERIMENTAL GROUP	
	MEAN	SD	MEAN	SD
PRE-TEST	48.56	2.42	47.62	3.06
POST-TEST	16.88	1.98	8.06	3.54

Comparison of management of decubitus ulcer among the bedridden patients in the control and experimental group.

Effectiveness of aloe vera gel in management of decubitus ulcer among the bedridden patients in experimental and control group.

That the mean in experimental group (8.06) was lower than mean in control group (16.88). Calculated 't' value 21.89 was more than tabled 't' value 1.98 at the degree of freedom 198. Calculated p value is .00001. Both t and p value show the significant difference between control group wound assessment score and experimental group wound assessment score at post-test. Calculated Hedge's G is 2.81 which showed that aloe vera dressing application cause large effect on wound healing which can be observed easily.

Association of post-test decubitus ulcer status among bedridden patients with selected socio-demographic variables.

That in control group, post-test decubitus ulcer status was significantly were not significantly associated with socio-demographic variables age, sex, education, occupation, family income, length of hospital stay, associated diseases and body built.

That in experimental group, post-test decubitus ulcer status was significantly were not significantly associated with socio-demographic variables age, sex, education, occupation, family income, length of hospital stay, associated diseases and body built.

DISCUSSION:

In present study the pre- test, distribution of bedridden patients in experimental group were having the mean of 47.62 with SD of 3.06 and in control group the mean of bedridden patients were 48.56 with SD of 2.42.

In the present study post -test distribution of bedridden patients in experimental group were having the mean of 8.06 with SD of 3.54 and in control group the mean of bedridden patients were 16.88 with SD of 1.98.

CONCLUSION:

This study was to evaluate the efficacy of Aloe Vera Gel in management of pressure ulcers among bedridden subjects admitted at selected hospitals Jaipur. The findings of the study showed that Aloe Vera Gel dressing was effective in healing of pressure ulcers in subjects. There was association between healing of pressure ulcer and demographic variables. As complementary therapy which is effective in healing of pressure ulcers in bedridden subjects without any complications, it can be adopted by every health care professional to improve healing and reduce the time period of healing among subjects with superficial pressure ulcers.

REFERENCES:

1. https://en.wikipedia.org/wiki/Pressure_ulcer pressure ulcer affects more than 3.1 million adults worldwide annually .Ghanee R, Gavami H. Pressure ulcer in intensive care unit. Bimonthly J Urmia Nurs Midwifery Fac. 2010;8(2):90–103.
2. <https://patient.info/doctor/pressure-ulcers-pro>
3. Tatiana V. Boyko,1,,2 Michael T. Longaker,1,,3 and George P. Yang.1Review of the Current Management of Pressure Ulcers.Adv Wound Care. 2018 Feb 1; 7(2): 57–67.
4. Patient risk factors for pressure ulcer development: Systematic review International Journal of Nursing Studies Volume 50, Issue 7, July 2013, Pages 974-1003
5. Vanderwee K, Clark M, Dealey C, et al. Pressure ulcer prevalence in Europe: a pilot study. J Eval Clin Pract. 2007;13(2):227–235.
6. Jiang Q, Li X, Qu X, et al. The incidence, risk factors and characteristics of pressure ulcers in hospitalized patients in China. Int J Clin Exp Pathol. 2014;7(5):2587–2594.
7. Bredeisen IM, Bjørø K, Gunningberg L, et al.

- Patient and organisational variables associated with pressure ulcer prevalence in hospital settings: A multilevel analysis. *BMJ Open*. 2015;5:8.
08. v.s. chauhanat.el. The prevalence of pressure ulcers in hospitalized patients in a university hospital in India. Article in *Journal of Wound Care* 14(1):36-7 •
 09. Jiang Q, Li X, Qu X, et al. The incidence, risk factors and characteristics of pressure ulcers in hospitalized patients in China. *Int J ClinExpPathol* 2014; 7(5): 2587-94
 10. Moore Z, Johanssen E, van Etten M. A review of PU prevalence and incidence across Scandinavia, Iceland and Ireland (Part I). *J Wound Care* 2013; 22(7): 361-362, 364-
 11. Ensieh Ramezanpour¹, Amir Emami Zeydi¹, Incidence and risk factors of pressure ulcers among general surgery patients *Journal of nursing and midwifery sciences*, year 2018 volume :5 issue:4 page 159-164
 12. Pressure ulcer summit 2018: An interdisciplinary approach to improve our understanding of the risk of pressure-induced tissue damage Lisa J. Gould MD, PhD <https://onlinelibrary.wiley.com/doi/abs/10.1111/wrr.12730>
 13. Biji M Sankaran¹, Santam Chakraborty², Burden and outcomes of pressure ulcers in cancer patients receiving the Kerala model of home based palliative care in India: Results from a prospective observational study ,Year : 2015 | indian journal of palliative care Volume : 21 |
 14. The Burden of Pressure Ulcers in Spain. Soldevilla Agreda JJ, Torra I Bou JE, *Wounds : a Compendium of Clinical Research and Practice* [01 Jul 2007, 19(7):201-206]
 15. Pressure ulcer summit 2018: An interdisciplinary approach to improve our understanding of the risk of pressure-induced tissue damage Lisa J. Gould CS, Gregory Bohn A, M, Ruth Bryant N, First published: 23 May 2019 <https://doi.org/10.1111/wrr.12730>
 16. Thomas DR. The new F-tag 314: prevention and management of pressure ulcers. *J Am Med Dir Assoc* 2006; 7:523.
 17. Qixia Jiang,¹ Xiaohua Li,¹ Xiaolong Qu. The incidence, risk factors and characteristics of pressure ulcers in hospitalized patients in China. *Int J ClinExpPathol*. 2014; 7(5): 2587–2594.. /
 18. Pressure ulcers: a renewed awareness. Baranoski S *Nursing*. 2006 Aug; 36(8):36-41; quiz 42.
 19. Soozani A, Raei M, Montazeri A, Nasiri E, Hamidzadeh A, Pourheidari M. The effect of education on knowledge and performance of nurses in prevention and control of pressure sore. *Mod Care Scient Quart Birjand Nurs Midwifery Fac*. 2012;9(1):16–
 20. Potter P, Perry A, Stockert P, Hall A. *Fundamental Nursing*. Vol. 2, Chapter 48. 2013. pp. 1176–1232.