“A STUDY TO ASSESS THE KNOWLEDGE AND PRACTICE OF FEMALE HEALTH WORKER (FHW), REGARDING IDENTIFICATION OF HIGH RISK PREGNANCY, WORKING AT SELECTED SUB CENTERS UNDER BLOCK PRIMARY HEALTH CENTER, SIRSI AND AMER IN JAIPUR DISTRICT, RAJASTHAN.”

Mr. Narshi Ram Yadav

PhD Scholar JJTU. M.Sc. (N), Obstetrics & Gynecological Nursing, Faculty,
Govt. College of Nursing, Jaipur, Rajasthan, India

Corresponding Email Id- narshiyadav255@gmail.com

Abstract

Introduction: - Women are considered as “vulnerable” or “special risk” group because of their child bearing state. In every minute, a women dies somewhere from pregnancy related causes. Maternal conditions are the 7th leading cause of death in adults aged 15-49 years as per WHO estimates. In India, Community Auxiliary Health Worker and Public Health Nurse or Female Health Worker (FHW) have been given the responsibility of identifying and dealing with high risk pregnant women in rural area.

Method & Material: The descriptive design was used in this study. 50 Female health workers worked at selected sub-centers, Jaipur were sampled through random sampling technique was used. A structured knowledge questionnaire was developed to assess knowledge regarding identification of high risk pregnancy and a checklist for practices of FHWs regarding identification of high risk pregnancy Result: The findings reveal that 76% Female Health Workers have adequate knowledge and 66% have appropriate practice regarding identification of high risk pregnancy. There is significant correlation between knowledge and in-service training and practice and age, in- service training, residence and type of appointment. There is highly positive correlation between knowledge and practice of Female Health Worker regarding identification of high risk pregnancy. Based on these finding it can be concluded that the Female Health Worker needs more competency, based on in-service training to improve their skill.

Conclusion: The FHWs have lack of knowledge & practice regarding identification of high risk pregnancy. So for awareness among FHWs regarding identification of high risk pregnancy, conduct a Orientation programme for FHWs regarding identification of high risk pregnancy.

Key words: Female health Worker, High risk pregnancy, Sub centre Sirsi & Amer, Jaipur.

INTRODUCTION:

Women are considered as “vulnerable” or “special risk” group because of their child bearing state. In every minute, a women dies somewhere from pregnancy related causes. Maternal conditions are the 7th leading cause of death in adults aged 15-49 years as per WHO estimates.

A pregnancy can be considered a high risk for a variety of reasons, called “risk factors”. These factors can be divided into Maternal and Fetal factors.

In pursuance of the National Health Policy 1983, a number of health programmes were launched in India to improve health of mothers and children.

The objectives of all these programmes were convergent and aimed at improving the health of mothers and young children, and to provide them facilities for prevention and treatment of major diseases.

The health care professions have been utilizing the concept of risk to define population requiring special kinds of medical intervention, nursing care, and complex facilities and equipments. Screening for risk which is a continuous process in more careful surveillance and anticipatory guidance of the risk prone mothers. It also helps in making decisions by the health care providers and avoidance of unnecessary interventions and cost effective care.

In India, Community Auxiliary Health Worker and Public Health Nurse or Female Health Worker (FHW) have been given the responsibility of identifying and dealing with high risk pregnant women in rural area. Thus the FHW, who are responsible for majority of delivery, must have adequate knowledge, which will definitely help to improve their practice and assist in the reduction of maternal and neonatal mortality and morbidity in this country.

STATEMENT OF THE PROBLEM:

“A study to assess the knowledge and practice of Female Health Worker (FHW), regarding identification of high risk pregnancy, working at selected sub centers under Block Primary Health Center, Sirsi and Amer in Jaipur district, Rajasthan.”

OBJECTIVES:-

1. To assess the knowledge of FHW regarding identification of high risk pregnancy.
2. To observe the practice of FHW regarding identification of high risk pregnancy.
3. To find out the relationship between knowledge of FHW and their practice in identification of high risk pregnancy.
4. To seek association between knowledge of FHW in identification of high risk pregnancy with selected variables i.e. age, academic qualification, working experience, residence, participated in in-service training programme, last date of attending in-service training and type of appointment.
5. To seek association between practice of FHW in identification of high risk pregnancy with selected variables i.e. age, academic qualification, working experience, residence, participated in in-service training programme, last date of attending in-service training and type of appointment.

RESEARCH HYPOTHESIS

H1: -There will be significant co-relation between knowledge score and practice score of Female Health Worker regarding identification of high risk pregnancy.

H2: -There will be significant association between knowledge of Female Health Worker regarding identification of high risk pregnancy and selected demographic variables such as- age, academic qualification, working experience, in-service training, last date of attending in- service training and type of

appointment.

H3: -There will be significant association between practice of Female Health Worker regarding identification of high risk pregnancy and selected demographic variables such as- age, academic qualification, working experience, in-service training, last date of attending in- service training and type of appointment.

METHODOLOGY

The descriptive design was used in this study. 50 Female health workers worked at selected subcentres, Jaipur were sampled through random sampling technique was used. A structured knowledge questionnaire was developed to assess knowledge regarding identification of high risk pregnancy and a checklist for practices of FHWs regarding identification of high risk pregnancy

STATISTICAL ANALYSIS

Data was analyzed by using percentage, mean, mode, median and standard deviation c.v. etc. Chi square test was used to find out association between Knowledge and Practice of FHWs regarding identification of high risk pregnancy.

RESULTS

The study was conducted in selected sub centers under Block Primary Health Centers Sirsi and Amer in Jaipur district, Rajasthan. The subjects of the study were fifty Female Health Workers. Data obtained were analyzed and interpreted by using descriptive and inferential statistics. Data related to personal background were analyzed in terms of frequency and percentage distribution. Chi- square (χ^2) test is applied to interpret the significant association. The study finding revealed that, maximum numbers of Female Health Workers belonged to the age group ≥ 36 yrs (46%). Majority were secondary (52%). 58% were working since ≥ 10 years and maximum number lived out side the sub center campus but within the villages of sub center (56%). 68% had taken in-service training, most of them received it within one year and 74% FHWs are permanent. According to knowledge score majority of Female Health Worker are in below 70% (mean Knowledge score is 68.8% and range of score is 40-80%) Whereas according to practice scores majority falls in group below 60% (mean practice score is 54.14% and range of score is 32.5-70%). Knowledge

score above 50% is considered as adequate Knowledge and below 50% is considered as inadequate knowledge. Regarding practice score, appropriate practice is considered above 50% score and inappropriate below 50% score.

Distribution of FHWS according to their knowledge score and practice score.

This section describes the distribution of female health workers according their knowledge and practice score in term of frequency and percentage.

Table No.1

Frequency and Percentage distribution of knowledge scores level

N=50

Score	Level	Frequency	Percentage
< 50%	Inadequate	12	24%
50 or more	Adequate	38	76%

Data represented in table indicates that majority of study participants (76%) were adequate in knowledge regarding identification of high risk pregnancy.

Figure:- Bar diagram shows samples distribution [in %] participated in study according to their knowledge level.

Table No.2

Frequency and Percentage distribution of Practice scores level

N=50

Score	Level	Frequency	Percentage
< 50%	Inappropriate	17	34%
50% or more	Appropriate	33	66%

Data represented in table indicates that majority of study participants (66%) were appropriate in practices regarding identification of high risk pregnancy.

Figure:- Bar diagram shows samples distribution [in %] participated in study according to their practices level.

Table No.3

Mean and Standard Deviation of Knowledge and Practice scores

N=50

Measure	Knowledge score	Practice score
Mean	15.56	21.54
Standard Deviation	2.55	3.82

Table No. depicted that Mean Knowledge and Mean Practice scores are 15.56 and 21.54 respectively. Both are above 50%.

SD of knowledge scores is ± 2.55 and SD of practice scores is ± 3.82

76% Female Health Workers have adequate knowledge and 66% have appropriate practice regarding identification of high risk pregnancy. There is significant correlation between knowledge and in-service training and practice and age, in-service training, residence and type of appointment. There is highly positive correlation between knowledge and practice of Female Health Worker regarding

identification of high risk pregnancy. Based on these finding it can be concluded that the Female Health Worker needs more competency, based on in-service training to improve their skill.

CONCLUSION:

The FHWs have lack of knowledge & practice regarding identification of high risk pregnancy. So for awareness among FHWs regarding identification of high risk pregnancy, conduct a Orientation programme for FHWs regarding identification of high risk pregnancy.

REFERENCES:

1. Elizabeth Stepp Gilbert, (2007). Manual of high risk pregnancy and delivery, (4th ed.). Elsevier India Private Ltd. New Delhi, India. P.P. 25-42.
2. D.K. James, et al.(2006). High risk pregnancy management options, (3rd edition). Elsevier India Pvt. Ltd. New Delhi India. P.P. 3-10.
3. The White Ribbon Alliance, A Field Guide For Implementing Best Practices In Safe Motherhood, The White Ribbon India.
4. Evans, A. T., and K. R. Niswander, (2000). Manual of Obstetrics, (6th ed.) Hagerstown, MD. Lippincott Williams and Wilkins,.
5. Klossner, N.J. (2006). Introductory Maternity Nursing. (1st ed.), Texas Lippincott Williams and Wilkins.
6. Anderson, J. & Dursten, B. H, "Thesis and Assignment writing New Delhi", Wiley eastern Ltd,1989

■ ■ ■