A COMPARATIVE STUDY TO ASSESS THE PREVALENCE RATE OF CAD AMONG TYPE 2 DM PATIENT AND NON TYPE 2DM PATIENT IN SELECTIVE HOSPITALS ETAWAH DISTRICT.

Devanand¹, Lokesh Tyagi², Anupam³, Mr. Krishna Kant⁴

¹B.Sc. Nursing Scholar, Faculty College of Nursing, UPUMS, Saifai, Etawah

^{2,3,4}Faculty College of Nursing, UPUMS, Saifai, Etawah

Corresponding Email: tyagikk2018@gmail.com

ABSTRACT

Introduction: Coronary artery disease (CAD) is a major cause of death in the western countries and it is the becoming major cause of death in developing countries. This increase may be due to the rising prevalence of many risk factor such as diabetes mellitus, which is one of the most important of these risk factor .the prevalence of diabetes mellitus increasing globally and it has reached in the worldwide.

Material & Methods: Study was done at selected hospital in Etawah district with 50 samples, were selected by convenient sampling technique. descriptive comparative design was adopted. The standard tool (Framingham scale) and demographic tool was use to assess the risk of CAD among TYPE2 DM and Non DM patient .Research design is Non Experimental simple descriptive comparative study.

Result: Coronary artery disease and type2 DM has significant relationship with some of the demographical variables and selected health related outcome 52% patient of CAD with type 2 DM and 48% patient of CAD with other demographical variables. So prevalence of CAD is more common in type2 DM patient. Age, marital status and hypertension have significant relationship at the level of $p>0.05$. Body build, BMI, systolic BP and blood pressure have significant at the level of $p>0.05$.

Conclusion: This study has found coronary artery disease has significant relationship with type2 DM. This research also found in significant relationship between age, marital status and, body build, BMI and hypertension.

Keyword: Assess, prevalence, DM patient and Non Type 2DM.

Introduction

Coronary artery disease (CAD) is a major cause of death in the western countries and it is the becoming major cause of death in developing countries. This increase may be due to the rising prevalence of many risk factor such as diabetes mellitus, which is one of the most important of these risk factor .the prevalence of diabetes mellitus increasing globally and it has reached in the worldwide.

Type-2 diabetes mellitus is a strong risk factor of CAD and expert consider type-2 DM an equivalent to established CAD risk patient with have 2 to 4 times greater risk of developing CAD than non-diabetic patient.

Type 2 diabetes mellitus (T2DM) has emerged as a pandemic and is associated with increased cardiovascular morbidity and mortality. India, with >60

million

People with diabetes, has the second largest diabetic population of the world. The projection shows that this number will increase to 100 million by 2030 (>90% T2DM).¹ although diabetes rates are lower in rural than urban India, the

Overall diabetes burden is higher in rural India as 68% of India lives in "Villages". Nonetheless, the prevalence of diabetes in rural areas is increasing rapidly

Diabetic patient exhibit an increased risk for developing of atherosclerosis CAD for many reason including metabolic factor like hyperglycemia, dislipidemia and insulin resistance, which lead to endothelial cell vascular smooth muscle dysfunction, impaired platelet function and abnormal coagulation.

Inflammation play an important role in atherosclerosis .

Inflammation activation in type-2 DM result from obesity and insulin resistance in which in acute phase reaction occurs large no of inflammatory and non-inflammatory cytokine are released from adipose tissue endothelial dysfunction is generally present in diabetic patient with CAD and evidence by high level endothelin-1 and low level of nitric oxide.

PROBLEM STATEMENT

A comparative study to assess the prevalence rate of CAD among Type 2 DM patient and Non Type 2DM patient in selective hospitals Etawah district

OBJECTIVES:

1. To assess the prevalence rate of CAD among Type2 DM patient.
2. To assess the prevalence rate of CAD among non DM patient.
3. To compare the prevalence rate of CAD with Type 2 DM and non DM patient.
4. To identified the correlation between CAD and Type 2 DM with other modifiable and non-modifiable risk factor.

METHODOLOGY AND MATERIAL

Research approach:-The selection of research approach is the basic procedure for the research of enquires. The research approach helps the researcher to determine what data to collect and how to analyze it.

In this study the research approach is quantitative approach.

Research design:-A research design refers to the plan of organization of scientific investigation.

In this study Non Experimental simple descriptive comparative design was adopted.

POPULATION: The population of my study is the coronary artery disease people who are admitted in the selected hospital Etawah district.

SETTING: It is the physical setting where the study will be conducted .For my study the, setting will be selected hospitals of Etawah district is the physical location and condition in which data collection take place in a study.

SAMPLE: Sample in the study is CAD patient are admitted in selective hospital Etawah district.

SAPMLE TECHNIQUE: Convenient sampling technique will be used.

Study variable: coronary artery disease among type-2 DM patient and non-Type-2 patient.

Attributed variable: age, sex, obesity, life style, etc.
SAMPLE SIZE:50 Coronary artery disease patientsadmitted in selective hospital Etawah district.

TOOL AND TECHNIQUE:

TOOL	TECHNIQUE
Demographic questionnaires	Interview
PART-B Observation	Physiological measurement
PART- C Framingham scale	Interview

ANALYSIS AND INTERPRITATION OF DATA

Method of organizing, shorting and scrutinizing data in a way that a research question can be answered or meaningful inferences can be drawn.

SECTION: TABLE 1: Frequency and percentage distribution of socio-demographic variable.

Table no. 2 : Frequency and percentage distribution of health related outcomes

S. No.	Health related outcomes		Frequency (f)	Percentage (%)
1	Body build	Thin	6	12%
		Normal	16	32%
		Obese	28	56%
2	BMI	<18.5	6	12%
		18.5-24.9	16	32%
		>25	28	56%
3	Fatigue	Yes	38	76%
		No	12	24%
4	Blood pressure	Yes (HTN)	32	64%
		NO (HTN)	18	36%
5	Systolic BP	120-129	3	6%
		130-139	9	18%
		140-159	17	34%
		>160	21	42%
6	dyspnoea	Present	34	68%
		Absent	16	32%
7	Type 2 DM	Yes	26	52%
		No	24	48%
8	Total cholesterol level	160-199	8	16%
		200-239	21	42%
		240-279	10	20%
		>280	11	22%
9	HDL level	<40	8	16%
		41-50	23	46%
		51-60	14	28%
		>60	5	10%

Table no. 3 : Relation between Type 2 DM and CAD with selected demographic variables.

s. no.	Socio demographic variable	Chi-square Test (X ²)	(df)	P Value	S*/NS
1	Age	13.86	5	>0.05	S*
2	Gender	0.418551	1	<0.05	NS
3	Education	0.300228	3	<0.05	NS
4	Marital status	5.92332	1	>0.05	S*
5	Type of residence	0.06883	1	<0.05	NS
6	Type of family	0.155554	1	<0.05	NS
7	Religion	0.876914	1	<0.05	NS
8	Occupation	0.24227	2	<0.05	NS
9	Socio economic condition	0.210267	2	<0.05	NS
10	Type of work	0.3145	2	>0.05	NS
11	Life style	0.378073	2	<0.05	NS
12	Habit	0.019093	2	<0.05	NS
13	Hypertension	4.63	1	>0.05	S*

Table no 4 :socio demographic variables

s. no.	Socio demographic variable	Chi-square Test (X ²)	(df)	P Value	S*/NS
1	Body build	6.3057	2	5.99	S*
2	BMI	6.9602	2	5.99	S*
3	Fatigue	0.01024	1	3.84	NS
4	Blood pressure	4.1305	1	3.84	S*
5	Systolic BP	7.9243	3	7.82	S*
6	Dyspnoea	0.4231	1	3.84	NS
8	Total cholesterol	0.363117	3	7.82	NS
9	HDL level	0.483823	3	7.82	NS

8NS- Non significant, *S significant

RESULT

- Majority of 32% CAD patient were belonging to 51-60 years age groups 60% male and 40% female, 52% were belonging to illiterate and 14% high education qualification, 80% were married, 62% belonging from rural area, 64% patient were living in nuclear family and 36% were from joint family, 74% were Hindus and other 26% were Muslims, 48% were private job and 34% were agriculture 18% government job, majority of patient socio-

economic status is 50% 5000-10000 Rs/month , 66% people were belonging to sedentary work, 72% were having moderate stress, 60% not have any bad habit and 26% were smoking or 14% were smoking and alcohol intake, 64% were suffering from hypertension ,52% suffering from Type 2 DM and 48% were never diagnosis of CAD or 34% were diagnosis 1-5 year before, 61% were taking treatment of DM continue and 4% irregular or 35% were never taking treatment of Type 2 DM

- Majority 56% of body build belong to obese range, 56% patient were having BMI >25, 76% patient feel fatigue, 64% patient having hypertension, 42% patient having >160 systolic BP , 68% patient feel dyspnoea , 52 % patient having Type 2 DM , 42% patient having total cholesterol level 200-239 mg /dl , 46% patient having HDL level 41-50 mg/dl.
- Age, marital status, and hypertension have significant relationship with Type 2 diabetes mellitus and CAD at the level of P= 0.05 . Other selective demographic variable i.g. age, gender, education status, type of residence, type of family, religion, occupation, socio economic condition, type of work, life style, habit were found non significant at p= 0.05 level of significant with their degree of type 2 DM and Non DM in CAD patient.
- Body build, BMI, blood pressure, systolic BP, was significant at the level of p>0.05 and fatigue, dyspnoea total cholesterol, HDL level were found non significant at the level of p>0.05 with there relation of type 2 DM to CAD.

CONCLUSION

This study implies that there is significant relationship between type 2 diabetes mellitus with age, marital status and hypertension and other health related outcome in case of body

build, BMI, blood pressure and systolic BP.

Conflict of interest- Nil

Source of funding- Self

Ethical clearance/ permission-taken from institution advisory committee

REFERENCES-

- RC Turener, H Mlins, HAW Neil, IM Stratton, SE

- Manley, DR Matthews, RR Holman: risk factor of coronary artery disease in non –insulin dependent DM United kingdom prospective diabetics study (UKPS23) :A review article (1998) available from-www.bmj.com
2. Goran K Hansson: Inflammation, atherosclerosis, and coronary artery disease. *New England journal of medicine* 302, .A review article 2005 .Available from – www.nejm.org
 3. Lukkariinen, Hannele, Maija: treatment coronary artery disease improve quality of life in the long term.: a review article 2006 volume 55: available from www.journals.lww.com
 4. Mrs.Hepsi Rachel Charles , prevalence of cardiac risk factor among people attending an exhibition.; A review article of 2014 *IORS Journal of nursing and health science* .Available from www.iosrjournal.org
 5. Abdel-Ellah Al-Shudifat, Asgeir Johannessen, Mohammed Azab: Risk factors for coronary artery disease in patients undergoing elective coronary angiography in Jordan .A review article of 2017, Available from (<http://www.creativecommons.org/licenses/by/4.0/>),
 6. Jatinder Mokta¹, Kiran Mokta², Asha Ranjan³, Mehak Garg, Prevalence of Cardiovascular Risk Factors among Diabetic Population and Awareness of Diabetes among Diabetic Patients: A Population Based Himalayan Study ,A review of 2016,available from ,www.nursingjournal.com
 7. Sanne A. E. Peters¹ & Rachel R. Huxley² & Naveed Sattar³, Sex Differences in the Excess Risk of Cardiovascular Diseases Associated with Type 2 Diabetes: Potential Explanations and Clinical Implications, Published online: 26 May 2015# The Author(s) 2015. This article is published with open access at Springerlink.com
 8. Anjana RM, Pradeepa R, Deepa M, et al. ICMR–INDIAB Collaborative Study Group. Prevalence of diabetes and prediabetes (impaired fasting glucose and/or impaired glucose tolerance) in urban and rural India: phase I results of the Indian Council of Medical Research–India Diabetes (ICMRINDIAB) study. *Diabetologia* 2011; 54:3022–3027.
 9. Madhu Basavegowda, Kavitha Hanumanahally Shankarappa, Ashok Nagaralu Channabasappa, Srinath Kenkere Marulaiah, Magnitude and pattern of hypertension among diabetics; risk prediction for stroke and myocardial infarction, a review of 2014,available from, *Journal of Mahatma Gandhi Institute of Medical Sciences*
 10. Angelo Avogaro, Md Carlo Giorda, Md Marina Maggini, PhD Eduardo Mannucci, Md Roberto Raschetti, PhD Flavia Lombardo, PhD:- Incidence Of Coronary Heart Disease In Type 2 Diabetic Men And Women . A Review Of 2007 ,Available From *Diabetes Care*, Volume 30, Number 5, May 2007,([Http://Www.Creativecommons.Org/Licenses/By/4.0/](http://Www.Creativecommons.Org/Licenses/By/4.0/))