“A STUDY TO ASSESS THE KNOWLEDGE AND ATTITUDE REGARDING HEALTH HAZARDS OF JUNK/FAST FOOD CONSUMPTION AMONG ADOLESCENTS IN SELECTED HIGH SCHOOLS OF HUBBALLI”. ..

Lucy Gladies E¹, Sanjay Peerapur², Somashekarayya Kalmath³

¹Lecturer, Department of Medical Surgical Nursing, KLES Institute of Nursing Sciences, Hubballi, Karnataka, India.

²Principal & HOD Medical Surgical Nursing KLES Institute of Nursing Sciences, Hubballi Karnataka, India.

³Asso. Professor HOD Paediatric Nursing KLES Institute of Nursing Sciences, Hubballi Karnataka, India.

ABSTRACT:

Background: A long time ago, people ate foods that were very good for their health. They ate foods that did not contain chemicals, preservatives, artificial colors & artificial flavors. That's the reason they did not have any problem on their health & they lived longer than now. Unfortunately, today's world has been adapted to a system of consumption of foods which has several adverse effects on health.

Objectives: 1. To assess the knowledge and attitude regarding the health hazards of junk/fast foods consumption among adolescents. 2. To find out the correlation between the knowledge and attitude regarding the health hazards of junk/fast foods consumption among adolescents. 3. To find the association between level of knowledge and attitude of adolescents regarding the health hazards of junk/ fast foods consumption with their selected socio demographic variables

Methodology: A descriptive study was conducted among 100 adolescents of selected schools of Hubballi. Samples were selected using non probability; simple random sampling technique. Descriptive research design was used for the study. Data was collected by structured knowledge questionnaire and attitude scale. Data analysis was done using descriptive and inferential statistics.

Results: Overall results revealed that out of 100 samples 18 (17.8%) had good knowledge 62 (62.2%) had average knowledge and 20 (20%) had below average knowledge. Regarding attitude 11 (11.1%) had positive attitude 71 (71.1%) had favorable and 18 (17.8%) had negative attitude.

Conclusion: The study concluded that, adolescents having average knowledge and showed a favorable attitude regarding health hazards of junk food consumption so a proper education in regard to healthy habits of eating can reduce the risks of health hazards of junk food consumption.

Key words: Adolescents, Health hazards, Junk/Fast food.

INTRODUCTION

The world economy has become fast faced, and this has forced people to transform their life style in order to adapt to the changing pace. Family structures have evolved drastically with modern mothers having outside careers rather than being housewives, such mothers get out of work worn out with little strength to prepare a proper meal for the family¹.

Food is nutritious substance used to maintain life growth. Fast/Junk food is the term given that can be prepared and served very quickly which is sold in a restaurant or store with low quality preparation and packaged form for take away, ² It mainly includes chips, hot pies, wraps, rolls, burgers, shakes, pao bhaji,

panipuri, kebabs, pizzas, chicken, soups, Chinese food like noodles, Gobi Manchurian, French fries.etc ³

Fast food are rich in fat and calories, which effects the body causing gas, bloating, elevated heart rate and energy crashes. Continuous consumption of these can cause obesity, atherosclerosis, heart disease and diabetes.⁴

Nutritional balance is not maintained easily with junk food as there is no control over how they are cooked. Some are cooked with a lot of oil and butter and high content of fat, sodium and sugar ⁴, which can lead to reduced immunity, increased susceptibility to disease, impaired physical, mental development and reduced productivity. They also contain flavoring agents like

monosodium glutamate, caffeine, casein, excess sodium etc which may lead to many skin diseases vomiting, rashes, asthma and many other minor as well as major diseases⁵

In India during recent years, there has been an increasing trend towards the sale and consumption of street food. This phenomenon is more seen among adolescence³. A nutritional status of adolescents is highly influenced by various eating patterns, Particular areas of concern have included intake of more dietary fats in comparison to fruits and vegetables.

Globalization and urbanization in developing countries, where adolescent eating behavior is under spotlight.⁴ Adolescents are typically fond of eating 'junk food' not only for its taste, but because of peer group habits.³

A recent study has revealed that a full 48% of soda fountains at fast food restaurants contain coli form bacteria – which grows in feces. 11% contained E. coli. It's extremely sickening to see that even our soft drinks are now infected by food-borne illnesses that usually derive from meat.⁶

National Nutrient Survey (2009) has stated low nutrient is common among adolescent depending on their food choices included hunger or cravings, appeal of food, time and parents, and convenience of food.⁷

Nurse plays an important role in early identification of health problems, nutritional screening for adolescents and nutritional education. Through planned teaching programme, the nurse educates the adolescents regarding health hazards of junk food and helps to develop right attitude and practice healthy life styles. Health education measures have to be taken for changing the diet of adolescents who have fallen into the junk food habits.

Objectives of the study:

1. To assess the knowledge and attitude regarding the health hazards of junk/fast foods consumption among adolescents. .
2. To find out the correlation between the knowledge and attitude regarding the health hazards of junk/fast foods consumption among adolescents.
3. To find the association between pre test level of knowledge and attitude of adolescents regarding the health hazards of junk/ fast foods consumption with their selected socio demographic variables.

METHODOLOGY:

Research approach : Descriptive approach

Research Design : Descriptive Survey design.

Sampling technique& sample size: probability simple random sampling technique and 100

setting of the study : M.R.Shakare English Medium School. Hubballi

tools used :

1. Structured knowledge questionnaire to assess knowledge regarding health hazards of Junk food consumption among adolescents

Section I: Socio-demographic variables of subjects

Section II: Structured Knowledge Questionnaire which contains:

Part I: Items on general aspect of nutrition and junk /fast food

Part II: Items on health hazards of junk/fast food consumption

Part III: Items on prevention of health hazards of junk /fast food consumption

2. Attitude scale [modified 5 point likert scale]

Procedure of Data collection

The formal permission from the Principal of KLES' M.R.Sakhare English Medium School, Hubballi. The written consent was obtained from the participants. The pre-test was conducted by using structured knowledge questionnaire and attitude scale to assess the knowledge and attitude on health hazards of junk/fast food consumption. Data collected was then tabulated and analyzed.

Results:

Findings related to socio-demographic variables of subjects

Majority of the subjects 51(51%) belonged to the age group of 15years, 40(40%) were from 14years, 9(9%) were 13years old. In terms of gender the maximum number of subjects 87(87%) were males and 13(13%) were females. Majority of the subjects 36(36%) were from the class 10th, 33(33%) were from 8th std & 31(31%) were from 9th std. The maximum number of subjects 80(80%) were belonging to urban area and 20(20%) were from rural area. Majority of subjects 69(69%) were belonging to nuclear family and 31(31%) to joint family. Majority of the subjects 38(37.8%) family income were in the range of Rs5000-

10000, 38(37.8%) Rs 15000 and above, 22(22.2%) Rs 10000-15000, 2 (2.2%) were below Rs 5000 per month. The maximum of them 78(77.8%) consume junk food 1-2 times/week, minimum 11(11.1%) consume no junk food, 9(8.9%) consume 3-4 times/week, & very less that is 02(2.2%) consume 5-6 times /week. The majority of subjects 56(55.5%) received no pocket money, 22(22.2%) received in the range of Rs 0-100 per month, 15(15.6%) received Rs100-200 per month, & only 7(6.7%) received above Rs 200 of pocket money per month. Majority of subjects 47(46.7%) received information from television, 25(24.5%) through Mobile & Internet, 22(22.2%) through news papers, 04 (4.4%) received through health professionals & 02(2.2%) of them got information through Peer group and social circle

Section I: Analysis and Interpretation of knowledge scores of students regarding health hazards of junk food consumption.

Area of analysis	Mean	Median	Mode	Standard deviation	Range
Knowledge	12	13	14	2.5	16

Table no.1: reveals that the mean value of knowledge scores was 12, median 13, mode 14 standard deviation 2.5 and range was 16.

Table No 2: Mean, Median, Mode, Standard deviation and range of attitude score of subjects regarding health hazards of junk food consumption.
n = 100

Area of analysis	Mean	Median	Mode	Standard deviation	Range
Attitude	98	96	85	10	50

Table no.2: reveals that the mean value of attitude scores was 98, median 68, mode 96 standard deviation 10 and range was 50.

Table No 3: Frequency and percentage distribution of knowledge scores of subjects regarding health hazards of junk food consumption among adolescence

Knowledge score	Frequency (f)	Percentage (%)
Good (15.51 and above)	18	17.8
Average (11.09 - 15.51)	62	62.2
Poor (below 11.09)	20	20.0

Table 3 reveals 18 (17.8%) of subjects had good knowledge, 62 (62%) average knowledge and 20 (20%) had poor knowledge regarding health hazards of junk food consumption,

Graph 1: The bar graph represents the percentage distribution of level of knowledge scores of subjects regarding health hazards of junk food consumption
Table No 4: Frequency and percentage distribution of attitude scores of subjects regarding health hazards of junk food consumption among adolescence

Attitude score	Frequency (f)	Percentage (%)
Positive (70.08 and above)	11	11.1
Favorable (70.08 - 55)	71	71.1
Negative (below 55)	18	17.8

Table No 4 reveals that 11(11.1%) had positive attitude, 71(71.1%) had favourable attitude and 18(17.8%) negative attitude regarding health hazards of junk food consumption.

Graph2: The pyramids graph represents percentage distribution of level of attitude scores of subjects regarding health hazards of junk food consumption.

Table No 5: Analysis and interpretation of data to find out correlation between knowledge scores and attitude scores

X	Y	Karl pearson coefficient of correlation (r_{xy})
14	98	$r = 0.40$ ($0 < r_{xy} < 1$) Moderately positive co relation

Table No 5: reveals that $r_{xy} = 0.40$, ($0 < r_{xy} < 1$), there exists a moderately positive co relation between knowledge and attitude scores of students.

Section II: Analysis and interpretation of data to find out association between knowledge scores and attitude scores with selected demographic variables

Since the calculated chi- square value is more than tabulated there is an association between knowledge scores and few selected demographic variables that are in place of living, consumption of junk food, pocket money received per month, source of information.

Since the calculated chi- square value is more than tabulated value there is an association between attitude scores and few selected demographic variables that is in age and consumption of junk food per week.

CONCLUSION:

Based on the findings of the study, the following conclusions were drawn

1. The overall knowledge scores of adolescent were average.
2. The majority of adolescents had favourable attitude regarding junk food consumption.
3. There was a moderately positive co relation between

knowledge and attitude

4. There was an association between knowledge scores and few selected demographic variables that are in place of living, consumption of junk food, pocket money received per month, source of information in regard to knowledge scores.
5. There was an association between attitude scores and few selected demographic variables that is in age and consumption of junk food per week.

RECOMMENDATIONS:

1. A similar study that can be done on large and wider sample size and for a longer period of time would be more pertinent in making broad generalizations.
2. An experimental study can be conducted for assessing the knowledge on health hazards of junk food consumption.
3. A descriptive study can be conducted to assess knowledge, attitude and practices regarding health hazards of junk food consumption.
4. A similar study can be replicated in different setting.

REFERENCES:

1. Stevens B. The future of the family to 2030. Nutrition journal. [serial online]. Dec 2008 [cited on 2014 dec]; 26(2):10-16. Available from URL: <http://www.oecd.org/futures/42551944.pdf>.
2. Morris I. How fast food affects you negatively. Nutrition journal. [serial online]. 2010 sep 8 [cited on 2013 dec 12]; 14(4):20-22. Available From URL: <http://www.wonderlist.com/10worst food effects of fast food>.
3. Williams J. Mc Donald's fast food. [serial online]. January 2010 [cited on 2013 Dec 14]; 12(2)4-8. Available from URL:.
4. Rail K. Effects of fast food a survey in adults. [serial online]. 2000 march [cited on 2014 Dec 16]; 10(6)30-32 Available from URL: <http://www.myfit.ca>
5. Puri S, Bhatia V, Impact of a diet and nutrition related education package on the awareness and practices of school children of Chandigarh. The internet journal of epidemiology, 2008, volume 6, Pp.20-25.
6. Park M. Soda fountains contained fecal bacteria study found. [serial online]. 2010 janaury [cited on 2014 dec 20]; 64(4) 126-32. Available from URL: <http://edition.cnn.com/2010/HEALTH/01/08/soda.fountain.bacteria/>
7. Paediatrics Today, Vol 12 No.(2), Jan-Feb 2008, p. 14-15.