ISSN (O): 2582-693X

Original Article

PREVALENCE OF SMARTPHONE ADDICTION DURING COVID-19 PANDEMIC AMONG NURSING STUDENTS

MEHAK¹, PRIYANKA², RAJINDER KAUR³, RAKSHANDA⁴, SHAGUN SHARMA⁵,
SHUBHPREET⁶, SUNIL KUMAR GARG⁷, RAMANDEEP KAUR⁸

¹⁻⁶ Post Basic B.Sc. Nursing students at SINPMS, Badal, Sri Muktsar Sahib, Punjab.

⁷ Associate professor, M.Sc. (Paediatrics) and Research Guide, SINPMS, Badal, Sri Muktsar Sahib, Punjab.

⁸ Tutor/ Demonstrator, M.Sc. (Community health) and Research Co-Guide, SINPMS, Badal, Muktsar, Pb.

Corresponding E-mail: shagunsharma19091996@gmail.com

ABSTRACT

Introduction and aim- India is the second largest country with over 900 million mobile users in world. The evolution of mobile has shown its existence in increasing the range and maintaining the relation in public and communication but also badly affecting in form of giving out one personal information to friends, public or social groups in virtual touch. The statistic says more than 1 billion people in the world are using mobile phone. In addition internet excess has increased the more attachment of an individual with his smartphone. The study is planned to assess the prevalence of smartphone addiction among nursing students during COVID-19 pandemic. **Material and Methods-** A non-experimental cross sectional survey study was conducted on 420 nursing students studying in selected nursing colleges of Punjab. Data was collected through online mode. **Results-** The results revealed that 158 (37.6%) of nursing students were addicted to smartphone whereas 262 (62.4%) nursing students were not addicted to smart phone. **Conclusion-** There is a need to increase the awareness regarding use of smartphone's among nursing students in under to reduce the prevalence of smartphone addiction. **Keywords-** Prevalence, smartphone, addiction.

Received	Accepted	Available online
22/06/2021	28/06/2021	30/06/2021

INTRODUCTION

India is the second largest mobile hub with over 900 million mobile users in world. The mobile technology has invaded human life and nothing is left personal. In today's world smartphones are being over used not only for communication but also badly affecting in form of giving out our personal information to friends, public or social groups in virtual touch. Even an adolescent in a family consume mobiles and waste much of his time with mobiles than family or other important work. It was found that 22% of young children aging 6-9years, 60% in the age group of 10-14 years and 84% of teen aged 15-18 years have their own mobile phones.

As the Smartphone functions and applications are increasing endlessly, they are creating over dependence on these gadgets. Overuse of smartphone has also impacted badly in negative physical health and psychosocial impact among users. People are so involved with their phone that they frequently check their mobile for any missed communication without realizing their addiction. Such addiction leads to risk on mental health, bullying, eye strain, health

impact, lack of sleep and losing of self-control.

According to Worldwide scenario, the age group of 25-34 is found to have the highest smartphone usage rate of 62%. 53% of smartphone users are male and 47% are female. Indian teen's currently driving smartphone's market in India. The age groups of 16-18 years using smartphone's have shown a rapid rise from 5% in 2012, 25% in early 2014. Recently in 2013, there were around "51 million" smartphone users in India and rate rise from year 2012 was 90%.

The smartphone addiction magnitude in India ranged from 39% to 44% as per fixed effect calculated. Smartphone addiction among Indian teens can not only damage interpersonal skills, but also it can lead to significant negative health risks and harmful psychological effect on Indian adolescents.

In sudden COVID-19 outbreak, governments of many countries including India have imposed a lockdown to contain the spread of COVID-19. This lockdown involved closure of educational institutions and social establishments to limit

movement. With restriction on face to face contact or physical contact in the COVID-19 pandemic, smartphone are playing a key role in lockdown. Mostly students are spending more time on various screens at home which is seriously affecting their health and study reveals that the student are making excessive use of their smartphones during the lockdown.

The present study was taken up with an objective to assess the prevalence of smartphone addiction during COVID-19 pandemic among the nursing students of selected nursing colleges of Punjab.

MATERIAL AND METHOD

Research design: A quantitative research approach, Non-experimental survey design was used to assess the prevalence of smartphone addiction among nursing students studying in nursing colleges of Punjab.

Sample: The target population of the present study consists of nursing students of nursing colleges of Punjab. The proposed sample was selected by convenience sampling technique as the participants agreed to participate in the study. A total number of 420 nursing students responded

for the Smartphone addiction scale.

Research variable-

The research variable included in the study was prevalence of smartphone addiction among nursing students. **Research setting-** In early stage of research study it was planned to conduct the study in nursing colleges of Punjab. But due to COVID-19 pandemic and lock-down the data collected by online mode from participants.

Development and description of tool(s)- In order to achieve the research objectives, a structured questionnaire was prepared in form of google form to collect online data from participants to assess the prevalence of mobile phone addiction during COVID-19 pandemic. 420 nursing students studying in different nursing college of Punjab responded to the smartphone addiction scale. The tool included:- **Section A:** Questions related to Socio-demographical variables. **Section B:** Items related to Smartphone addiction (Smartphone addiction Scale)

Socio-demographical variables: This section consist of items for obtaining information about subjects such as class of participants, age (in years), gender, religion,

marital status, place of residence, nursing course in which studying, college, number of smartphone using, years of smartphone use & occupational status of parents.

Smartphone addiction scale (SAS):

Smartphone addiction scale (SAS) was used to assess the prevalence of smartphone addiction. In this scale 15 items related to addiction were used. This tool consist a five-point Likert scale (1.Strongly Disagree 2.Disagree 3. Neutral 4.Agree 5.strongly agree) with a maximum score of 75 and minimum score of 15.

Ethical considerations: Ethical approval to conduct the study was obtained from institutional research and ethical committee of SINPMS, Badal, Muktsar, Punjab. Permission to conduct the study through online mode was obtained from concerned authority. Informed consent was obtained from the participants through online mode to participate in the research project. **Data collection:** Data collection was done in the month of September, 2020 on 420 nursing students of selected college of Punjab. Prior to the data collection permission was obtained through online mode. Written consent was taken from participants after

assurance of confidentiality and anonymity.

The collected data were analyzed using descriptive and inferential statistics. Different charts and diagrams were used to depict the findings.

RESULTS:

Out of 420 participants (nursing students), the data revealed that according to their age, maximum 221 (52.6%) of the nursing students were more than 20 years of age whereas 199 (47.4%) of the nursing students were in the age group of 18-20 years.

Gender-wise distribution of subjects revealed that 387 (92.1%) nursing students were females and only 33 (7.9%) were males.

Religion-wise distribution revealed that 228 (54.3%) nursing students were Sikh, 162 (38.6%) were Hindu, 16 (3.8%) were Christian, 9 (2.1%) were Muslim and 5 (1.2%) belong to other category.

Distribution of nursing students according to marital status shows that maximum 397 (94.5%) were unmarried, 22 (5.2%) were married and only 1 belong to divorced and any other category.

Distribution regarding place of residence of

nursing students revealed that maximum 268 (63.8%) were residing out of hostel/on rent/ own house/ relative's house/ any other whereas, only 152 (36.2%) were residing in hostel.

Nursing course the nursing student were studying revealed that maximum 314(74.8%) were studying in B.Sc. Nursing, 50 (11.9%) were studying in GNM, 40 (9.5%) were in Post-Basic B.Sc. nursing and only 16 (3.8%) were studying in M.Sc. nursing or above.

The ownership status of nursing college the nursing students were studying revealed that maximum 306 (72.9%) were studying in Private/Charitable/Autonomous University/Any other nursing college whereas only 114(27.1%) were studying in a government nursing college or a college Constituent to Government University.

Regarding number of smartphones the nursing students were using, the data revealed that maximum 405 (96.4%) were using only 1 smartphone whereas 15 (3.6%) nursing students were using more than one smartphone.

Regarding the duration of smartphone use, the nursing students revealed that maximum

271 (64.5%) were using it from 1-5 years, 87 (20.7%) using from more than 5 years whereas only 62 (14.8%) were using smartphone from less than 1 year.

Table 1:- Mean smartphone addiction assessment score among nursing students.

N=420

Smartphone	N	Minimum	Maximum	Mean	Std. Deviation
Addiction Score	420	18.00	75.00	42.6881	9.31908

The above table depicts that mean smartphone addiction assessment score achieved by 420 nursing students was 42.6881 +/- 9.31908 (Mean+/-SD). The minimum score achieved was 18 and maximum was 75.

Table 2:- Prevalence of Smartphone addiction among Nursing students

N=420

Addiction status	Frequency	Percent
Addicted	158	37.6
Not Addicted	262	62.4
Total	420	100.0

The above table revealed the prevalence of smartphone addiction among nursing students. The results revealed that 158

(37.6%) of nursing students were addicted to smartphone whereas 262(62.4%) nursing students were not addicted to smartphone.

Table 3:- Association of prevalence of smartphone addiction among Nursing students with their demographic variables.

N=420

Demographic Variable		Addicted (f)	Non-addicted (f)	Degree of freedom (df)	Chi-square value
Place of residence	Hostel	94	58	1	0.0295*
	Out of hostel/On rent/ Own House/ Relative's House/ Any other	168	100		

S= Significant

p<.05 level of

significance

The above table reveals significant Chi-square value which means there is significance association between the smartphone addiction and place of residence of nursing student.

There was no significance association of addiction with other demographic variables.

DISCUSSION

In support of the research objective 1 the findings of the study revealed that

prevalence of smartphone addiction was found among 158 (37.6%) nursing students whereas 262 (62.4%) nursing students were not addicted to smartphone.

These finding were supported by **Ammarti R, Kakunjee A(2018)** cross sectional study conducted in a medical university in south India among 328 students to assess the smartphone addiction, The data was collected using a structural interview schedule revealed that 46% of students were using smartphone about 4-6 hours per day. The students reported maximum use of smartphone for the purpose of social networking sites. Out of 328 students 36.8% of students were addiction to smartphone. Nearly half of male and quarter of female students were found addiction to smartphone.

Another study conducted by **Sethuraman RA, Rao S, Vincent (2018)**, a cross sectional study to evaluate smartphone addiction among 192 medical students in the Andaman and Nicobar Island. A self administered SAS scale was used and assessed that out of 192 subjects who completed the questionnaire 85.40% (n=164) had smartphone addiction.

The data in support of objective 2 regarding association of smartphone addiction with selected their demographic variable revealed significance association of smartphone addiction with the place of residence (chi-square value= .029 S^*) of nursing students. These findings were supported by **Renuka K, Gopalakrishna (2019)**, a cross sectional study conducted to evaluate the prevalence of smartphone addiction in Kanchipuram District Tamil Nadu among 400 sample by calculated using the formula $4PQ/L2$. In this study 405 participants 191 participants were non smart phone users and 214 were smart phone users. The study revealed prevalence of smart phone addiction was 27.6%. Male respondents were more addicted than the female (OR-1.94, 95%CI: 1.12-3.77, $p=0.01$). There was a statistically significant association between subjects <45 years of age and smart phone addiction (OR-2.33, 95% CI: 1.31-4.13, $p=0.003$) compared to older age group. Likewise respondents who were class III (modified BG Prasad Scale) and above were more addicted compared to subjects below class III (OR-2.29, 95% CI: 1.32-3.98, $p=0.002$).

Abdon APV, Mesquite R carried out a cross sectional study to evaluate the prevalence of smartphone addiction and associated factor in Brazil among 274 adult. The adult were study for a socio demographical/ lifestyle questionnaire Physical Activity Questionnaire, and the Mobile Phone Addiction Test. The researchers found, in all 47.8% ($n=131$) of participants were mildly addicted and 44.9% ($n=123$) were moderately addicted. Addiction was associated with the amount of use($r=0.40$; $p=0.000$), age ($r=0.20$; $p=0.001$) and neck pain (OR=1.912; $p=0.008$). In the multivariate analysis, strong smartphone addiction was associated with amount of use (OR=1.766; $p=0.000$) and neck pain (OR=1.172; $p=0.037$). The prevalence of smartphone addiction was high in their sample and associated with the amount of smartphone use and neck pain.

Limitations and future perspective:

- ☐ ☐ The study is limited to nursing students of selected nursing colleges of Punjab.
- ☐ ☐ Small number of sample subjects limits the generalization of the study.

□□Difficulty found in face to face communication with research participants due to COVID-19 situation.

Recommendations:

Following recommendations were made-

- A similar study can be replicated on large number of subjects to generalize the findings.
- An experimental study can be conducted to assess the impact of modification strategies regarding smartphone use/addiction among nursing students
- A comparative study can be conducted to assess the prevalence of smart phone addiction in Punjab with other states.

REFERENCES:

1. Chen B, Liu F, Ding S, Ying X, Wang L, Wen Y. Gender differences in factors associated with smartphone addiction: a cross-sectional study among medical college students. *BMC Psychiatry*. 2017 Oct 10;17(1):341. doi: 10.1186/s12888-017-1503-z. PMID: 29017482; PMCID: PMC5634822.
2. Fernandes B, Biswas UN, Tan-Mansukhani R, Vallejo A, Essau CA. The impact of COVID-19 lockdown on internet use and escapism in adolescents. *Journal of Clinical Psychology with Children and Adolescent*. September 2020 - Vol. 7 no. 3 - pp 59 -65 doi: 10.21134 / rpcna.2020.mon.2056
3. Davey S, Davey A. Assessment of Smartphone Addiction in Indian Adolescents: A Mixed Method Study by Systematic-review and Meta-analysis Approach. *Int J Prev Med*. 2014;5(12):1500-1511.