JOURNAL

OF LIVING ARCHITECTURE

A GREEN ROOFS FOR HEALTHY CITIES PUBLICATION

Volume 8 Number 1 Pages 19-40

2021

Observations on the survival capacity of 118 plant taxa on a green roof in a semi-arid climate: 12 year update

Amy Schneider¹, Michelle Landis², Jennifer Boussetot^{2,*}

¹Denver Botanic Gardens, 1007 York Street, Denver, CO 80206,

²Department of Horticulture and Landscape Architecture, Colorado State University, Ft. Collins, CO 80523

*corresponding author: Jennifer.Boussetot@colostate.edu

ABSTRACT

The Green Roof at Denver Botanic Gardens was built in the fall of 2007. Since installation, data on the green roof have been collected on 118 plant taxa 43 of which have survived for over 10 years. Plants were grouped based on their metabolic and growth type: succulents, creeping forbs, upright forbs, graminoid, and shrubs. Overall, shrubs and succulents have displayed the highest survival rates, showing the greatest potential for use on green roofs in semi-arid conditions. Graminoids survived about half the time and creeping and upright forbs had the lowest overall survival in semi-arid Colorado. Species survivability rates were calculated based on the number of plants of that species originally installed on the green roof.

Key words: Green Roof, Long-Term Species Evaluations, Semi-Arid

INTRODUCTION

Green roofs are constructed ecosystems installed on rooftops in cities to help mitigate the negative effects of urbanization and add space for biodiversity. Green roof environmental benefits include: urban heat island mitigation, stormwater runoff management, utilization as urban agriculture sites, restoration of urban biodiversity, building energy savings, and aesthetics (Getter and Rowe, 2006; Oberndorfer et al., 2007). Several US cities such as Denver, CO,

This peer-reviewed article is provided free and open-access.

Philadelphia, PA, Seattle, WA, New York, NY, Chicago, IL, (and more) recognize these benefits and have developed incentives to increase green roof installations.

Green roofs that contain a diversity of plant species and plant forms have been shown to outperform monocultures, providing greater ecosystem services (Lundholm & Cadotte, 2015). Diverse plant communities with many plant forms, such as shrubs, grasses, and herbaceous perennials increase the total cooling effect that the green roof system can offer (Cao et al., 2019), lead to greater stormwater detention (Lundholm et al., 2010), and result in an increase in arthropod richness on the rooftop (Braaker et al., 2017).

Plant species diversity also benefits individual plants on a green roof. A range of plant forms that grow at different heights, causing shading, can influence overall plant success on a green roof (Aguiar et al., 2019). Plant species selection based on substrate depth and moisture retention of the substrate results in optimal survival rates (Graceson et al., 2014; Nektarios et al., 2011). Long-term diverse plant taxa evaluations on green roofs in North America have seldomly been published. Only one other long-term study in North America by Lubell et al. (2017), is found in literature.

The harsh environment on green roofs can cause many challenges for plants because of higher intensity of light, higher and lower air and substrate temperatures, relatively shallow substrate depth, wind gusts, low water availability, and more intense storm effects. Therefore, taxa planted must be selected for both survivability and enhanced benefits (Shafique et. al, 2018, Reyes et al., 2016). Sedum spp. are commonly used on green roofs because of their ability to efficiently utilize moisture and survive in dry locations. Green roofs that contain a diverse range of plant taxa have been shown to outperform monocultures, providing greater ecosystem services (Lundholm and Cadotte, 2015).

Plant provenance could play a role on plant survivability on green roofs, but more research must be conducted, especially in North America (Dvorak and Volder 2010). A ten-year study of native plant survival conducted in the Northeastern U.S. by Lubell et al (2017) found that approximately 50% of the initial plant taxa perished. In a three-year evaluation of native plants conducted in Michigan, 67% of native plant taxa had perished when no supplemental irrigation was added after establishment (Monterusso et al., 2005).

To date, 118 plant taxa have been trailed on the semi-arid Denver Botanic Gardens (DBG) green roof over the past 12 years. The DBG green roof, along the Front Range of Colorado, is used to demonstrate potential plant taxa for use on green roofs with similar environments and limited moisture. Choosing plant taxa suited to the climate, and especially the precipitation of an area, can change the overall air temperature cooling effects, reducing or enhancing the success of a green roof (Chagolla-Aranda et al., 2017). Plant taxa in this study were characterized by plant metabolic pathway and growth form, as in evaluations conducted by MacIvor and Lundholm (2011). The objective was to evaluate plant taxa in this observational study for long-term success as potential green roof taxa used in a semi-arid climate. Additionally, we discuss how plant taxa provenance may have influenced survivability.

MATERIALS AND METHODS

Located along the Front Range of Denver, Colorado, USA in USDA plant hardiness zone 5 at an elevation of 1609 meters above sea level, the DBG Green Roof was installed on the south side of a one-story building, the Boettcher Memorial Tropical Conservatory. Established in the fall of 2007, this is one of Denver's earliest green roof installations (Figure 1).

The green roof is 195 m² with substrate depth gradually increasing from 10 cm at the northwest edge to 45.7 cm near the southeast corner, which creates a small berm. This green roof is considered semi-intensive, which allows us to evaluate a range of plant taxa suited to variable substrate depths. We did not control for substrate depth or microclimates on this green roof as this is an observational study of survivability only. The substrate consists of 80% expanded shale and 20% compost/composted bark, by volume. During the initial planting in 2007, mycorrhizae were added to the substrate, and in 2009 local compost was applied at the rate of approximately 6 mm on the surface. Since then, no supplemental nutrients have been applied.

Figure 1. The Denver Botanic Gardens (DBG) green roof in September 2013. Photo by Amy Schneider.

Environmental Conditions

Microclimates are created by shrubs and the three-foot-high concrete parapet walls that surround the east, south and western edges of the space. In the winter, the roof heats up during the day due to reflection from the surrounding surfaces and drops to freezing temperatures

overnight. The extreme seasonal fluctuations on the green roof, low moisture holding capacity of the substrate, and shallow depth of the substrate make survival for plants difficult.

The Colorado Front Range is a temperate and semi-arid region averaging from 18°C in warmer months to 2°C in the cooler months with an average annual precipitation rate of about 350 mm. Precipitation at DBG was collected at the southwest corner of the gardens (Table 1). Due to low precipitation and extreme summertime heat, irrigation was installed on the green roof at planting time. Half of the green roof was irrigated by an underground drip and the other half by popup sprinkler heads. For initial plant establishment, between 2007-2009, approximately 100-150 mm of water through irrigation was provided per month throughout the growing season with some supplemental hand watering. From 2011 on, the irrigation system is always activated in July with an annual irrigation rate of 200 mm or less. In 2009, 2011, 2014, and 2015, annual precipitation at DBG was exceptionally high, over 500 mm. A total of 1020 mm of supplemental irrigation has been applied to the green roof between the years of 2010-2019.

Table 1. Total monthly precipitation (in mm) and first/last snow at the Denver Botanic Gardens (DBG) from 2007 to 2019 (CoCoRaHS, n.d; accessed January 2020).

Month	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Jan	4.3	11.4	12.2	4.1	13.7	11.7	7.4	27.2	10.7	8.9	22.6	10.7	45.5
Feb	0.8	14.7	3.8	19.1	27.2	50.5	26.4	8.4	60.5	23.6	4.1	10.4	23.4
Mar	33.3	20.6	33.3	47.5	8.6	Trace	40.9	27.7	22.1	63.5	21.3	50.5	52.3
Apr	70.1	29.5	114.0	56.1	22.1	31.8	69.6	38.9	51.6	10.4	61.7	32.8	21.3
May	97.5	57.2	61.0	55.9	111.8	37.1	47.0	102.6	161.8	72.1	61.2	67.6	99.6
Jun	12.7	17.0	85.1	52.3	43.4	10.4	13.5	30.5	219.2	49.3	28.7	28.2	30.7
Jul	26.2	5.8	91.2	30.5	168.9	57.9	70.9	118.6	40.6	17.8	29.7	59.7	44.7
Aug	55.6	108.7	25.9	11.2	17.8	9.9	42.7	38.4	23.6	39.4	22.6	24.6	32.3
Sep	18.0	37.6	21.3	3.0	36.1	68.1	139.2	52.6	7.1	1.3	38.6	43.9	20.0
Oct	30.2	18.3	67.8	14.2	47.2	31.0	17.5	27.9	54.4	2.5	32.0	29.2	7.4
Nov	5.6	5.6	26.9	15.0	19.6	9.1	4.8	16.8	59.4	17.8	6.6	9.1	
Dec	42.4	13.2	14.5	7.4	43.4	5.8	10.2	18.5	27.9	20.1	3.0	2.5	
Total	396.7	339.6	557.0	316.2	559.8	323.3	490.0	508.0	738.9	326.6	332.2	369.3	377.2
Last Snow	4/14	5/14	4/25	5/12	4/23	4/3	5/2	5/12	5/10	4/29	5/18	4/24	5/21
First Snow	10/22	11/13	10/10	11/10	10/26	10/5	10/4	11/11	11/5	11/17	10/9	10/31	10/10

Plants

During the past 12 growing seasons, from 2007-2019, 118 plant taxa have been trialed on the green roof. Many of the taxa were obtained from onsite propagation at DBG, grown from seed or cuttings, and a few presented spontaneously. Most of the 118 plant taxa trialed are generally available in the local green industry.

Planting dates for the plant taxa varied between 2007-2019 (Table 2, 3, and 4). Plants were installed on the green roof for aesthetic design purposes and were therefore massed. As plants perished they were either replanted or replaced by a different plant taxa. In 2007, during the installation, 53 plant taxa were planted at the beginning of the trial, ten new taxa were planted in 2008, five planted in 2009, 16 more were added in 2010, 20 in 2011, seven in 2012, two in

2014, one in 2016, and 4 new plant taxa were planted in 2019 (including the spontaneous *Muhlenbergia* sp.). No new plants were planted in 2013, 2015, 2017, or 2018.

For this evaluation, plants were categorized by plant growth form characteristics, as in a study completed in Halifax, Canada (MacIvor and Lundholm, 2011). The plant groups represented in this study include: succulents, creeping forbs (which in this evaluation only represents non-succulents), upright forbs (represents all non-carpet growing forbs), graminoids, and woody shrubs. The creeping forbs were also subcategorized into bun forming forbs because they do not spread as rapidly as the other creeping forbs. All plant taxa were monitored for 12 years and designated as ‘perish’ or ‘survive’. The ‘survive’ category was then subdivided into 2 groups: plant taxa surviving over 10 years and plant taxa currently surviving under 10 years.

Plant taxa provenance information was included to determine if provenance could help explain survival rate in each plant category. In the succulent category, 15 of 22 planted taxa are native to North America, in the creeping forbs category, 15 of the total 47 are native, 13 of the 24 upright forbs, all 8 of the graminoids (excluding *Muhlenbergia* sp. and *Andropogon* sp. found in 2019), and of the woody shrubs, 12 of the 15 are native to North America.

Table 2. All taxa surviving more than 10 years by scientific name, common name, plant growth form, months surviving (two dates means replanting), and provenance.

Scientific Name	Common Name	Form	Months since planting (year planted)	Provenance (Kew science, n.d.)
<i>Acantholimon armenum</i>	acantholimon, spike thrift	creeping forb, bun	137 (2008)	Asia minor
<i>Acantholimon litvinovii</i>	acantholimon, spike thrift	creeping forb, bun	137 (2008)	Central Asia (Kirgizstan, Uzbekistan)
<i>Agave parryi</i>	Parry’s agave	succulent	137 (2008)	S.W. United States, N. Mexico
<i>Amorpha fruticosa</i> ‘Nana’	dwarf false indigo	woody shrub	125 (2009)	S. United States*
<i>Arctostaphylos</i> ‘Lauren’s Best’	Lauren’s best manzanita	woody shrub	143 (2007)	W. North America*
<i>Arctostaphylos patula</i>	greenleaf manzanita	woody shrub	143 (2007)	W. United States, N.W. Mexico
<i>Arctostaphylos</i> x <i>coloradoensis</i> ‘Cascade’	cascade manzanita	woody shrub	143 (2007)	W. United States*
<i>Cercocarpus breviflorus</i>	hairy mountain mahogany	woody shrub	143 (2007)	S.W. United States (AZ, NM, TX), N. Mexico
<i>Chamaebatiaria millefolium</i>	desert sweet	woody shrub	143 (2007)	W. United States
<i>Chilopsis linearis</i>	desert willow	woody shrub	143 (2007)	S.W. United States (incl. Georgia)
<i>Dianthus anatolicus</i>	Anatolian pink	creeping forb, bun	137 (2008)	Asia minor
<i>Draba hispanica</i>	Spanish draba	creeping forb	143 (2007)	N.W. Africa, S.W. Europe (Spain)
<i>Echinocereus fendleri</i>	Fendler’s hedgehog cactus	succulent	125 (2009)	S.W. United States

<i>Echinocereus triglochidiatus</i>	kingcup cactus	succulent	143 (2007)	S.W. United States (AZ, NM, CO)
<i>Echinocereus triglochidiatus</i> white sands strain	White sands kingcup cactus	succulent	143 (2007)	S.W United States
<i>Echinocereus viridiflorus</i>	nylon hedgehog cactus	succulent	143 (2007)	C. United States, NE. Mexico
<i>Echium amoenum</i>	red feathers	upright forb	143 (2007)	Middle East (Iran, Transcaucasus)
<i>Ericameria nauseosa</i> subsp. <i>nauseosa</i> var. <i>nauseosa</i>	rubber rabbitbrush	woody shrub	143 (2007)	C. United States
<i>Eriogonum umbellatum</i> var. <i>aureum</i>	sulphur flower buckwheat	creeping forb	143 (2007), 137 (2008)	W., N.W. United States (excl. WA, CA)
<i>Escobaria missouriensis</i>	Missouri foxtail cactus	succulent	143 (2007)	C., M.W. United States
<i>Escobaria vivipara</i>	spiny star	succulent	143 (2007)	C., S.W. North America
<i>Euonymus nanus</i> var. <i>turkestanicus</i>	Turkistan burning bush	woody shrub	143 (2007)	C. Asia (Altai Mountains)
<i>Hesperaloe parviflora</i>	red yucca, hummingbird yucca	succulent	143 (2007), 137 (2008)	S. United States (TX), N.E. Mexico
<i>Heterotheca jonesii</i>	Jones' false goldenaster	creeping forb	143 (2007)	W. United States (Utah)
<i>Iris germanica</i> dwarf	dwarf Germanic iris	upright forb	143 (2007)	S.E. Europe (Yugoslavia)
<i>Maihuenia poeppigii</i>	chupasangre	succulent	137 (2008)	S. South America (Argentina, Chile)
<i>Nolina microcarpa</i>	beargrass	succulent	143 (2007)	S.W. United States (AZ, NM), Mexico
<i>Opuntia aurea</i>	golden pricklypear	succulent	143 (2007)	W., S.W. United States (AZ, UT)
<i>Opuntia phaeacantha</i>	tulip pricklypear	succulent	143 (2007)	W., S.W. United States, N. Mexico
<i>Opuntia polyacantha</i>	plains pricklypear	succulent	143 (2007)	C., M.W. United States, Canada
<i>Pediocactus simpsonii</i>	mountain ball cactus	succulent	143 (2007)	Central Western United States
<i>Penstemon pinifolius</i>	pineleaf penstemon	creeping forb	125 (2009)	S.W. United States (AZ, NM), Mexico
<i>Phedimus kamschaticus</i> / <i>Sedum kamschaticum</i>	Russian stone crop	succulent	143 (2007)	E., N.E. Asia
<i>Phemeranthus calycinus</i>	largeflower flameflower	succulent	137 (2008), 101 (2011)	C., M.W., S. United States, Mexico
<i>Phlox bifida</i> 'Betty Blake'	cleft phlox	creeping forb	143 (2007)	C. E. United States and VT*
<i>Physaria bellii</i>	front range twinpod	creeping forb	143 (2007)	M.W. United States, CO
<i>Salvia greggii</i>	autumn sage	woody shrub	137 (2008)	S. United States (TX), NE. Mexico
<i>Salvia pachyphylla</i>	mountain desert sage	upright forb	143 (2007)	S.W. United States, Mexico
<i>Sedum lanceolatum</i>	spearleaf stonecrop	succulent	143 (2007)	W., N.W. North America
<i>Thelesperma ambiguum</i>	Colorado greenthread	upright forb	143 (2007)	S.W. United States (TX and NM), N.E. Mexico
<i>Veronica liwanensis</i>	Turkish speedwell	creeping forb	143 (2007)	Asia minor

				(Transcaucasus, Turkey)
<i>Veronica pectinata</i>	woolly speedwell	creeping forb	143 (2007)	Asia minor (Bulgaria, Turkey, Turkey-in-Europe)
<i>Veronica thymoides</i>	thyme-leaf speedwell	creeping forb	143 (2007)	Asia minor (Turkey)

Note: * includes plants that are cultivars

Table 3. All taxa surviving less than 10 years by scientific name, common name, plant growth form, months surviving, and provenance.

Scientific Name	Common Name	Form	Months since planting (year planted)	Provenance (Kew science, n.d.)
<i>Andropogon</i> spp.	big bluestem	graminoid	n/a (2019)	n/a
<i>Andropogon ternarius</i>	splitbeard bluestem	graminoid	84 (2012)	S., M.W., S.E. United States, N.E. Mexico
<i>Andropogon virginicus</i>	broomsedge bluestem	graminoid	84 (2012)	E. North America (incl. CA), N. South America
<i>Atriplex confertifolia</i>	shadescale sagebrush	woody shrub	113 (2010)	W. United States N.E. Mexico
<i>Bouteloua gracilis</i> 'Blonde Ambition' (spontaneous)	blue grama grass	graminoid	n/a (2019)	C., M.W., S.W. United States, Canada, Mexico*
<i>Chrysanthemum weyrichii</i>	dwarf chrysanthemum	creeping forb	101 (2011)	C. Europe, C., N. Asia
<i>Ephedra minuta</i>	miniature joint fir	woody shrub	113 (2010)	S., C. China (China South-Central, Qinghai)
<i>Erigeron vetensis</i>	early bluetop fleabane	creeping forb	113 (2010)	M.W. United States
<i>Fallugia paradoxa</i>	apache plume	woody shrub	60 (2014)	S.W. United States, N. Mexico
<i>Genista tinctoria</i> 'Royal Gold'	Dyer's greenweed, Dyer's bloom	creeping forb	60 (2014)	Europe, Middle East
<i>Genista lydia</i>	lydia broom	creeping forb	36 (2016)	S.E. Europe to Syria
<i>Ipomopsis aggregata</i>	scarlet gilia	upright forb	113 (2010)	S. W. United States (CO, NM), N. Mexico
<i>Muhlenbergia montana</i>	mountain muhly	graminoid	84 (2012)	M.W. United States (incl. CA), N., C. Mexico
<i>Muhlenbergia</i> spp. (spontaneous)	muhly	graminoid	n/a (2019)	n/a
<i>Silene schafta</i>	autumn catchfly	creeping forb	113 (2010)	Middle East (Iran, Transcaucasia)
<i>Sporobolus heterolepis</i>	prairie dropseed	graminoid	n/a (2019)	M.W., N., E. United States
<i>Teucrium graphaloides</i> (2)	germander	creeping forb	n/a (2019)	S.W. Europe (Spain, Portugal), Middle East

Note: * includes plants that are cultivars

Table 4. All taxa that 'perished' by scientific name, common name, plant growth form, months surviving (two dates means replanting), and provenance.

			Months alive since Planting	
--	--	--	-----------------------------	--

Scientific Name	Common Name	Form	(year planted)	Provenance (Kew science, n.d.)
<i>Acantholimon acerosum</i>	acantholimon, spike thrift	creeping forb, bun	72 (2008)	Asia minor (Lebanon-Syria, Turkey, Transcaucasia), Middle East
<i>Aethionema schistosum</i>	fragrant Persian stonecress	creeping forb	60 (2007)	Asia minor (Turkey)
<i>Anacyclus maroccanus</i>	Moroccan chamomile	creeping forb	<12 (2011)	N.W. Africa (Morocco)
<i>Antennaria parvifolia</i> ‘McClintock’	dwarf pussytoes	creeping forb	60 (2007)	C. W. United States, Canada, N. Mexico*
<i>Arenaria alfacarensis</i>	Spanish sandwort	creeping forb	n/a (2007)	S.W. Europe (Spain)
<i>Artemisia</i> sp.	wormwood	creeping forb	n/a (2007)	Middle East
<i>Baileya multiradiata</i>	desert marigold	upright forb	n/a (2009)	S.W. United States, N. Mexico
<i>Braya alpine</i>	smooth rockcress	upright forb	<12 (2010)	Central Europe (Austria, Italy)
<i>Campanula incurva</i>	Evia bellflower	upright forb	24 (2011)	S. Europe (Greece, E. Aegean Is.)
<i>Campanula kemulariae</i>	bellflower, blue bells	upright forb	<12 (2010)	Asia minor (Transcaucasia)
<i>Clematis columbiana</i> var. <i>tenuiloba</i>	rock clematis	creeping forb	<12 (2011)	M.W., N.W. United States
<i>Cotula hispida</i>	silver cotula	creeping forb	72 (2007) 60 (2010)	S. Africa
<i>Cotyledon orbiculata</i>	pigs ear	succulent	<12 (2012)	S.W. Africa
<i>Cytisus purgans</i> ‘Spanish Gold’	Spanish gold broom	woody shrub	96 (2007)	W. Europe (Spain, France)*
<i>Delosperma cooperi</i>	hardy ice plant	succulent	96 (2007)	S. Africa (Free State, Lesotho)
<i>Delosperma nubigenum</i>	yellow ice plant	succulent	96 (2007)	South Africa (Free State, Lesotho)
<i>Delosperma</i> ‘Kelaidis’	Mesa Verde ice plant	succulent	72 (2007), 84 (2008)	S. Africa*
<i>Draba streptocarpa</i>	pretty draba	creeping forb, bun	n/a (2007)	M.W. United States (CO, NM, WY)
<i>Dryas octopetala</i> var. <i>hookeriana</i>	Hooker’s mountain-avens	creeping forb	<12 (2010)	N.W., M.W. United States, North America
<i>Erigeron elatior</i>	tall fleabane	upright forb	89 (2011)	M.W. United States
<i>Erigeron leiomerus</i>	rockslide yellow fleabane	creeping forb	n/a (2011)	M.W. United States
<i>Erysimum capitatum</i> var. <i>purshii</i>	alpine wallflower	upright forb	24 (2011)	C. W. United States N.W. North America
<i>Euphorbia polychroma</i> ‘Candy’	yellow cushion spurge	upright forb	<12 (2008)	C., S.E. Europe, Asia Minor*
<i>Fendlera rupicola</i> var. <i>wrightii</i>	Wright’s fendlerbush	woody shrub	n/a (2007)	S.W. United States, N. Mexico
<i>Festuca brachyphylla</i>	alpine fescue	graminoid	<12 (2012)	W., N. North America,, N. Asia, N. Europe (Sweden)
<i>Hedeoma</i> clone	pennyroyal	upright forb	<12 (2011)	S. United States (NM)
<i>Herniaria glabra</i>	green carpet	creeping forb	n/a (2007)	Europe, W. Asia, Middle East, N. Africa, S. China
<i>Herniaria glabra</i> ‘Sea	Sea foam green carpet	creeping forb	n/a (2008)	Europe, W. Asia, Middle

Foam'				East, N. Africa, S. China*
<i>Heteropappus</i> sp.	Heteropappus	creeping forb	<12 (2010)	Eurasia
<i>Hypericum olympicum</i>	Mount Olympus St. John's wort	upright forb	48 (2011)	S. Europe, Asia Minor (Turkey)
<i>Ipomopsis rubra</i>	standing cypress, scarlet gilia	upright forb	n/a (2008)	S. United States
<i>Leucanthemum atlanticum</i>	daisy (collected in Morocco)	creeping forb	<12 (2011)	S., C. Europe
<i>Muhlenbergia emersleyi</i>	bullgrass	graminoid	36 (2012)	S.W. United States, Mexico
<i>Olsynium biflorum</i>	grass widow	upright forb	<12 (2011)	S. South America (S. Argentina)
<i>Panicum virgatum</i>	switchgrass	graminoid	96 (2007)	C., E. North America, Mexico
<i>Paronychia kapela</i> subsp. <i>serpyllifolia</i>	nailwort	creeping forb, bun	n/a (2007)	S. Europe
<i>Penstemon angustifolius</i>	broadbeard beardtongue	upright forb	36 (2010)	M.W. United States
<i>Penstemon caespitosus</i>	mat penstemon	creeping forb	72 (2007)	S. United States (WY, CO, UT)
<i>Penstemon cyananthus</i>	Wasatch beardtongue	upright forb	24 (2010)	N.W. United States (WY, MT, ID, UT)
<i>Penstemon fendleri</i>	Fendler's penstemon	upright forb	<12 (2007)	S. United States, N.E. Mexico
<i>Penstemon grahamii</i>	Graham beardtongue	creeping forb	<12 (2010)	W. United States (CO, UT)
<i>Penstemon linarioides</i> subsp. <i>coloradensis</i>	Colorado narrowleaf beardtongue	creeping forb	60 (2007)	S.W. United States
<i>Petrophytum caespitosum</i>	mat rockspirea	creeping forb	<12 (2010)	W. United States. N.E. Mexico
<i>Phacelia campanularia</i> (seed)	California bluebells	upright forb	<12 (2009)	W. United States (CA)
<i>Poa fendleriana</i>	muttongrass	graminoid	36 (2012)	W. United States, N. Mexico, W. Canada
<i>Potentilla hyparctica</i>	arctic cinquefoil	creeping forb	n/a (2007)	N. Asia, N. Europe (Sweden), N.W., N.E. North America
<i>Potentilla nepalensis</i> 'Shogran'	Nepal cinquefoil	creeping forb	<12 (2010)	E. Asia, W. Himalayas*
<i>Putoria calabrica</i>	stinking madder	upright forb	<12 (2011)	S. Europe, Asia minor, N. Africa
<i>Salvia cryptantha</i>	salvia	upright forb	<12 (2011)	Asia minor (Turkey)
<i>Salvia</i> 'Eskihar'	Eskihar sage	creeping forb	<12 (2011)	Asia minor*
<i>Scutellaria prostrata</i>	prostrate skullcap	creeping forb	<12 (2011)	Himalayas
<i>Scutellaria scordifolia</i>	skullcap hyssop	creeping forb	<12 (2010)	S. Mexico
<i>Sedum rupestre</i>	blue stonecrop	succulent	48 (2011)	Europe, Asia minor
<i>Streptanthus cordatus</i>	heartleaf twistflower	creeping forb	36 (2010)	W. United States
<i>Teucrium aroanium</i>	gray creeping germander	creeping forb	53 (2010)	S. Europe (Greece)
<i>Teucrium graphaloides</i>	germander	creeping forb	65 (2011)	S.W. Europe (Spain, Portugal), Middle East
<i>Thymus neiceffi</i>	juniper leaf thyme	creeping forb	108 (2007)	Europe, N. Africa, Asia

<i>Townsendia eximia</i>	Townsend's daisy	upright forb	n/a (2007)	S.W. United States (CO, N.M)
<i>Vella spinosa</i>	fine broom, spiny broom	creeping forb	<12 (2011)	S. Europe (Spain)
<i>Ziziphora bungeana</i>	Ziziphora	upright forb	<12 (2011)	Middle East, C. Asia

Note: * includes plants that are cultivars

Data Collection and Analysis

Each year in early autumn, data were collected on each of the plant taxa on the green roof. Plants were observed and recorded in the categories of survive or perish, and when possible, the number of remaining plants were recorded. Additionally, in April 2020 data was collected on the number of individual plants that have overwintered for each taxa. This was then used to determine the calendar months of survival and the relative percent survival rate for each taxa since planting. Additionally, data on plant provenance was noted to help interpret success rate of these taxa on a green roof in a semi-arid climate.

RESULTS AND DISCUSSION

Succulents

The succulent category had the overall highest survival rate in this evaluation (Figure 2). This is not surprising as succulents have proven to be successful on green roofs around the globe (Erwin et al., 2019; Nektarios et al., 2015; Vahdati et al., 2017) with some exceptions (Nektarios et al., 2020). From 2007-2009, 19 succulent taxa were planted (*Hesperaloe parviflora* and *Delosperma* 'Kelaidis' planted twice). Sixteen out of 19 taxa, or 84%, have survived. Later, between 2010-2012, three succulent taxa were planted, and of them, one, or 33%, survived. In total, 22 succulent taxa have been trialed on the rooftop, of them, 17, or 77%, have survived.

Many of the succulents on the DBG green roof have been successful and grown nearly exponentially, including: *Agave parryi*, *Nolina microcarpa*, *Opuntia aurea*, *Opuntia phaeacantha*, *Opuntia polyacantha*, *Phedimus kamschaticus*, *Phemeranthus calycinus*, and *S. lanceolatum*. Four of the total six succulents that perished did so in 2015, all likely due to a sudden November 2014 freeze, including *S. rupestre*, *D. cooperi*, *D. nubigenum*, and one of the two *D.* 'Kelaidis'. This suggests that the genus *Delosperma* may not be suitable for our low-irrigation, semi-arid, temperate green roof; all the species trialed in the genus perished in this evaluation. Remarkably, others have found the genus *Delosperma* to be resilient and ideal for use in water limited green roofs (Bousselot et al., 2010; Bousselot et al., 2011; Klein and Coffman 2015; Thuring et al., 2010).

Although less evaluated on green roofs in general, the monocarpic *A. parryi* has shown incredible vigor. Starting from four plants in 2008, with two blooming in 2011, it has resulted in 19 plants between both reseeding and runners, increasing in individuals by 475%. *Maihuea poeppigii* has survived, although it was observed as perished in 2013 and 2014; this likely due to early fall dormancy or its branched, dry aesthetic. *Cotyledon orbiculata*, which was mistakenly installed by a helper on the green roof, quickly perished the year it was planted,

2012, as soon as it became cold. *C. orbiculata* is only hardy to zone 9b-12, making it too tender for survival outdoors in Colorado in USDA hardiness zone 5.

Interestingly, of all 22 succulent taxa that were planted, all 15 that are native to North America have survived. Only two have survived that are not native to the United States: *M. poeppigii*, native to South America and *P. kamtschaticus*, native to Asia. Of the five succulent taxa that have perished, four are native to Africa: *C. orbiculata*, *D. cooperi*, *D. ‘Kelaidis’*, and *D. nubigenum*. This may indicate that *Delosperma* species native to Africa are not as likely to survive on a USDA plant hardiness zone 5 low-water green roof in a semi-arid, temperate climate in the United States. The other succulent to perish, *S. rupestre*, is native to Europe.

Succulent Plant Taxa Survival Chart															
Plant Taxa	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Survival %
<i>Agave parryi</i>		4			3	3	2,6	2,6					19	19	475%
<i>Echinocereus fendleri</i>			mass		6	6								4	n/a
<i>Echinocereus triglochidiatus</i>	18				18	18	18	18						4	22%
<i>Echinocereus triglochidiatus</i> white sand strain	2				2	2	2	2						1	50%
<i>Echinocereus viridiflorus</i>	11				8	8								2	18%
<i>Escobaria missouriensis</i>	mass				2	6	3	3						3	n/a
<i>Escobaria vivipara</i>	20				20	9	3	3						3	15%
<i>Hesperaloe parviflora</i>	8				8	8	6	6						4	50%
<i>Hesperaloe parviflora</i> (2)		5			5	5	3	3						4	80%
<i>Maihuenia poeppigii</i>		27			6	6	0	0		~	~	~		~	n/a
<i>Nolina microcarpa</i>	3				6	6	6	6						5	167%
<i>Opuntia aurea</i>	mass				mass	mass	mass	mass						mass	n/a
<i>Opuntia phaeacantha</i>	mass				mass	mass	mass	mass						mass	n/a
<i>Opuntia polyacantha</i>	mass				mass	mass	mass	mass						mass	n/a
<i>Pediocactus simpsonii</i>	10				10	5	3	3							n/a
<i>Phedimus kamtschaticus</i>	5				mass	mass	mass	mass						mass	n/a
<i>Phemeranthus calycinus</i>														mass	n/a
<i>Phemeranthus calycinus</i> (2)					12									mass	n/a
<i>Sedum lanceolatum</i>	64				mass	mass	mass	mass						mass	n/a
<i>Cotyledon orbiculata</i>						5									0%
<i>Delosperma cooperi</i>	12				mass	mass	6	6	0						0%
<i>Delosperma ‘Kelaidis’</i>	40				2	2	0								0%
<i>Delosperma ‘Kelaidis’</i> (2)		8			2	2	2	2	0						0%
<i>Delosperma nubigenum</i>	21				mass	mass	2	2	0						0%
<i>Sedum rupestre</i>					31	mass	mass	mass	0						0%

Figure 2. Succulent survival by year from 2007-2020. The plants categorized as ‘survive’ are listed alphabetically by genus at the top while those that are categorized as ‘perish’ are listed alphabetically by genus at the bottom. Note: #,# second number represents number of pups, ‘mass’ = many individual plants that have grown together over time.

Creeping Forbs

In the first two years after the green roof was installed, fall 2007- September 2009, 26 creeping forb taxa were planted (with *Eriogonum umbellatum* var. *aureum* being planted twice but counted as one taxa; Figure 3). After 12 years, 12 of the 26 plant taxa have survived, or 46%. Over the next two planting seasons, 2010-2012, 19 creeping forbs were planted (including a second *Cotula hispida*, therefore, only 18 ‘new’ taxa were planted). Of the 19 creeping forb taxa only 3, or 16% survived. Between the seasons of 2013-2018 two taxa were planted, *Genista tinctoria* and *G. lydia*, both of which have survived. In the past 12 years (2007-September 2019) 17 of the total 47 species, or 36%, have survived. *C. hispida*, *E. umbellatum* var. *aureum*, and *G. tinctoria* were trialed twice but were only counted as taxa once. Both plantings of *C. hispida* have perished, both *E. umbellatum* var. *aureum* survived, and *G. tinctoria* (2) was planted in 2019 therefore only data on the first planted *G. tinctoria* is included in the data as perished.

Since the original evaluation in 2012, seven years later, seven more creeping forb plant taxa have perished; *Acantholimon acerosum*, *Aethionema schistosum*, *Artemisia* sp., *C. hispida*, *Penstemon caespitosus*, *Streptanthus cordatus* and *Thymus neiceffi* (Schneider et al., 2014). Most form taproots, which have previously been shown to be less ideal for extensive green roofs (Nagase and Dunnett 2010). Many creeping forbs trialed on the green roof can reseed which has made tracking them more difficult, although that could be a factor in overall plant species success. In the original evaluation, *Erigeron vetensis*, *Heterotheca jonesii*, and *Veronica pectinata* were marked as having perished, but since, has reappeared through seeding. *E. vetensis* has declined in number but has had great overall success and has bloomed again (authors observation, April 2020). *E. leiomerus* was planted in 2011, perished between 2012-2014, then appeared in 2015, surviving for five years until perishing again. *T. neiceffi* survived on the green roof for nine years before perishing and should be reevaluated.

Of the subcategory of bun forming creeping forbs, three of the six, or 50%, have survived. *A. armenum*, *A. litvinovii*, and *Dianthus anatolicus* have been successful in survival, surviving for 11 total years so far. Although technically successful, each species has declined in number without showing any remarkable new growth. This confirms what was stated in the original study, the small size and slow growth of bun forming creeping forbs may be unfavorable for covering large areas. *A. acerosum* survived for six years before perishing and could be retried as its two relatives *A. armenum* and *A. litvinovii* have been successful.

Of the 47 trialed creeping forb plant taxa, 16 are native to the United States. Of the 16 that are native, seven have survived, or 44%. Thirty-one of the trialed plants are non-native, of them, 13 survived, or 42%. Despite the results being almost equal, creeping forbs that are native to this country, and especially our ecoregion, theoretically should be more likely to survive these harsh conditions, however more evaluation would be required.

Creeping Forbs Plant Taxa Survival Chart															
Plant Taxa	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Survival %
<i>Acantholimon armenum</i>		5			1	1	1	1					1	1	20%
<i>Acantholimon litvinovii</i>		5			1	1	1	1	2					1	20%
<i>Chrysanthemum weyrichii</i>					16	mass	mass	mass		~	~			1	6%
<i>Dianthus anatolicus</i>		28	19		10	11	17	17	10				10	10	36%
<i>Draba hispanica</i>	3				10	mass							~	1	33%
<i>Erigeron vetensis</i>				27		0								6	22%
<i>Eriogonum umbellatum</i> var. <i>aureum</i>	15		12		9	9	9	9	2					3	20%
<i>Eriogonum umbellatum</i> var. <i>aureum</i> (2)		10			6	9	mass	mass	2				2	2	20%
<i>Genista tinctoria</i> 'Royal Gold'								1					1	3	300%
<i>Genista lydia</i>										1				4	400%
<i>Heterotheca jonesii</i>	32					0								mass	n/a
<i>Penstemon pinifolius</i>			6		18	5	5	5						mass	n/a
<i>Phlox bifida</i> 'Betty Blake'	1				1	1	5	5						1	100%
<i>Physaria bellii</i>	64				17									7	11%
<i>Silene schafta</i>				23	10	10	10+	10+						7	30%
<i>Teucrium graphaloides</i> (2)													3	3	100%
<i>Veronica lithanensis</i>	9				5	5	7	8						2	22%
<i>Veronica pectinata</i>	mass				0									2	n/a
<i>Veronica thymoides</i>	13				6	6	6	6						1	8%
<i>Acantholimon acerosum</i>		4			3	2	2	1	0						0%
<i>Aethionema schistosum</i>	3				3	1	0								0%
<i>Anacyclus maroccanus</i>					6	0									0%
<i>Antennaria parvifolia</i> 'McClintock'	64				1	0									0%
<i>Arenaria alfacarensis</i>	16				0										0%
<i>Artemisia</i> sp.	32					10									0%
<i>Clematis columbiana</i> var. <i>temuloba</i>					2	0									0%
<i>Cotula hispida</i>	9				mass	mass	0								0%
<i>Cotula hispida</i> (2)				21	10	20	mass	mass	0						0%
<i>Draba streptocarpa</i>	32				0										0%
<i>Dryas octopetala</i> var. <i>hookeriana</i>				10	0										0%

<i>Erigeron leiomerus</i>					6	0											0	0%
<i>Herniaria glabra</i>	2					0												0%
<i>Herniaria glabra</i> 'Sea Foam'	18					0												0%
<i>Heteropappus</i> sp.				2		0												0%
<i>Leucanthemum atlanticum</i>						9												0%
<i>Paronychia kapela</i> subsp. <i>serpyllifolia</i>	16					0												0%
<i>Penstemon casepitosus</i>	11					2	2	0										0%
<i>Penstemon grahamii</i>				1		0												0%
<i>Penstemon linarioides</i> subsp. <i>coloradensis</i>	12					2	0											0%
<i>Petrophytum caespitosum</i>				3		0												0%
<i>Potentilla hyparctica</i>	16					0												0%
<i>Potentilla nepalensis</i> 'Shogran'				3		0												0%
<i>Salvia</i> 'Eskihar'						3	0											0%
<i>Scutellaria prostrata</i>						20	0											0%
<i>Scutellaria scordifolia</i>				4		0												0%
<i>Streptanthus cordatus</i>				32		30												0%
<i>Teucrium aroanium</i>				4	4												0	0%
<i>Teucrium graphaloides</i>					8								~	0				0%
<i>Thymus neiceffi</i>	9				mass	mass	2	2									0	0%
<i>Vella spinosa</i>					3	0												0%

Key
Survive
Seed
Perish
Unknown
~ Struggling

Figure 3. Creeping forbs survival by year from 2007-2020. The plants categorized as 'survive' are listed alphabetically by genus at the top while those that are categorized as 'perish' are listed alphabetically by genus at the bottom.

Upright Forbs

To date, a total of 24 upright forb taxa have been trialed on the DBG green roof (Figure 4). From installation in falls 2007- 2009, ten total upright forb taxa were planted. Of the ten, four, or 40% survived. Between fall 2010- 2012, 14 more upright forbs were planted for trial and out of the 14 planted only one, or 7%, survived. The total survival rate for upright forb taxa on the green roof is five of 23, or 22%.

As previously found in the five-year evaluation (Schneider et al., 2014), most upright forbs in low-irrigation green roofs have limited use in semi-arid climates. However, five of the plant taxa, *Echium amoenum*, *Ipomopsis aggregata*, *Iris germanica* dwarf, *Salvia pachyphylla*, and *Thelesperma ambiguum* have survived 12 years, demonstrating that upright forb taxa should be considered for semi-arid environments on green roofs after careful trial. Note: *Hypericum olympicum* in the 2014 evaluation was listed as woody shrub (Schneider et al., 2014). In this evaluation it is considered an upright forb as it performs more as a perennial than a subshrub in Colorado as it does not have above ground woody material that overwinters.

In this evaluation, the upright forb category has had the lowest success rate for semi-arid climates on a green roof. Of the 17 perished upright forbs, 11 taxa perished within one year then all the rest except *E. elatior* and *H. olympicum* have perished within three years of planting.

This suggests that these upright forbs that are not adapted to semi-arid, temperate green roofs will perish quickly. *E. amoenum*, *I. aggregata*, and *I. germanica* dwarf are quite successful and have grown exponentially, eventually massing. From the results of this evaluation, these species are recommended for use on semi-arid green roofs.

Erigeron elatior and *S. pachyphylla* have consistently survived, however, *E. elatior* has perished in 2020, although both were mistakenly marked as perished in the original study. This may be due to their appearance, such as early dormancy, when they were counted in the fall (Schneider et al. 2014). *E. elatior* survived for eight years before eventually perishing; it is an alpine, therefore, premature die back is normal. However, the species that perished may have done so due to the lack of winter snow cover and nighttime temperatures that reached -1°C to 4°C.

Although three *S. pachyphylla* were originally planted during the installation of the green roof in 2007, only one has remained. *Baileya multiradiata* seeds were sown in 2009 but did not germinate until 2016 when it appeared for two years then perished; this lifecycle is normal as *B. multiradiata* is biennial, but it is unknown if the plant flowered before dying. In general, *B. multiradiata* germination is erratic which could play a role in their low germination rate in this evaluation.

On the DBG green roof, *T. ambiguum* has survived for 12 years but is now showing a decline in number, although it has varied in its relative vigor seasonally, showing potential for its use for green roofs in semi-arid climates. As discussed in the original study, *Ipomopsis* species reseed which allow them to move around the roof to areas that provide ideal conditions (Schneider et al. 2014); Overall, their reseeding could allow them to be more successful than non-seeding upright forbs. Originally *I. rubra* was planted on the green roof in 2007 and did survive and reseed up until the first evaluation (Schneider et al, 2014) but then eventually died out to be replaced entirely by its close relative, *I. aggregata*. While *I. aggregata* was first planted in 2010, this biennial quickly flowered, reseeded and continued to dominate the shallow portions of the green roof. One of the surviving plants, *I. germanica* dwarf develops underground rhizomes which may benefit its survival due to its horizontal growth habit.

Provenance in the upright forbs category include taxa ranging from North America to southeastern Asia. Of the surviving seven upright forbs, five, or 71%, are native to the United States. The other two surviving taxa are *E. amoenum*, which is native to the Middle East, and *I. germanica* dwarf, native to southeastern Europe. Of the taxa that perished, eight of the 16, or 50%, are native to the United States. The data from this evaluation suggests that upright forbs that are native to our continent may be more likely to survive on a semi-arid green roof, although further evaluation is necessary.

Upright Forbs Plant Taxa Survival Chart															
Plant Taxa	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Survival %
<i>Echium amoenum</i>	4				4	4	4	4					8	mass	n/a
<i>Iris germanica</i> dwarf	26				26	26								mass	n/a
<i>Ipomopsis aggregata</i>				18										mass	n/a
<i>Salvia pachyphylla</i>	3				1	1	1	1					1	1	n/a
<i>Thelesperma ambiguum</i>	32				mass	mass								3	n/a
<i>Baileya multiradiata</i>			seed	0						~	~	0			0%
<i>Braya alpine</i>				12	0										0%
<i>Campanula incurva</i>					6	2	0								0%
<i>Campanula kemulariae</i>				1	0										0%
<i>Erigeron elatior</i>					8	8	8	8						0	0%
<i>Erysimum capitatum</i> var. <i>purshii</i>					31	28	0								0%
<i>Euphorbia polychroma</i> 'Candy'		mass	0												0%
<i>Hedeoma</i> clone					3	0									0%
<i>Hypericum olympicum</i>					4	4	4+	4+	0						0%
<i>Ipomopsis rubra</i>		16								0					0%
<i>Olsynium biflorum</i>					4	0									0%
<i>Penstemon angustifolius</i>				2	1	1	0								0%
<i>Penstemon cyananthus</i>				6	6	0									0%
<i>Penstemon fendleri</i>	8	0													0%
<i>Phacelia campanularia</i>			seed	0											0%
<i>Putoria calabrica</i>					32	0									0%
<i>Salvia cryptantha</i>					5	0									0%
<i>Townsendia eximia</i>	24				0										0%
<i>Ziziphora bungeana</i>					17	0									0%

Figure 4. Upright forbs survival by year from 2007-2020. The plants categorized as 'survive' are listed alphabetically by genus at the top while those that are categorized as 'perish' are listed alphabetically by genus at the bottom.

Graminoids

A total of 11 graminoid species have been trialed on the DBG green roof (Figure 5). Out of the 11, four of them have been planted or seeded in 2019, therefore will not be included in the overall survival rate. Between fall 2007- 2009 one graminoid was planted, *Panicum virgatum*, which survived eight years before perishing. Between fall 2010-2012, six more graminoids were trialed. Of the six, three survived, or 50%. The surviving graminoids planted between 2007-2019 include *A. ternaries*, *A. virginicus*, and *M. montana*, each of them have survived for seven years, showing great potential. Graminoids overall (excluding plantings in 2019) have a three of seven, or 43% survival rate. This indicates that specific graminoid plant taxa should continue to be trialed to determine successful candidates for semi-arid green roofs. As of April 2020, all graminoids planted in 2019 have survived but will need a longer evaluation period to determine their overall success.

Of the graminoids on the green roof, *A. ternarius* had the highest survival at 40%, followed by *A. virginicus* as a close second. All three of the surviving graminoids have decreased in number, *M. montana* more than the others, starting with 64 plants and as of 2019 only three are surviving. Although all have decreased in number, the graminoids that remain are showing great potential.

Our data suggest that the genus *Festuca* is not a good candidate for low water green roofs in high elevation, semi-arid climates. Sixty-four *F. brachyphylla* were planted and all perished within a year. Similarly, an evaluation in arid Iran found two relatives, *F. aurundinacea* and *F. ovina*, to be unsuccessful (Vahdati et al., 2017) although Nektarios et al. (2014) reported that *F. aurundinacea* was a successful green roof plant with irrigation. Erwin et al. (2019) also found that three *Festuca* spp. performed well.

P. virgatum survived for eight years on the DBG roof top and suddenly perished in 2015 yet thrived for Erwin et al. (2019). This could be due to differences in climate or substrate depth, although *P. virgatum* should be re-trialed to confirm this conclusion. It is speculated that *P. virgatum*, along with *M. emersleyi* and *Poa fendleriana*, suddenly perished in 2015 because of a November freeze in 2014 (author observation). And it is suspected that they would have otherwise survived.

Sporobolus heterolepis, planted in 2019, has been established for one year and has been successful in producing another plant. The success of *S. heterolepis* on the DBG green roof contrasts with what was found by Tran et al. (2019) that this graminoid did not thrive with lower precipitation levels. *Bouteloua gracilis* made a spontaneous appearance in 2019 and was successful overwintering.

Graminoids Plant Taxa Survival Chart															
Plant Taxa	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Survival %
<i>Andropogon ternarius</i>						10	4	4					3	4	40%
<i>Andropogon virginicus</i>						10	7	7					3	3	30%
<i>Andropogon</i> sp.													3		n/a
<i>Bouteloua gracilis</i> 'Blonde Ambition'													1	1	100%
<i>Muhlenbergia montana</i>						64	64	64	5			3	3	3	5%
<i>Muhlenbergia</i> sp.													1		n/a
<i>Sporobolus heterolepis</i>													2	3	150%
<i>Festuca brachyphylla</i>						64	0								0%
<i>Muhlenbergia emersleyi</i>						10	10	10	0						0%
<i>Panicum virgatum</i>	6				6	6	6	6	0						0%
<i>Poa fendleriana</i>						64	2	2	0						0%

Figure 5. Graminoids survival by year from 2007-2020. The plants categorized as 'survive' are listed alphabetically by genus at the top while those that are categorized as 'perish' are listed alphabetically by genus at the bottom.

Woody Shrubs

In this evaluation, the woody shrubs were planted on or near the berm area on the green roof where substrate reaches its maximum depth of around 45 cm. Although less represented in literature, these taxa have shown good overall survivability in our study (Figure 6). Between 2007-2009, 12 taxa were planted, of the 12 planted, ten, or 83%, survived. Two taxa were planted between 2010-2012 and one of the two, or 50%, survived. In 2014, *Fallugia paradoxa* appeared and has continued to survive. Of the total 15 woody shrub taxa, 13, or 87%, have survived. Overall, woody shrubs show great potential for use on green roofs in a semi-arid climate.

One of the most remarkable observations on this green roof was the spontaneous arrival of *F. paradoxa*. It appeared in 2014 and has become established, blooming, and producing seed. *S. greggii* was the most successful woody shrub (technically a subshrub) on this semi-arid green roof, growing into a mass from 5 original plants throughout 11 years. The green roof has proven to be an ideal environment for this plant taxa sharing similar qualities to its natural environment such as high light and low moisture. *Chamaebatiaria millefolium* has also been quite successful, reseeding and doubling the original number of plants. This is likely due to its adapted xeric properties.

Although *Ericameria nauseosa* subsp. *nauseosa* var. *nauseosa*'s survival rate is listed as 80%, it has been vigorous and has seeded several times (some specific years unknown). It has emerged throughout the rooftop and is continuously being removed by the gardener seasonally. *Cercocarpus breviflorus* and *Chilopsis linearis* reappear seasonally and bloom. The surviving plants have maintained a steady population annually. No taxa have declined more than 50% and seem to maintain a steady number annually. One exception is *A. confertifolia*, at 17%. Despite their high success rate, woody shrubs are seldom mentioned in literature; this study suggests that they are candidates for further evaluation.

The data suggest that provenance may play a role on these taxa's survivability in a semi-arid climate. Eleven of the 13 surviving taxa, or 85%, are native to the United States. The other two surviving taxa, *Ephedra minuta* and *Euonymus nanus* var. *turkestanicus* are native to Asia. More shrubs native to Asia should be trialed to determine if provenance is a determining factor on success rate. Of the two taxa that perished, *Cytisus purgans* 'Spanish Gold' is native to Europe, and the other one, *Fendlera rupicola* var. *wrightii*, is native to the United States. The data collected suggests that provenance could be a factor in the success rate of woody plant taxa on green roofs.

Woody Shrubs Plant Taxa Survival Chart																
Plant Taxa	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	Survival %	
<i>Amorpha fruticosa</i> ‘Nana’			3		3	4	3	3	3				3	3	100%	
<i>Atriplex confertifolia</i>				6	6	2			1	1	1			1	17%	
<i>Arctostaphylos</i> ‘Lauren’s Best’	1				1	1	1	1						1	100%	
<i>Arctostaphylos patula</i>	3				2	2	2	2						2	67%	
<i>Arctostaphylos</i> x <i>coloradoensis</i> ‘Cascade’	2				1	2	2	2						1	50%	
<i>Cercocarpus breviflorus</i>	3				3	3	3	3						3	100%	
<i>Chamaebatiaria millefolium</i>	3				3	3	3	3					7	6	200%	
<i>Chilopsis linearis</i>	2				2	2	2	2	1				1	1	50%	
<i>Ephedra minuta</i>				6	6	5	6	6						3	50%	
<i>Ericameria nauseosa</i> subsp. <i>nauseosa</i> var. <i>nauseosa</i>	5						1	1		<div>Key</div> <div> Survive</div> <div> Seed</div> <div> Perish</div> <div> Unknown</div> <div> Struggling</div>			4	80%		
<i>Euonymus nanus</i> var. <i>turkestanicus</i>	5				5	5	5	5					4			
<i>Fallugia paradoxa</i>								1						1	100%	
<i>Salvia greggii</i>		5			5	5	5	5						4	mass	n/a
<i>Cytisus purgans</i> ‘Spanish Gold’	3				1	1	1	1	0						0%	
<i>Fendlera rupicola</i> var. <i>wrightii</i>	1				0										0%	

Figure 6. Woody shrub survival by year from 2007-2020. The plants categorized as 'survive' are listed alphabetically by genus at the top while those that are categorized as 'perish' are listed alphabetically by genus at the bottom.

CONCLUSIONS

Select plant taxa in each plant form category were successful on the green roof in this evaluation. Succulents showed greatest potential for survival on green roofs in a semi-arid climate. But the less studied shrub forms came in at a close second with only two taxa perishing. This evaluation demonstrates that all plant form categories can be utilized on a semi-arid green roof, with (forms) being most successful. Forms such as upright forbs, creeping forbs, and graminoids, in general, were less successful. Therefore, taxa from these plant form types should be carefully selected for green roof use.

All plant forms demonstrate yearly fluctuation of relative success within taxa (i.e. individual plants have perished or survived), however relative plant cover on the green roof has remained constant over time. One of the most unexpected observations was *F. paradoxa*'s spontaneous arrival on the green roof.

Seeding plants from all plant forms, such as *E. nauseosa* subsp. *nauseosa* var. *nauseosa*, *E. leiomerus*, and *I. aggregata* may increase success on a green roof due to vital seed dispersion and reappearance. Shade created by the installation of different plant heights such as woody shrubs may also increase the survivability of other plant forms, similar to how shade from solar

panels improved green roof plant survivability in another Colorado study (Bousselot et al. 2017). Between low precipitation and low quantities of supplemental irrigation, this green roof demonstrates plant taxa that may be used on green roofs in other semi-arid climates.

In some cases, provenance may explain survivability in each of the plant form categories. In general, plants that are native to North America were more likely to survive on the green roof than plants that were not native; however, not equally or consistently across all plant categories. Factors such as low levels of supplemental irrigation, variable shading by taller plants, and microclimates could have greatly influenced the relative success or failure of species no matter their provenance. Additional long-term evaluations for both taxa survivability and provenance among green roofs in other regions of North America are needed.

LITERATURE CITED

- Aguiar, A.C., S.A. Robinson and K. French. 2019. "Friends with benefits: The effects of vegetative shading on plant survival in a green roof environment." *PLoS ONE*, 14(11): 1–16. <https://doi.org/10.1371/journal.pone.0225078>
- Bousselot, J.M., J.E. Klett and R.D. Koski. 2010. "Extensive green roof species evaluations using digital image analysis." *HortScience*, 45(8): 1288-1292. <https://doi.org/10.21273/HORTSCI.45.8.1288>
- Bousselot, J.M., J.E. Klett and R.D. Koski. 2011. "Moisture content of extensive green roof substrate and growth response of 15 temperate plant species during dry down." *HortScience*, 46(3): 518-522. <https://doi.org/10.21273/HORTSCI.46.3.518>
- Bousselot, J., T. Slabe, J. Klett and R. Koski. 2017. "Photovoltaic array influences the growth of green roof plants." *Journal of Living Architecture*, 4(3): 9-18. <https://doi.org/10.46534/jliv.2017.04.03.009>
- Braaker, S., M.K. Obrist, J. Ghazoul, M. Moretti and K. Cuddington. 2017. "Habitat connectivity and local conditions shape taxonomic and functional diversity of arthropods on green roofs." *Journal of Animal Ecology*, 86(3): 521–531. DOI: [10.1111/1365-2656.12648](https://doi.org/10.1111/1365-2656.12648)
- Cao, J., S. Hu, Q. Dong, L. Liu, and Z. Wang. 2019. "Green roof cooling contributed by plant species with different photosynthetic strategies." *Energy and Buildings*, 195: 45–50. <https://doi.org/10.1016/j.enbuild.2019.04.046>
- Chagolla-Aranda, M.A., E. Simá, J. Xamán, G. Álvarez, I. Hernández-Pérez, and E. Téllez-Velázquez. 2017. "Effect of irrigation on the experimental thermal performance of a green roof in a semi-warm climate in Mexico." *Energy and Buildings*, 154:232–243. <https://doi.org/10.1016/j.enbuild.2017.08.082>
- CoCoRaHS. n.d. "Community Collaborative Rain, Hail and Snow Network." *CoCoRaHS*, www.cocorahs.org/.
- Dunnett, N. and A. Nagase. 2010. "Drought tolerance in different vegetation types for extensive green roofs: effects of watering and diversity." *Landscape and Urban Planning*, 97(4): 318-327.
- Dvorak, Bruce, and Astrid Volder. 2010. "Green Roof Vegetation for North American Ecoregions: A Literature Review." *Landscape and Urban Planning*, Elsevier. <https://doi.org/10.1016/j.landurbplan.2010.04.009>
- Erwin, J. and J. Hensley. 2019. "Plants with horticultural and ecological attributes for green roofs in a cool, dry climate." *HortScience*, 54(10): 1703–1711. <https://doi.org/10.21273/HORTSCI13893-19>
- Getter, K.L. and D.B. Rowe. 2006. "The role of extensive green roofs in sustainable development." *HortScience*, 41(5): 1276-1285.

- Graceson, A., J. Monaghan, N. Hall, and M. Hare. 2014. "Plant growth responses to different growing media for green roofs." *Ecological Engineering*, 69: 196–200.
<https://doi.org/10.1016/j.ecoleng.2014.03.067>
- Kew Science. n.d. Plants of the World Online. <http://www.plantsoftheworldonline.org/>.
- Klein, P.M. and R. Coffman. 2015. "Establishment and performance of an experimental green roof under extreme climatic conditions." *Science of the Total Environment*, 512: 82-93.
- Köhler, M. and P. Poll. 2010. "Long-term performance of selected old Berlin greenroofs in comparison to younger extensive greenroofs in Berlin." *Ecological Engineering* 36: 722-729.
- Kotsiris, G., P.A. Nektarios, N. Ntoulas and G. Kargas. 2013. "An adaptive approach to intensive green roofs in the Mediterranean climatic region." *Urban Forestry and Urban Greening*, 12(3): 380-392. <https://doi.org/10.1016/j.ufug.2013.05.001>
- Lubell, J.D., B. Connolly and K.N. Jones. 2017. "Ten-year persistence of native plant species on a green roof in Northeast :US." *Native Plants Journal*, 18(3): 227–234.
<https://www.muse.jhu.edu/article/680008>.
- Lundholm, J.T. and M. Cadotte. 2015. "Green roof plant species diversity improves ecosystem multifunctionality." *Journal of Applied Ecology*, 52(3): 726–734. <https://doi.org/10.1111/1365-2664.12425>
- Lundholm, J., J.S. MacIvor, Z. MacDougall, and M. Ranalli. 2010. "Plant species and functional group combinations affect green roof ecosystem functions." *PLoS ONE*, 5(3): 1–11.
<https://doi.org/10.1371/journal.pone.0009677>
- Monterusso, M.A., D.B. Rowe, and C.L. Rugh. 2005. "Establishment and persistence of *Sedum* spp. and native taxa for green roof applications." *HortScience*, 40(2), 391–396.
- Nektarios, P.A., I. Amountzias, I. Kokkinou and N. Ntoulas. 2011. "Green roof substrate type and depth affect the growth of the native species *Dianthus fruticosus* under reduced irrigation regimens." *HortScience*, 46(8): 1208-1216. <https://doi.org/10.21273/HORTSCI.46.8.1208>
- Nektarios, P.A., I. Kokkinou and N. Ntoulas. 2020. "The effects of substrate depth and irrigation regime, on seeded *Sedum* species grown on urban extensive green roof systems under semi-arid Mediterranean climatic conditions." *Journal of Environmental Management*, p.111607.
- Nektarios, P.A., N. Ntoulas, G. Kotopoulis, T. Ttoulou and P. Ilia. 2014. "*Festuca arundinacea* drought tolerance and evapotranspiration when grown on two extensive green roof substrate depths and under two irrigation regimes." *European Journal of Horticultural Science*, 79(3): 142-149.
- Nektarios, P.A., N. Ntoulas, E. Nydrioti, I. Kokkinou, E.M. Bali and I. Amountzias. 2015. "Drought stress response of *Sedum sediforme* grown in extensive green roof systems with different substrate types and depths." *Scientia Horticulturae*, 181: 52-61.
<https://doi.org/10.1016/j.scienta.2014.10.047>
- Oberndorfer, E., J. Lundholm, B. Bass, R. Coffman, H. Doshi, N. Dunnett, S. Gaffin, M. Köhler, K. Liu, and B. Rowe. 2007. "Green roofs as urban ecosystems: ecological structures, functions, and services." *Bioscience*, 57(10): 823-833. doi:10.1641/b571005
- Reyes, R., W. Bustamante, J. Gironás, P.A. Pastén, V. Rojas, F. Suárez, S. Vera, F. Victorero and C.A. Bonilla. (2016). "Effect of substrate depth and roof layers on green roof temperature and water requirements in a semi-arid climate." *Ecological Engineering*, 97: 624–632.
<https://doi.org/10.1016/j.ecoleng.2016.10.025>
- Schneider, A., M. Fusco, J. Bousselot. 2014. "Observations on the survival of 112 plant taxa on a green roof in a semi-arid climate." *Journal of Living Architecture*, 1(5): 10-30.
<https://doi.org/10.46534/jliv.2014.02.01.010>
- Shafique M., R. Kim, M. Rafiq. 2018. "Green roof benefits, opportunities and challenges – A review." *Renewable and Sustainable Energy Reviews*, 90: 757-773
<https://doi.org/10.1016/j.rser.2018.04.006>

- Thuring, C.E., R.D. Berghage and D.J. Beattie. 2010. "Green roof plant responses to different substrate types and depths under various drought conditions." *HortTechnology*, 20(2): 395-401. <https://doi.org/10.21273/HORTTECH.20.2.395>
- Tran, S., J.T. Lundholm, M. Staniec, C.E. Robinson, C.C. Smart, J.A. Voogt and D.M. O'Carroll. 2019. "Plant survival and growth on extensive green roofs: A distributed experiment in three climate regions." *Ecological Engineering*, 127: 494–503. <https://doi.org/10.1016/j.ecoleng.2018.09.027>
- Vahdati, N., A. Tehranifar and F. Kazemi. 2017. "Assessing chilling and drought tolerance of different plant genera on extensive green roofs in an arid climate region in Iran." *Journal of Environmental Management*, 192: 215–223. DOI: [10.1016/j.jenvman.2017.01.027](https://doi.org/10.1016/j.jenvman.2017.01.027)