Open Access

Mesenchymal Stem Cell Transplantation for COVID-19

Attapon Cheepsattayakorn^{1*}, Ruangrong Cheepsattayakorn²

¹10th Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand

² Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

***Corresponding Author:** Attapon Cheepsattayakorn, 10th Zonal Tuberculosis and Chest Disease Center, 143 Sridornchai Road Changklan Muang Chiang Mai 50100, Thailand. Tel: +66 53140767; +66 53276364; Email: Attapon1958@gmail.com

Received Date: 01-04-2020; **Accepted Date:** 03-04-2020; **Published Date:** 10-04-2020

Copyright © 2020 by Cheepsattayakorn A, et al. All rights reserved. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

In-vitro, Mesenchymal Stem Cell (MSC) populations with potentials of similar multi-lineage differentiation have been obtained from several Bone Marrow (BM) and non-bone marrow tissues, including umbilical cord, placenta, amniotic fluid, adipose tissue, and peripheral blood [1-10]. The clonogenic BM-human MSCs fraction ranges from 10 to 100 Colony-Forming Unit-Fibroblast (CFU-F) per 10⁶ Marrow Mononuclear Cells (MNCs) [11]. BM-human MSCs are characterized by lacking CD11b, CD14, CD19, D34, CD45, CD79 α , and Human Leukocyte Antigen (HLA)-DR expression; positive expression of surface antigens CD73, CD90, and CD105; multipotency (i.e., chondrogenic, osteogenic, and adipogenic); and their adherence to plastic [11]. By the year 2000, clinicians increasingly had become interested in intravenously applied MSC therapy [12]. A previous study demonstrated that both human and murine MSCs can induce immune suppression by attracting and killing auto reactive T cells via FasL, therefore stimulating Transforming Growth Factor-beta (TGF- β) production by macrophages and generation of regulatory T cells [13]. The dying T cells that is caused by the interaction involving the MSC-induced Monocyte Chemoattractant Protein-1 (MCP-1) secretion in turn activate macrophages to produce TGF- β , then stimulating regulatory T cells and promoting immune tolerance [14]. The capacity of MSCs for *in-vivo* differentiation and engraftment and by their efficacy in promoting wound healing highlighted its clinical relevance [15-21].

In 2006, the International Society for Cellular Therapy came up with the guidelines for MSC characterization for standardization the MSC biology, definition, isolation, and characterization criteria, *in-vivo* relevance, and ethical and institutional regulations for its

Cheepsattayakorn A | Volume 1; Issue 1 (2020) | JCIM-1(1)-01 | Editorial

Citation: Cheepsattayakorn A, Cheepsattayakorn R. Mesenchymal Stem Cell Transplantation for COVID-19. J Clin Immunol Microbiol. 2020; 1 (1):1-4.

DOI: <http://dx.doi.org/10.46889/JCIM.2020.1101>

clinical use [11]. Since the COVID-19 pandemic, there are several ongoing trials that have been studied in China, such as the ClinicalTrials.gov identifiers: NCT04252118, NCT04273646, NCT04276987, NCT04293692, NCT04302519, NCT04288102, etc. for fighting against severe COVID-19 or COVID-19 pneumonia [22-27]. MSCs can decrease the overproduction of immune cells caused by a reaction to the COVID-19 and decrease excessive levels of inflammatory substances, contributing to regulating the immune system and recovering to the normal status, particularly of the elderly patients [28].

In conclusion, human MSCs are currently being evaluated as a stem cell treatment for a number of diseases, particularly severe COVID-19 and have been demonstrated to be safe in clinical trials. Nevertheless, further studies are urgently needed to investigate and optimize a number of variables in the human MSC culture environment by developing a bioprocess that can be operated in accordance with the Good Manufacturing Product (GMP).

References

1. Williams AR, Hare JM. Mesenchymal stem cells: biology, pathophysiology, translational findings, and therapeutic implications for cardiac disease. *Circ Res* 2011; 109 (8): 923-940.
2. Lee OK, Kuo TK, Chen WM, Lee KD, Hsieh SL, Chen TH. Isolation of multipotent mesenchymal stem cells from umbilical cord blood. *Blood* 2004; 103 (5): 1669-1675.
3. Wang HS, Hung CS, Peng ST, Huang CC, Wei HM, Guo YJ, et al. Mesenchymal stem cells in the Wharton's jelly of the human umbilical cord. *Stem Cells* 2004; 22 (7): 1330-1337.
4. Tondreau T, Meuleman N, Delforge A, Dejeneffe M, Leroy R, Massy M, et al. Mesenchymal stem cells derived from CD133-positive cells in mobilized peripheral blood and cord blood : proliferation, Oct4 expression, and plasticity. *Stem Cells* 2005; 23 (8): 1105-1112.
5. Vellasamy S, Sandrasaigaran P, Vidyadaran S, George E, Ramasamy R. Isolation and characterization of mesenchymal stem cells derived from human placenta tissue. *World J Stem Cells* 2012; 4 (6): 53-61.
6. Anker PS l't, Scherjon SA, Kleijburg-van der Keur C, Noort WA, Claas FH, Willemze R, et al. Amniotic fluid as a novel source of mesenchymal stem cells for therapeutic transplantation. *Blood* 2003; 102 (4): 1548-1549.
7. De Coppi P, Bartsch Jr G, Siddiqui MM, Xu T, Santos CC, Perin, et al. Isolation of amniotic stem cell lines with potential for therapy. *Nat Biotechnol* 2007; 25 (1): 100-106.
8. Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, et al. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell* 2002; 13 (12): 4279-4295.
9. Gimble J, Guilak F. Adipose-derived adult stem cells: isolation, characterization, and differentiation potential. *Cytotherapy* 2003; 5 (5): 362-369.
10. Zvaifler NJ, Marinova-Mutafchieva L, Adams G, Edwards CJ, Moss J, Burger JA, et al. Mesenchymal precursor cells in the blood of normal individuals. *Arthritis Res* 2000; 2 (6): 477-488.

11. Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells, the International Society for Cellular Therapy position statement. *Cytotherapy* 2006; 8: 315-317.
12. Horwitz EM, Gordon PL, Koo WK, Marx JC, Neel MD, McNall RY, et al. Isolated allogeneic bone marrow-derived mesenchymal cells engraft for cell therapy of bone. *Proc Natl Acad Sci* 2002; 99: 8932-8937.
13. Akiyama K, Chen C, Wang D, Xu X, Qu C, Yamaza T, et al. Mesenchymal stem cell-induced immunoregulation involves Fas ligand/Fas-mediated T cell apoptosis. *Cell Stem Cell* 2012; 10 (5): 544-555.
14. Choi H, Lee RH, Bazhanov N, Oh JY, Prockop DJ. Anti-inflammatory protein TSG-6 secreted by activated MSCs attenuates zymosan-induced mouse peritonitis by decreasing TLR2/NF-Kb signaling in resident macrophages. *Blood* 2011; 118: 330-338.
15. Karp JM, Leng Teo GS. Mesenchymal stem cell homing: the devil is in the details. *Cell Stem Cell* 2009; 4 (3): 206-216.
16. Sivanathan KN, Gronthos S, Rojas-Canales D, Thierry B, Coates PT. Interferon-gamma modification of mesenchymal stem cells : implications of autologous and allogeneic mesenchymal stem cell therapy in allotransplantation. *Stem Cell Rev* 2014; 10 (3): 351-375.
17. Nemeth K, Keane-Myers A, Brown JM, Metcalfe DD, Gorham JD, Bundoc VG, et al. Bone marrow stromal cells use TGF-beta to suppress allergic responses in a mouse model of ragweed-induced asthma. *Proc Natl Acad Sci* 2010; 107 (12): 5652-5657.
18. Ren G, Zhang L, Zhao X, Xu G, Zhang Y, Roberts AI, et al. Mesenchymal stem cell-mediated immunosuppression occurs via concerted action of chemokines and nitric oxide. *Cell Stem Cell* 2008; 2 (2): 141-150.
19. English K, French A, Wood KJ. Mesenchymal stromal cells: facilitators of successful transplantation? *Cell Stem Cell* 2010; 7 (4): 431-442.
20. Murphy MB, Moncivais K, Caplan AI. Mesenchymal stem cells: environmentally responsive therapeutics for regenerative medicine. *Exp Mol Med* 2013; 45: e54.
21. Tyndall A. Mesenchymal stem cell treatments in rheumatology: a glass half full? *Nat Rev Rheumatol* 2014; 10 (2): 117-124.
22. Beijing 302 Hospital, China. ClinicalTrials.gov Identifier: NCT04252118. Mesenchymal stem cell treatment for pneumonia patients infected with 2019 novel coronavirus. Available at: https://clinicaltrials.gov/ct2/show/NCT04252118?show_xprt=Y (accessed on March 28, 2020).
23. Wuhan Union Hospital, China. ClinicalTrials.gov Identifier: NCT04273646. Study of human umbilical cord mesenchymal stem cells in the treatment of novel coronavirus severe pneumonia. Available at: https://clinicaltrials.gov/ct2/show/NCT04273646?show_xcprt=Y (accessed on March 28, 2020).
24. Ruijin Hospital, China. ClinicalTrials.gov Identifier: NCT04276987. A pilot clinical study on inhalation of mesenchymal stem cells exosomes treating severe novel coronavirus pneumonia. Available at: https://clinicaltrials.gov/ct2/show/NCT04276987?show_xprt=Y (accessed on March 28, 2020).
25. Puren Hospital Affiliated to Wuhan University of Science and Technology, China. ClinicalTrilas.gov Identifier: NCT04293692. Therapy for pneumonia patients infected by 2019 novel coronavirus. Available at: https://clinicaltrials.gov/ct2/show/NCT04293692?show_xprt=Y (accessed on March 28, 2020).

26. CAR-T (Shanghai) Biotechnology Co., Ltd., China. ClinicalTrials.gov Identifier: NCT04302519. Novel coronavirus induced severe pneumonia treated by dental pulp mesenchymal stem cells. Available at: https://clinicaltrials.gov/ct2/show/NCT04302519?show_xprt=Y (accessed on March 28, 2020).
27. Beijing 302 Hospital, China. ClinicalTrials.gov Identifier: NCT04288102. Treatment with mesenchymal stem cells for severe coronavirus disease 2019 (COVID-19). Available at: <https://clinicaltrials.gov/ct2/show/NCT04288102> (accessed on March 28, 2020).
28. Liying Z. China.org.cn. Mesenchymal stem cell therapy shows promise in treating COVID-19. Published online on March 7, 2020. Available at: www.china.org.cn/china/2020-03/07/content_75785868.htm (accessed on March 28, 2020).