Open Access

Opinion Article

Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology

Win L Chiou^{1*}

¹Chiou Consulting Inc, 8552 Johnston Road, Burr Ridge, Illinois, 60527, USA

***Corresponding Author:** Win L Chiou, Chiou Consulting Inc, 8552 Johnston Road, Burr Ridge, Illinois, 60527, USA; E-mail: win@chioucounseling.com

Received Date: 09-03-2021; **Accepted Date:** 01-04-2021; **Published Date:** 08-04-2021

Copyright © 2021 by Chiou WL. All rights reserved. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Currently, the U.S. Food and Drug Administration's policy on sunscreens assumes that exposure to sunlight, mainly ultraviolet (UV) rays, damages the skin, the damage is cumulative and any sun exposure should be avoided or minimized. Sunscreens are approved to help prevent early skin aging, sunburn and skin cancer. This work presents provocative perspectives on some issues related to use of sunscreens shown below. Skin aging is predominantly caused by the intrinsic nutritional factor. Sunlight damage is unlikely cumulative due to extremely efficient self-repair/self-defense mechanisms of the body. Regular exposure to moderate non-burning sunlight increases immunity and protects against sunburn, melanoma, COVID-19 mortality, high blood pressure, etc. White and non-white populations may need sun protection only when the UV Index is $\geq 5-6$ and $\geq 7-8$, respectively. Modelling analyses suggest that sunscreens with SPF 8 and 2-6 may be adequate for whites and non-whites, respectively. There are inherent risks of sunburn and melanoma when sunscreen is used intermittently mainly due to unavoidable SPF-independent missing applications and may serve as a false security; this may mainly account for exponential increases in melanoma incidences in recent decades; thus, the more intermittent use of sunscreen, the more incidence of sunburn and skin cancer. 10am-2pm sunlight avoidance guidelines may need modification. Extensive human studies suggest that intense UVB from sunburn, not UVA or sub-erythral UVB, may cause melanoma; this hypothesis is supported by mouse and fish models. Without exposure to burning sunlight, we

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. J Dermatol Res. 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

may not need to use broad-spectrum sunscreens or UVB-based sunscreens in our daily lives. Potential implications of this work may be immense in terms of saving lives and improving health of consumers, reducing costs associated with marketing and use of sunscreens, as well as reducing environmental pollution.

Keywords

Food and Drug Administration; Sunscreen Policy; Sunlight Benefits; Aging/Anti-Aging Theory; Enhanced Immunity; Melanoma Incidence/Etiology; COVID-19 Mortality; Sunscreen Modelling; UV Index

Introduction

In the United States broad-spectrum sunscreens with a Sun Protection Factor (SPF) equal to or greater than 15 or 30 are currently marketed as over-the-counter drugs to help prevent early skin aging, sunburn and skin cancer. Their efficacy and safety have been generally regarded as facts [1-3]. It is commonly accepted that exposure to sunlight which has been regarded as a carcinogen damages the skin and the damage is cumulative [1,2,4,5]. Hence, the current FDA's policy recommends zero tolerance toward sunlight exposure. In fact, infants under six months are advised to avoid sun exposure even on cloudy days [1,5]. It also adopts a one-size-fits-all product for every user in terms of SPF [5]. However, it appears that no theoretical or clinical rationale for this policy has been published to date. This is relevant in view of well-known marked differences in skin sensitivity toward sunlight exposure between whites and non-whites [6-8]. In the U.S., the non-whites presently account for about 40% of the population and may become the majority in 2050 [9,10]. Also, until recently no pharmacodynamic modelling studies regarding dosing evaluation have ever been published, although this kind of study has been generally required for new drug development and approval [11]. In this regard, some organic sunscreens were unexpectedly found to be significantly absorbed into the body, triggering the requirement for safety studies [12].

The main goal of this article is to present provocative perspectives on some issues related to the use of sunscreens based on an updated review of the literature with the hope that it may stimulate further discussions and studies. The major topics to be covered are described below under each heading.

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. J Dermatol Res. 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

New Alternative Skin Aging/Anti-Aging Theory

Skin aging is caused by the intrinsic chronological factor and the extrinsic factor due to exposure to sunlight and environmental pollutants and possibly smoking [13,14]. Aging due to sunlight exposure, i.e., photo aging, may account for up to 80% or 90% of total skin aging [13,14].

Numerous studies have elegantly demonstrated that UV radiation can simultaneously reduce collagen synthesis and increase collagen degradation thereby resulting in a reduction of skin collagen with age in an accelerated manner [1,2,15-19]. However, this concept seems inconsistent with a widely cited pioneering study in 1975 demonstrating that total collagen contents from skin exposed or not exposed to sunlight in Caucasian whites over a wide range of age were practically the same [20]. Furthermore, the aging kinetics of collagen surprisingly followed a first- order kinetics, i.e., logarithmic linear decline, without showing the expected accelerated aging [21]. Results were similar for the parallel exponential aging kinetics of superficial microvascular density data from sites exposed or not exposed to sunlight in humans [21,22]. These findings suggest that skin aging in ordinary people is predominantly due to the intrinsic factor [21,23].

A new cardiac index/nutrition aging theory was recently proposed to explain the aging kinetics of the whole body and its various organs or tissues [21]. The cardiac index, postulated as a whole body's aging marker, was found unexpectedly for the first time to decrease exponentially with a mean half-life of 82 years based on a re-analysis of previously reported data [21,24]. Obviously, a decreasing cardiac index can result in reduced delivery of nutrients to various organs and tissues including skin through vascular capillaries that are required to maintain their normal function and vitality [25]. Interestingly, this theory is consistent with two recent reports on anti-aging and tissue regeneration:

1. The topical use of propylene glycol, an FDA-approved calorie-generating nutrient, could quickly promote growth of tissues such as hair, nail, skin and gums apparently through the rejuvenation of each tissue's stem/progenitor cells [26]
2. The rejuvenation of hearts in old rodents could quickly produce unexpected systemic effects, reverse the aging process of hearts (including removal of scar tissue and repair of congestion), skin, hair, kidney and muscle and potentially increase the life span of rodents or humans [11,21,27]

Exposure to excessive (UV Index, skin-sensitivity and time-length dependent) sunlight can obviously accelerate intrinsically induced aging symptoms such as age spots, wrinkles and sagging. In this new aging theory, formation of age or dark spots is attributed to born or acquired deficiencies in superficial microcirculation and formation of fine lines and wrinkles may result from a defense mechanism that could reduce the effective skin surface area to

minimize its water loss [21]. As expected, age spots can also occur in areas usually not exposed to sunlight such as thighs and trunks and increase in number and size with age [21]. Based on the conventional aging theory, facial wrinkles and age spots are formed from exposure to sunlight. However, this theory does not seem statistically reasonable because the whole face, not an isolated area, is virtually uniformly exposed to sunlight all the time [21].

Benefits of Non-Burning Sunlight: Enhanced Immunity and Reduction of Sunburn, Melanoma and Mortality from COVID-19

This subject has been well reviewed and only a few points will be highlighted. Although exposure to intense or burning sunlight is well known to be immunosuppressive, moderate sunlight may increase immune function and this may be partly related to enhanced synthesis of vitamin D, which has numerous potential health benefits [28-36]. Sunlight can increase production or release of nitric oxide that may dilate blood vessels, increase blood flow to tissues and enhance immunity, as well as to reduce blood pressure, heart attack and stroke; nitric oxide can be induced by solar UVA and UVB rays [29-32,37,38]. It was stated that the risk of sunlight avoidance is equivalent to smoking in reducing life expectancy [39]. Obviously, chronic exposure to wind or heat of any sources which can increase loss of water from the skin may also enhance the aging process.

Instead of causing sunburn and skin cancer as expected from the current policy, regular exposure to non-burning sunlight may ironically be protective against sunburn and melanoma. For example, outdoor professional workers had less incidences of melanoma than indoor workers in countries with a temperate climate [29-32,40]. In the state of Washington, regular exposure to sunlight was protective against melanoma in children with sun-tanned skin [41]. In Germany, outdoor activities in the absence of sunburn during childhood were protective against melanoma [42]. In a British study, spending weekends sunbathing at a beach (UV Index < 6 or 7) without wearing sunscreen was protective against melanoma [43]. Surprisingly, melanoma incidences were inversely proportional to a high continuous pattern of sun exposure from a meta-analysis [44]. The protection of non-burning sunlight against sunburn may be primarily attributed to increased melanin secretion that serves as a natural sunscreen; protection against melanoma may be attributed to reduced sunburn incidence and enhanced immunity or repair mechanism against DNA mutation. Interestingly, sunlight exposure could also increase the survival rate in melanoma patients [29,31,45].

Apparently after realizing the potential benefits of sunlight, Australia in 2014 and New Zealand in 2019 started adopting the World Health Organization guidance recommending their residents to enjoy sunshine when the UV Index is below 3 and to seek sun protection only when the UV Index is ≥ 3 [5,46-48]. These two countries are known to have the highest melanoma

incidences in the world. In view of the above, in the U. S. we may need sun protection only when the UV Index is ≥ 5 or 6 for whites and ≥ 7 or 8 (even 9 for short periods) for non-whites [5].

During the current COVID-19 pandemic, the potential role of sun exposure has received much attention. For example, in a study involving 88 countries their mortality rates were positively correlated with their distance from the equator [49]. In Iceland (mean peak UV Index being about 1.5) and Singapore (mean peak UV Index being about 11.8) where both countries had similar populations with similar GDPs per person in 2019 (about 67,000 vs 66,000 USD) and similar infected cases as of May 1, 2020, the mortality rate in Iceland was, however, surprisingly 79 times higher compared to Singapore [49]. In another study, the COVID-19 mortality rates in continental metropolitan France were negatively correlated with their average annual sunlight hours [50]. In the U.S., the mortality rates of various states were also generally higher with higher latitudes [51]. Despite the limitation of potential numerous confounding factors, the above examples, especially between Singapore and Iceland, seem to strongly support the role of sunlight in increasing the immunity which can suppress and kill the COVID-19 virus. Furthermore, a preliminary review showed that globally COVID-19 fatality rates paralleled vitamin D deficiency rates which may also be related to sun exposure [52]. Hence, non-burning sunlight exposure and blood measurements and oral supplements of vitamin D may be considered as part of a defence strategy against COVID-19.

In various states in the U.S., COVID-19 mortality rates were zero or negligible in young people under age 16-20 [53]. This interesting phenomenon is generally consistent with their highest cardiac indexes and highest immune function compared to other age groups [21,54].

Skin Cancer: A Rare Disease in Non-Whites

Skin cancer has been conveyed to the public as the most commonly occurring cancer in the U.S. However, this can only be applied to light-skinned Caucasians [6,7]. For non-whites such as blacks, Hispanics and Asians in the U.S. and many other countries, skin cancer is a rare disease and generally has no correlation with sun exposure [6,7,55,56]. Interestingly, their melanomas often occur at sites usually not exposed to sunlight such as the soles of feet, palms of hands, beneath toenails, groin areas and oral gums [6,7,55,56]. Early detection is emphasized for successful treatment whereas late detection is often associated with higher morbidity and mortality (6,7,55,56). Therefore, one may need to have a different health message for different constituents.

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. J Dermatol Res. 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

Cumulative Damage vs Extremely Efficient Self-Repair and Self-Defence Mechanisms: Implications

Skin damage due to sunlight may not cumulate [5,21]. For example, erythema may heal itself in days and a more serious blistering sunburn may be healed with minimum intervention within weeks [5,21,57]. Generally, mutation damage to DNA can be virtually completely repaired or removed through apoptosis within weeks or months and normal cells and mutated cells in our body are constantly maintained in a homeostatic state [5,32,58,59]. Therefore, our body has extremely efficient self-defense and self-repair mechanisms that may account for insignificant adverse effects of sunlight exposure on skin aging as discussed above and for the age-independent incidences of sunburn, basal cell carcinoma, squamous cell carcinoma and melanoma [5,21,29].

Interestingly, it has been stated that melanoma is a rare disease because only a very tiny fraction of mutant DNAs could escape the repair mechanism and develop into melanoma [44]. This tiny fraction may represent some of those mutant DNAs induced probably only by the intense UVB from sunburn, not from the non-burning UVA or UVB as commonly assumed (more discussion below). Thus, there may exist a critical level of DNA damage that is needed for developing into melanoma.

Peak Noon Hours Concept and UV Index Monitoring

Presently, the public is advised to avoid sun exposure between 10 am and 2 pm due to peak sunlight intensity (1, 2). Such a recommendation may need to be revised. This is because the peak UV Index may vary greatly with the location and time of year. For example, in winter and early spring most of the U.S. has a UV Index below 3 to 5, thus there may be no need to seek sun protection for most of us. In fall and late spring most UV Indexes are probably below 6 to 7, thus there may be no need to exercise this precaution for non-whites. It seems that the importance and regular monitoring of UV Index should be properly conveyed to our general public.

Pharmacodynamics modelling: New Efficacy Evaluation and SPF Recommendations

A new Skin UV Index (SUI) was introduced recently [11]. It refers to sunlight heat intensity on the skin in terms of UV Index after sunscreen application. It was estimated that SUIs were all below 3 when sunscreens with an SPF ranging from 4 to 100 were applied according to the

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. J Dermatol Res. 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

label instruction. i.e., 2 mg/cm² and the UV Index was 10 [11]. This suggests that for Caucasian white's use of these sunscreens may be effective against sunburn and melanoma and any reported inefficacies of sunscreens may be mainly attributed to various degrees of non-compliance with the label instruction [11]. SUIs were also estimated when only 50% and 25% of labelled amounts were applied as they may often occur in the real world [11]. The present modelling analysis may thus provide a novel rationale as to why regular use of an SPF 8 sunscreen was effective against melanoma in Australia [60]. It was suggested that in the U.S., SPF 8 sunscreen may be adequate for whites and SPF 2-6 sunscreens may be sufficient for non-whites [11]. The above findings seem important in view of recent discussions on the use of sunscreens with SPF's higher than 50 or 100 for achieving greater efficacy [1,11,61-64].

It is noteworthy that the well-known pivotal study of AC Green, et al., in 2011 that has been universally used as a gold standard for setting the current minimum SPF 15 requirement has been reported to have serious shortcomings in the study design, data interpretation and application [3,11]. Therefore, the justification of its use as a regulatory supporting evidence is now in doubt [3,11].

Risks of Sunburn and Melanoma from Sunscreen

In recent decades, there has been an exponential increase in melanoma incidences in many countries [65]. An intriguing question arises as to whether increased use of sunscreen has played a role and this issue has been extensively reviewed [11]. Because in the U.S., sunscreens have been primarily used intermittently for sunbathing at beaches or vacation at high-altitude snow resorts under intense sunlight, an unavoidable, SPF-independent missing application of sunscreen on the body may result in sunburn on these areas that may later develop into melanoma [11]. Conceivably, missing application may most often occur on the trunk of the body or lower legs. Therefore, it is not surprising that melanoma has been indeed found to occur most often in these two areas [57,66]. It is noteworthy that after one week of sun vacation and wearing a broad-spectrum SPF 30 sunscreen at a beach in Spain, all 25 Danish vacationers were found to experience sunburn [67]. The author of this study referred to such a sun vacation as a sunburn vacation. It is important to emphasize that even 1% or 0.1 % of missing application may result in that missed area to develop into sunburn. The missed areas may also include face, neck, eyelids, ears and lips. It was hypothesized that the increasing use of sunscreen may primarily account for the reported exponential increase in melanoma in many Western countries over the last few decades mainly due to their intermittent use [11,65]. Thus, the use of sunscreen can protect or harm the skin. The potential risks of sunburn and melanoma or perhaps other skin cancers from use of sunscreen during intermittent exposure may probably need to be reflected on product labels [11].

Alternative Hypothesis on Etiology of Melanoma in Caucasian Whites: Implications

Solar UV rays reaching the Earth's surface contain UVA and UVB [1,2]. The UVA is much more abundant and can penetrate deeper into the skin than UVB to cause more damage to DNA, collagen, etc. [1,2,32,66,68,69]. In the last two decades, the major cause of skin aging and melanoma has been largely or predominantly attributed to UVA radiation [1,2,32,66,68,69]. As discussed above, regular exposure to non-burning sunlight may not have a significant adverse effect on skin aging. Could this kind of phenomenon also occur in the etiology of skin cancers, especially melanoma? A review of literature suggests that this may also occur as discussed below.

Since chronic exposure to non-burning sunlight has been repeatedly shown to be protective against melanoma, therefore, it seems unlikely that the non-burning solar UVA or UVB may cause melanoma [5,29-31,40-44]. Because sunburn is a biomarker for melanoma (70) and five or more serious sunburns in a lifetime could double the melanoma risk and an increased risk of melanoma with increasing number of sunburns was observed during all life periods, hence, the UVB-dominated burning solar ray may be hypothesized to predominantly cause the melanoma in humans [71,72]. This is consistent with the above discussion that use of sunscreens may account for increasing incidences of melanoma in recent decades due to intermittent exposure [11]. Also, melanoma incidences were reported to be proportional to incidences of intermittent high-UV-intensity sun exposures [44]. Thus, the above discussion suggests that there may exist a critical level of DNA damage that is needed for it to develop into melanoma. Obviously damage from severe sunburn may be more critical than the simple superficial erythema [66]. The present hypothesis seems to be supported by animal studies. For example, only the erythematous dose of UVB, not UVA or repeated sub-erythematous UVB doses, could induce melanoma in mice and only the erythematous dose of UVB could induce melanoma development in neonatal mice [73,74]. Results from two other mouse studies also seem to support the present theory [75,76]. Furthermore, only UVB, not UVA, could induce melanoma in a fish model [66,75-77].

The above discussion suggests that for the general public we may need to reconsider the need of using broad-spectrum sunscreens or even traditional UVB-based sunscreens for protecting us from photo aging and melanoma or perhaps also other types of skin cancers in our daily lives. We may need to emphasize that the best way to prevent skin cancer is to prevent sunburn by avoiding exposure to excessive burning sunlight, by seeking shade or by wearing protective clothing and hats [1,2,5,40]. Although visible and infrared light wavelengths can generate 50% of all free radicals in the skin and cause similar collagen damage, there may be no need to protect against them [78]. Interestingly, moisturizers containing hyaluronic acid, glycerin or

propylene glycol have unique anti-skin-aging properties, a 50% glycerin solution was tested to have an SPF 2 and used to protect against sunburn [11,79-81-83].

Concluding Statements: Implications

Obviously, the potential implications of the above perspectives, if implemented, may be quite immense in terms of saving lives and treatment costs of patients suffering from skin cancer due to their intermittent use of sunscreens, savings of costs associated with the use and marketing of sunscreens, reduction of environmental pollutions from sunscreens and improvements in mental and physical health, as well as in life expectancy of people from regular exposure to moderate non-burning sunlight [32]. Clearly, unnecessary use of sunscreen in our daily lives may deprive us from the potential benefits of sunlight [21]. It is my view that if needed, inorganic sunscreens such as zinc oxide and titanium dioxide should be our preferred choices because of their non-absorbability and inertness and use of organic sunscreens should be strongly discouraged mainly because of their potential long-term adverse effects on users and the environment [1,12,82,83].

Conflict of Interest

Winlind Skincare LLC: As president of this practically one-person (WLC, age 82), residence-based family company with negligible activities.

Source of Funding:

Self-funding

References

1. Statement from FDA Commissioner Scott Gottlieb, M.D. on new FDA actions to keep consumers safe from the harmful effects of sun exposure and ensure the long-term safety and benefits of sunscreens. [Last accessed on March 25, 2021] <https://www.fda.gov/news-events/press-announcements/statement-fda-commissioner-scott-gottlieb-md-new-fda-actions-keep-consumers-safe-harmful-effects-sun>
2. Sunscreen FAQs, American Academy of Dermatology Association. [Last accessed on March 25, 2021] www.aad.org/media/stats-sunscreen
3. Waldman RA, Grant-Kels JM. The role of sunscreen in the prevention of cutaneous melanoma and nonmelanoma skin cancer. *J Am Acad Dermatol*. 2019;80:574-6.
4. Nole G, Johnson AW. An analysis of cumulative lifetime solar ultraviolet radiation exposure and the benefits of daily sun protection. *Dermatol Ther*. 2004;17(1):57-62.
5. Chiou WL. We may need to reconsider when to apply sunscreen in our daily life. *Ann Dermatol Res*. 2019;3:7-10.

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. *J Dermatol Res*. 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

6. Bradford PT. Skin cancer in skin of color. *Dermatol Nurs*. 2009;21:170-7.
7. Wang Y, Zhao Y, Ma S. Racial differences in six major subtypes of melanoma: descriptive epidemiology. *BMC Cancer*. 2016;16:691.
8. Morris KL, Perma FM. Decision tree model vs traditional measures to identify patterns of sun-protective behaviors and sun sensitivity associated with sunscreen. *JAMA Dermatol*. 2018;154:897.
9. U.S. Census Bureau QuickFacts: United States. [Last accessed on March 25, 2021] <https://www.census.gov/quickfacts/fact/table/US/PST045219>
10. Gloster Jr HM, Neal K. Skin cancer in skin of color. *J Am Acad Dermatol*. 2006;55:741–60.
11. Chiou WL. Pharmacodynamic modeling of sunscreens: new efficacy evaluation, risks of sunburn and melanoma and very low to very high sun protection factor. *J Dermatol Res Ther*. 2020;1:20-9.
12. Matta MK, Zusterzeel R, Pilli NR, Patel V. Effect of sunscreen application under maximal use conditions on plasma concentration of sunscreen active ingredients: A randomized clinical trial. *JAMA*. 2019;321:2082-91.
13. Taylor CR, Stern RS, Leyden JJ, Gilchrest BA. Photoaging/photodamage and photoprotection. *J Amer Acad Dermatol*. 1990;22:1-15.
14. Kammeyer A, Luiten RM. Oxidation events and skin aging. *Ageing Res Rev*. 2015;21:16-29.
15. Fisher GJ, Datta SC, Talwar HS, Wang ZQ, et al. Molecular basis of sun-induced premature skin ageing and retinoid antagonism. *Nature*. 1996;379:335-9.
16. Jung M, Kelly K, McCullough J. Skin aging: pathogenesis, prevention and treatment, In: Rattan S, Kassem M. Prevention and treatment of age-related diseases. 2006;175-92.
17. Jin K. Modern biological theories of aging. *Aging Dis*. 2010;1:72-4.
18. Chung J, Seo JY, Choi HR, Lee MK. Modulation of skin collagen metabolism in aged and photoaged human skin in-vivo. *J Invest Dermatol*. 2001;117:1218-24.
19. Butov A, Volkov N, Anisimov V, Sehl ME, et al. A model of accelerated aging induced by 5-bromodeoxyuridine. *Biogerontol*. 2002;3:175-82.
20. Shuster S, Black M, McVitie E. The influence of age and sex on skin thickness, skin collagen and density. *Br J Dermatol*. 1975;93:639-43.
21. Chiou WL. Aging kinetics of human hearts and skin: New aging theories and implications in the use of sunscreens. *Dermatol Arch*. 2017;1:1-5.
22. Li L, Mac-Mary S, Sainthillier JM, Nouveau S, et al. Age-related changes of the cutaneous microcirculation *in-vivo*. *Gerontol*. 2006;52:142-53.
23. Shuster S. Osteoporosis, like skin ageing, is caused by collagen loss which is reversible. *J R Soc Med*. 2020;113:158-60.
24. Guyton A. Textbook of medical physiology (4th Ed). Saunders, USA. 1971:311.
25. Chiou WL. The phenomenon and rationale of marked dependence of drug concentration on blood sampling site (Part I). *Clin Pharmacokinet*. 1989;17:175-99.
26. Chiou WL. Compositions and methods for tissue regeneration. United States patent no. 10456366 issued on October 29, 2019.
27. Grignorian-Sharmagian L, Liu W, Fereydooni S, Middleton R. Cardiac and systemic rejuvenation after cardiosphere-derived cell therapy in senescent rats. *Eur Heart J*. 2017;38:2957-67.
28. Wright F, Weller RB. Risks and benefits of UV radiation in older people: More of a friend than a foe? *Maturitas*. 2015;81:425-31.
29. Hoel DG, Berwick M, de Gruijl F, Holick M. The risks and benefits of sun exposure 2016. *Dermatoendocrinol*. 2016;8:e1248325.
30. Grant WB. A review of the evidence supporting the vitamin D-cancer prevention hypothesis in 2017. *Anticancer Res*. 2018;38:1121-36.
31. Rhee H, de Vries E, Coomans CP, van de Velde P. Sunlight: For better or for worse? A review of positive and negative effects of sun exposure. *Cancer Res Frontiers*. 2016;2:156-83.
32. Powers JM, Murphy JEJ. Sunlight radiation as a villain and hero: 60 years of illuminating research. *Int J Rad Bio*. 2019;95:1043-9.

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. *J Dermatol Res*. 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

33. Narbutt J, Lesiak A, Sysa-Jedrzejowska A, Wozniacka A. Repeated low-dose ultraviolet (UV) B exposures of humans induce limited photoprotection against the immune effects of erythematous UVB radiation. *Br J Dermatol.* 2007;156:539-47.
34. Aranow C. Vitamin D and the Immune System. *J Invest Med.* 2011;59:881-6.
35. González Maglio DH, Paz ML, Leoni J. Sunlight effects on immune system: Is there something else in addition to UV-Induced immunosuppression? *Biomed Res Int.* 2016;1934518.
36. Cela EM, Gonzalez CD, Friedrich A, Ledo C. Daily very low UV dose exposure enhances adaptive immunity, compared with a single high-dose exposure. Consequences for the control of a skin infection. *Immunol.* 2018;154:510-21
37. Moncada S, Higgs EA. The discovery of nitric oxide and its role in vascular biology. *Br J Pharmacol.* 2006;147(1):S193-201.
38. Weller RB, Wang Y, He J, Maddux FW. Does incident solar radiation lower blood pressure? *J Am Heart Ass.* 2020;9:e13837.
39. Lindqvist PG, Epstein E, Landin-Olsson M, Ingvar C. Avoidance of sun exposure is a risk factor for all- cause mortality: results from the melanoma in southern Sweden cohort. *J Intern Med.* 2014;276:77-86.
40. Planta MB. Sunscreen and melanoma: Is our prevention message correct? *J Am Board Fam Med.* 2011;24:735-9.
41. White E, Kirkpatrick CS, Lee JA. Case-control study of malignant melanoma in Washington State. I. Constitutional factors and sun exposure. *Am J Epidemiol.* 1994;139:857-68.
42. Kaskel P, Sander S, Kron M, Kind P. Outdoor activities in childhood: a protective factor for cutaneous melanoma? Results of a case-control study in 271 matched pairs. *Br J Dermatol.* 2001;145:602-9.
43. Newton-Bishop JA, Chang YM, Elliott F, Chan M. Relationship between sun exposure and melanoma risk for tumors in different body sites in a large case-control study in a temperate climate. *Eur J Cancer.* 2011;47:732-41.
44. Gandini S, Sera F, Cattaruzza MS, Pasquini P. Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer.* 2005;41:45-60.
45. Berwick M, Armstrong BK, Ben-Porat L, Fine J. Sun exposure and mortality from melanoma. *J National Cancer Inst.* 2005;97:195-9.
46. Siegel J. America is getting the science of sun exposure wrong. Website for Cancer Council Australia. 2014. [Last accessed on March 25, 2021] <http://nautil.us/issue/14/mutation/america-is-getting-the-science-of-sun-exposure-wrong> or <https://www.cancer.org.au/cancer-information/causes-and-prevention/sun-safety/10-myths-about-sun-protection>
47. Whiteman DC, Neale RE, Aitken J, Gordon L. When to apply sunscreen: a consensus statement for Australia and New Zealand. *Aust N Z J Public Health.* 2019;43:171-5.
48. WHO UV Index. [Last accessed on March 25, 2021] <https://www.who.int/uv/publications/en/UVIGuide.pdf>
49. Whittemore PB. COVID-19 fatalities, latitude, sunlight and vitamin D. *Amer J Infection Control.* 2020;48:1042-4.
50. Lansiaux É, Pébay PP, Picard JL, Forget J. COVID-19 and vitamin D: Disease mortality negatively correlates with sunlight exposure. *Spat Spatiotemporal Epidemiol.* 2020;35:100362.
51. Marik PE, Kory P, Varon J. Does vitamin D status impact mortality from SARS-CoV-2 infection? *Med in Drug Disco.* 2020;6:100041.
52. Benskin LL. A basic review of the preliminary evidence that COVID-19 risk and severity is increased in vitamin D deficiency. *Front Public Health.* 2020;8:513.
53. Children and COVID-19: state-level data report, American Academy of Pediatrics. [Last accessed on March 25, 2021] <https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-covid-19-state-level-data-report>
54. Hirokawa K, Utsuyama M, Kasai M, Kurashima C. Aging and Immunity. *Acta Pathol Jpn.* 1992;42:537-48.
55. Chang JWC. Cutaneous melanoma: Taiwan experience and literature review. *Chang Gung Med J.* 2020;33:602-12.

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. *J Dermatol Res.* 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

56. Fujisawa Y, Yoshikawa S, Minagawa A, Takenouchi T. Clinical and histopathological characteristics and survival analysis of 4594 Japanese patients with melanoma. *Cancer Med.* 2019;8:2146-56.
57. Habif HP. *Clinical Dermatology*, 5th Ed. 2010. MOSBY, Elsevier. 747 and 863.
58. Anna B, Blazej Z, Jacqueline G, Andrew CJ. Mechanism of UV-related carcinogenesis and its contribution to nevi/melanoma. *Expert Rev Dermatol.* 2007;2:451-69.
59. Scott TL, Rangaswamy S, Wicker CA, Izumi T. Repair of oxidative DNA damage and cancer: Recent progress in DNA base excision repair. *Antioxid Redox Signal.* 2014;20:708-26.
60. Watts CG, Drummond M, Goumas C, Schmid H, et al. Sunscreen use and melanoma risk among young Australian adults. *JAMA Dermatol.* 2018;154:1001-9.
61. Linden KG. Sunscreen sun protection factor (SPF), Is higher better? *J Am Acad Dermatol.* 2018;78:911-2.
62. Williams JD, Maitra P, Atillasoy E, Wu MM. SPF 100+ sunscreen is more protective against sunburn than SPF 50+ in actual use: Results of a randomized double-blind, split-face, natural sunlight exposure clinical trial. *J Am Acad Dermatol.* 2018;78:902-10.
63. Kohli I, Nicholson CL, Williams JD, Lyons AB. Greater efficacy of SPF 100+ sunscreen compared to SPF 50+ in sunburn prevention during 5 consecutive days of sunlight exposure: A randomized double-blind clinical trial. *J Am Acad Dermatol.* 2020;82:869-77.
64. Ou-Yang H, Jiang LI, Meyer K, Wang SQ. Sun protection by beach umbrella vs sunscreen with a high sun protection factor: A randomized clinical trial. *JAMA Dermatol.* 2017;153:304-8.
65. Whiteman DC, Green AC, Olsen CM. The growing burden of invasive melanoma: projections of incidences rates and numbers of new cases in six susceptible populations through 2031. *J Invest Dermatol.* 2016;136:1161-71.
66. Autier P, Doré JF. Ultraviolet radiation and cutaneous melanoma: a historical perspective. *Melanoma Res.* 2020;30:113-25.
67. Petersen B, Thieden E, Philipsen PA, Heydenreich J. A sun holiday is a sunburn holiday. *Photodermatol Photoimmunol Photomed.* 2013;29:221-4.
68. Pudroma X, Duoqi G, Grigalavicius M, Jie D, Molecular mechanisms of UVA-induced melanoma. *J Environ Pathol Toxicol Oncol.* 2017;2(13):217-28.
69. Leonardi GC, Falzone L, Salemi R, Zanghi A, Spandidos DA. Cutaneous melanoma: From pathogenesis to therapy. *Int J Oncol.* 2018;52:1071-80.
70. Holman DM, Ding H, Guy GP, Watson M. Prevalence of sun protection use and sunburn and association of demographic and behavioral characteristics with sunburn among US adults. *JAMA Dermatol.* 2018;154:561-8.
71. Pfahlberg A, Kölmel KF, Gefeller O, Febim Study Group. Timing of excessive ultraviolet radiation and melanoma: epidemiology does not support the existence of a critical period of high susceptibility to solar ultraviolet radiation-induced melanoma. *Br J Dermatol.* 2001;144:471-5.
72. Dennis LK, Vanbeek MJ, Beane Freeman LE, Smith BJ. Sunburns and risk of cutaneous melanoma: does age matter? A comprehensive meta-analysis. *Ann Epidemiol.* 2008;18:614-27.
73. Van Schanke A, Jongsma MJ, Bisschop R, van Venrooij GMCAL. Single UVB overexposure stimulates melanocyte proliferation in murine skin, in contrast to fractionated or UVA-1 exposure. *J Invest Dermatol.* 2005;124:241-7.
74. Noonan FP, Recio JA, Takayama H, Duray P. Neonatal sunburn and melanoma in mice. *Nature.* 2001;413:271-2.
75. De Fabo EC, Noonan FP, Fears T, Merlino G. Ultraviolet B but not ultraviolet A radiation initiates melanoma. *Cancer Res.* 2004;64:6372-6.
76. Day CP, Marchalik R, Merlino G, Michael H. Mouse models of UV-induced melanoma: genetics, pathology and clinical relevance. *Lab Invest.* 2017;97:698-705.
77. Mitchell DL, Fernandez AA, Nairn RS, Garcia R. Ultraviolet A does not induce melanomas in a Xiphophorus hybrid fish model. *Proc Natl Acad Sci USA.* 2010;107:9329-34.
78. Zastrow L, Lademann J. Light - instead of UV protection: New requirements for skin cancer Prevention. *Anticancer Res.* 2016;36:1389-93.

Chiou WL | Volume 2; Issue 1 (2021) | JDR-2(1)-019 | Opinion Article

Citation: Chiou WL. Perspectives on the FDA's Sunscreen Policy: New Aging Theory, Sunlight Benefits, Enhanced Immunity, COVID-19 Mortality, Modelling Analyses, Melanoma Risks and Etiology. *J Dermatol Res.* 2021;2(1):1-13.

DOI: <http://dx.doi.org/10.46889/JDR.2021.2104>

79. Jegasothy SM, Zabolotniaia V, Bielfeldt S. Efficacy of a new topical nano-hyaluronic acid in humans. *J Clin Aesthet Dermatol*. 2014;7:27-9.
80. Chiou WL, Chiou LL. Aqueous compositions for facial cosmetics. United States Patent No. 6616923 issued on September 9, 2003.
81. Chiou WL. Compositions and methods for tissue regeneration. United States Patent No. 10456366 issued on October 9, 2019.
82. DiNardo JC, Downs CA. Why evaluate the sunscreen active oxybenzone (benzophenone-3) for carcinogenicity and reproductive toxicology or consider it unsafe for human use? *Dermatol Res*. 2019;1:1-3.
83. Downs CA, DiNardo JC, Stien D. 83 Benzophenone accumulates over time from the degradation of octocrylene in commercial sunscreen products. *Chem Res Toxicol*. 2021.