BJPS

BRAZILIAN JOURNAL OF PHARMACEUTICAL SCIENCES

*Brazilian
journal of
pharmaceutical
ciences*

Faculdade de Ciências Farmacêuticas
Universidade de São Paulo

V International Workshop on Rational Use of Medicines

III Workknowledge of Evidence-Informed Policy

III Symposium ISPE BrazIntRIG

I Symposium of ISPE Brazilian Student Chapters

Cataloguing in Publication Data (CIP)

-
- B827 Brazilian Journal of Pharmaceutical Science [recurso eletrônico] / USP, Faculdade de Ciências Farmacêuticas. – Vol. 1, n. 1 – Campina Grande, PB: Realize editora, 2021.
Edição especial contendo os Anais do V International Workshop on Rational Use of Medicines.
Texto em inglês.
DOI 10.46943/V.IWRUM.2021.01
ISSN 2175-9790 (versão online).
1. Ciências farmacêuticas. 2. Farmacoterapia. 3. Cuidado farmacêutico. 4. Prescrição de medicamentos. 5. Uso de medicamentos. I. Título.
21 ed. CDD 615-1
-

Elaborated by Giulianne M. Pereira

CRB 15/714

About the Publication:

Graphic Project and Publishing: Jefferson Ricardo Lima Araujo Nunes

REALIZE EVENTOS CIENTÍFICOS & EDITORA LTDA.

Rua: Aristίδes Lobo, 331 - São José - Campina Grande-PB | CEP: 58400-384

E-mail: contato@portalrealize.com.br | Telefone: (83) 3322-3222

V International Workshop on Rational Use of Medicines

III Workknowledge of Evidence-Informed Policy

III Symposium ISPE BrazIntRIG

I Symposium of ISPE Brazilian Student Chapters

ABSTRACTS

March 1-5, 2021

Sorocaba [Online] – Brazil

COORDINATION

Luciane Cruz Lopes - *Uniso*

ORGANIZATION COMMITTEE

Fabiane Raquel Motter - *Uniso*

Maíra Ramos Alves - *Uniso*

Cristiane de Cassia Bergamaschi - *Uniso*

Monique Elseviers - *University of Antwerp*

SCIENTIFIC COMMITTEE

Airton Tetelbom Stein - *UFMG*

Claudia Garcia Serpa Osorio de Castro – *Fiocruz*

Daniela Oliveira de Melo - *Unifesp*

Daniele Ribeiro de Araujo - *UFABC*

Denise Grotto - *Uniso*

Fabiane Raquel Motter - *Uniso*

Gladys Marques - *Uniso*

Jorge Barreto Maia - *Fiocruz/Brasília*

Lisiane Freitas Leal – *McGill University*

Luciane Cruz Lopes - *Uniso*

Monique Elseviers - *Uantwerp*

Renata de Lima - *Uniso*

COORDINATION OF THE WORK GROUPS

Daniela Oliveira de Melo - *Unifesp*

Denise Grotto - *Uniso*

Fabiane Raquel Motter - *Uniso*

Jorge Barreto Maia - *Fiocruz*

Lisiane Freitas Leal - *UFRGS*

Renata de Lima - *Uniso*

Silvio Barberato Filho - *Uniso*

Tais Freire Galvão - *Unicamp*

PRESENTATION APPRAISERS

Cristiane de Cássia Bergamaschi - *Uniso*

Daniela Oliveira de Melo - *Unifesp*

Daniele Ribeiro de Araujo – *UFABC*

Fabiane Raquel Motter – *Uniso*

Izabela Fulone - *Uniso*

Jardel Côrrea de Oliveira - *SMS-SC*

Lisiane Freitas Leal – *McGill University*

Silvio Barberato Filho - *Uniso*

Sueli Miyuki Yamauti – *Unifesp*

PREFACE

Producing and sharing knowledge have been the main goals of the Graduate Program in Pharmaceutical Sciences of the University of Sorocaba. With a focus on the rational use of medicines, the importance of the event is justified by highlighting one of the main concerns worldwide, with an important impact on society, health systems, institutions, and communities.

The improvement of professional practices depends on the engagement of researchers, health professionals, managers, students, and others interested in improving health policies, programs, services, and actions. Developing and applying scientific methods in producing and using the best evidence is the path we have chosen.

Therefore, everyone was invited to discuss relevant topics in this field of knowledge, including Drug Utilization Research; Health Technology Assessment; Global Health Systems and Environment; and Innovation and Development of Health Technologies. Experts from several countries in describing, interactive teaching approaches, implementation science and policy, environment and pharmaceutical care joined us sharing their experiences and encouraging debate.

We are sure that the social distancing, essential in this period, did not represent an obstacle in making new connections, and effective and bright collaborations that are able to transform reality.

Luciane Cruz Lopes, ScD, MSc

*Graduate Program in Pharmaceutical Sciences
V Workshop of Rational Use of Medicine Coordination*

SUMMARY***TA 01: Drug Utilization Research/ Promotion of Rational Use of Medicine***

Frequency and severity of adverse drug reactions to medications prescribed for Alzheimer's disease in a Brazilian city: a cross-sectional study	12
Analysis of pharmacotherapy among residents who use multiple drugs in Santo André, São Paulo	13
Home care through the best at home program (MEMC) - a case report	14
Free access and use of medicines by the elderly population in the municipality of Campinas/SP: results of a population survey	15
Analysis of serious drug interactions in paracetamol poisoning in Brazil	16
Promotion of rational use of medicine through social media	17
Profile of antimicrobial dispensing in primary health care in the context of the COVID-19 pandemic	18
Monitoring actions for promoting the rational use of antimicrobials contained in the national action plan for the prevention and control of antimicrobial resistance 2018- 2022 (PAN-BR)	19
Analysis of the competence of written communication of prescriptions carried out by Brazilian pharmacy students in the last year of the graduation course	20
Impact of polypharmacy on the nutritional status of institutionalized elderly	21
Potential drug-drug interactions of clinical importance between antihypertensives: prevalence and risks for patients in primary healthcare	22
Rethinking HIV/AIDS prevention public policies for the elderly based on the discourses of elders living with the virus	23
Use of the wise application as a tool in the therapeutic conduct of the elderly with chronic pain	24
Indiscriminate use of ascorbic acid during the COVID-19 pandemic in Brazil	25
COVID-19 pandemic induces increase in clonazepam consumption in the state of Alagoas, Brazil, 2020	26

Eating habits and physical exercise in patients with type 1 Diabetes mellitus: is there a difference before and after pharmacotherapeutic follow-up?.....	27
Evaluation of medication use by public school teachers in Pelotas-RS	28
Association between anticholinergic burden of medications and emergency care in the elderly assisted in primary health care, Sorocaba, SP, Brazil.....	29
Self-medication in Pernambuco: a preliminary analysis of SINAN data from 2017 to 2019.....	30
Pharmaceutical care in critical patients and the impact on the use of financial resources with medicines	31
Risk of infections during treatment of juvenile idiopathic arthritis with adalimumab versus other indications.....	32
Safety in the disposal of medicines: perspectives on rational use	33
Insulin therapy among users of a primary health care unit.....	34
Evaluation of a control system for the prescription of restricted antibiotics in a large Brazilian hospital.....	35
Trend in the use of atypical antipsychotics in the Brazilian Public Health System: a cross-sectional study.....	36
Is home blood pressure monitoring an opportunity to deprescribing antihypertensive drugs?.....	37
Opioids in Brazilian federal hospitals.....	38
Falls among the elderly assisted in primary health care: prevalence and associated factors.....	39
The role of pharmaceutical intervention in the management of drug shortage in an epidemic context.....	40
Pharmaceutical Services in the perspective of the users of public pharmacies in a municipality of Rio de Janeiro state: gaps that can impact access.....	41
Use of medicines by elderly people in primary care of the Unified Health System (SUS) in Brazil.....	42
Drug therapeutic prescribed according to initial complaint and sign/symptom to users of the infanto-juvenile psychosocial care center	43

Pharmacists' role in patient safety: evaluation of dispensing errors in a regional hospital.....	44
Use of medicines in child mental health: a description	45
Psychotropic medication usage by children referred to centers for psychosocial child and adolescent care	46
Profile of use of ceftazidime/avibactam in ICU patients in a tertiary hospital.....	47
Investigation of anti-infectives adverse reactions in hospitalized children in a university institute in the city of Rio de Janeiro	48
Anticoagulant and antiplatelet therapy in older patients: a Brazilian consensus of potentially inappropriate prescribing indicators.....	49
Interventions for depression and anxiety in diabetes mellitus patients: overview of systematic reviews and meta-analyses.....	50

TA 02: Health Technology Assessment

Prevalence of <i>Enterococcus faecium</i> resistant in poultry farming: systematic review.....	51
Prevalence of <i>Enterococcus faecium</i> resistant in commercial cattle: systematic review.....	52
Efficacy and effectiveness of mobile health technologies to reduce alcohol or tobacco consumption during pregnancy: scoping review.....	53
Wearable devices designed to improve medication adherence: a systematic map.....	54
Efficacy and safety of repellents marketed in Brazil against bites from <i>Aedes aegypti</i> and <i>Aedes albopictus</i> : a systematic review.....	55
Antimicrobial resistance in periodontal infections (MICROBE-DENT): systematic review.....	56
Critical appraisal of clinical practice guidelines for the treatment of urinary incontinence: a systematic review.....	57
Development of a nursing care guideline for patients with external ventricular drainage.....	58

Prevalence of <i>Enterococcus faecium</i> resistant in commercial aquaculture: systematic review.....	59
The incorporation of biological medicines into the Brazilian Unified Health System (SUS), 2000-2019	60
Equity in health and COVID-19: bibliometric approach of impact publications..	61
Access to innovative biological medicines in Brazil: an analysis of FDA and ANVISA drug approvals.....	62
Health technology assessment study on effectiveness and safety of OK-432 (picibanil) in cervical lymphatic malformations.....	63
Expectation for the development of health technology assessment in Brazil.....	64
Risk of osteonecrosis in bisphosphonate users submitted to dental procedures: systematic review and meta-analysis	65
Safety and efficacy of treatment with zoledronate compared with pamidronate in children with osteogenesis imperfecta: rapid review.....	66

TA 03: Global Health System & Environment

Integration of dental attention to the multi-professional work process in the basic health unit of vila do engenho located in the municipality of Itacoatiara.....	67
Avoidable public expenditures with selected antineoplastic drugs for breast cancer in Brazil	68
Right to free health care and high-cost medications: brief notes.....	69
Saber-SUS: Evidence based health and recommendations for SUS	70
Analysis of financial viability in the participation of the municipality of Maringá in the Paraná health consortium.....	71
Profile of health judicialization involving drugs from specialized component of pharmaceutical assistance in the state of Pará, Brazil	72
Profile of administrative demands for off label use of mofetilla mycophenolate, in the state of Pará	73

TA 04: Innovation and Development of Health Technology

A comparative analysis of capillary blood glucose with deproteinized glucose oxidase in relevant clinical glucose concentrations in diabetic Wistar rats after beheading.....	74
Use of limonene as a therapeutic agent against <i>Escherichia coli</i> infections	75
Effects of <i>Lentinula edodes</i> cereal bar consumption on antioxidant profile: a double blind randomized clinical trial.....	76
Indicators for safe medication use: systematic review.....	77
Comparison of fungiculture and viniculture residues as biosorbents for lead contamination.....	78
Doxycycline safety through hematological, hepatic, and renal profile in chronic pre-clinical exposure.....	79
Biogenic synthesis and characterization of silver nanoparticles based on <i>Beauveria bassiana</i>	80
Analysis in laboratory-scale of cadmium removal in aqueous solution utilizing bacterial nanocellulose.....	81
Antimicrobial evaluation of a bioproduct composed of bacterial nanocellulose, fibroin and nisin.....	82
Phytol-based nanoemulsion: a promising anti-cancer action.....	83
Clinical trials with hormonally active herbal substances on the reproductive health of men and women: systematic literature review.....	84
Validation of the x-ray fluorescence technique for the use in pharmaceutical sciences.....	85
Anti-inflammatory <i>in vivo</i> evaluation by the air pouch model of the crude hydroalcoholic extract of <i>Myrcia pubipetala</i> (EBH-MP).....	86
Use of citrus fruit-derived essential oil as a bactericidal agent on <i>Escherichia coli</i> strains.....	87
<i>In vitro</i> antidiabetic activity of medicinal plants used by communities in the city of Vitória de Santo Antão - PE, Brazil.....	88

Understanding the use of nanoparticles in drug delivery systems and its importance in the treatment and monitoring of breast cancer.....	89
Preliminary study of medicines used against androgenetic alopecia by mutagenicity parameter (Ames Test).....	90
Probiotic formulations for the treatment of intestinal constipation: development and evaluation.....	91
Preliminary toxicity assessment of silver nanoparticles (AgNPs) synthesized with curcumin (Ames Test)	92
Development of curcumin gel and choline-geranate for psoriasis treatment.....	93
Preliminary assay: safety assessment of doxycycline by mutagenicity assay in <i>Salmonella typhimurium</i> microsomes	94
The benefits of using the ethanolic extract of <i>Catharanthus roseus</i> L.	95
Biogenic synthesis of silver nanoparticles based on <i>Penicillium</i> sp to combating resistant pathogens.....	96
<i>In vitro</i> study of direct inhibition of cholinesterase enzyme by the organophosphate chlorpyrifos.....	97
Regional analysis of the implementation of pharmaceutical care in the health care network and its perception by pharmaceutical professionals	98

Frequency and severity of adverse drug reactions to medications prescribed for Alzheimer's disease in a Brazilian city: a cross-sectional study

Tânia Regina Ferreira¹; Luciane Cruz Lopes¹; Cristiane de Cassia Bergamaschi¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: There is a lack of national studies that assess the risks associated with the drugs provided under the Brazilian Public Health System for treating Alzheimer's disease.

Objectives: To determine the prevalence and severity of self-reported adverse drug reactions (ADRs) to medications prescribed to patients with Alzheimer's disease in the Brazilian public health system.

Methods: A cross-sectional study was carried out based on public data from the Medex system (information on dispensing data, known as exceptional dispensing medications) and interviews with patients and/or caregivers who get access to Alzheimer's drugs at a public pharmacy in Brazilian large city, between July and September 2017. The subjects were asked about ADRs and serious adverse events (SAEs) related to the use of donepezil, galantamine, rivastigmine, and memantine.

Results: Out of 285 patients enrolled on the database, 250 participated in the study (87.7%). Among the participants, approximately 63.0% were female, 70.3% aged ≥ 75 years, and 70.3% had comorbidities. Overall, 209 patients (83.6%) reported at least one ADR (total 1.14 ADRs). Rivastigmine was associated with the largest number of ADRs per patient (7.9 ADRs/patient). The predominant adverse effects were psychiatry disorders with common frequency (57.1%) and mild severity (89.0%). Six patients (2.4%) had SAEs that required hospitalization. After adjustment, the use of antipsychotics was associated with ADR (OR= 4.95; 95% CI: 1.45–16.93; $p=0.011$).

Conclusion: The number of ADRs reported was large and most were of common frequency and mild severity and were mainly related to psychiatry disorders. Then, it is important to verify if these disorders did not occur due to the use of an anticholinergic drug. Considering the fragility, the use of drugs, and comorbidities in patients, it is important to improve safety-related care in the use of drugs for treating this disease.

Funding: No funding to declare.

Ethics Statement: 1860724.

Analysis of pharmacotherapy among residents who use multiple drugs in Santo André, São Paulo

Felipe de Lara Janz¹; Dones Cláudio Janz Jr²

¹ Anhanguera University, São Paulo, São Paulo, Brazil

² State University of Ponta Grossa, Ponta Grossa, Paraná, Brazil

Context: Pharmaceutical Care (PC) emerged with aim of extending the pharmaceutical role in health care actions and benefiting patients through the rational use of drugs. Many patients, i.e. elderly and people with chronic disease use multiple medications and can present Drug Related Problems (DRPs).

Objectives: This study aims to investigate the occurrence of DRPs among patients in multi-drug therapy in Santo André, SP.

Methods: A case study was conducted to analyze the drug therapy of Santo André residents (over 18 years) in multi-drug therapy (using 2 or more drugs at the same time). We used the Dader method to analyze the drug therapy of Santo André residents (over 18 years) in multi-drug therapy (using 2 or more drugs at the same time). Questionnaires about sex, age, weight, height, physical activity, schooling, drug storage, and pharmacotherapy profile were applied. Problems leading to DRP were mainly classified into three domains (necessity, effectiveness, and safety - DRP 1 to 6)

Results: Of 12 patients included in this study (58.3% male and 41.7% female), 50.0% were aged 26-50 years, 25.0% 18-25 years, and 51-75 years. 50% of the patients were overweight; 33.3% were in the normality and 16.7% were obese. 66.7% studied only to the fundamental level. 66.66% did not perform physical activity. Drug storage was done in drawers, cabinets, and refrigerators. the main source of drug information is the physician (66.7%) and 75% make self-medication. DRP was detected in 33.3% of the cases. The most frequent DRP was PRM 1 (60%). DRP 2 and 6 were also found.

Conclusion: Patients using multi-drug therapy may present DRP, mainly related to necessity. Pharmacotherapeutic follow-up is an important tool to improve the drug rational use and quality of life.

Funding: Project supported by Funadesp.

Ethics statement: 289746.

Home care through the best at home program (MEMC) - a case report

Letícia Cristina Alves de Sousa¹; Bruno Rogério Ferreira¹; Isabela Jubé Wastowski¹

¹ Goiás State University, Goiânia, Brazil

Context: MEMC is a multidisciplinary Home Care program (HC) of the Unified Health System (SUS). The population is aging and with this increase in comorbidities, health complications, polypharmacy, and development of mental disorders. Receiving care at home brings a perspective of improving the quality of life and understanding your clinical condition.

Objectives: The objective was to report the case of an elderly patient attended by MEMC to show how home care interferes with the patient's health and quality of life.

Methods: This is a descriptive and qualitative case study carried out with an elderly patient, attended by the MEMC program; information from the patient's medical records from August 2019 to May 2020 was used.

Results: The case is a female patient, 76 years old and with low social status. The patient was diagnosed with Parkinson's disease, cerebral aneurysm, anemia, venous insufficiency, sacral eschar, schizophrenia, cerebrovascular accident (CVA), hypothyroidism, dementia, and polypharmacy. Food reeducation, patient positioning, bronchial aspiration, psychotherapy, and dressings were the actions performed. Currently, the patient is in good health, with scars, 90% regression, and improved nutritional status.

Conclusion: The MEMC is indicated for people who have temporary or definitive difficulties in locomotion or leaving the space of the house to reach a health unit, providing better conditions for medical care, as the treatment and monitoring are carried out at the patient's home. The recovery of the patient in the comfort of his home, together with family members, reduces the costs of hospitalization. The MEMC improved the eschar, the nutritional and psychological support. There was guidance on how to contain the patient and reduce the appearance of injuries.

Funding: No funding to declare.

Ethics statement: 4374388.

Free access and use of medicines by the elderly population in the municipality of Campinas/SP: results of a population survey

Nadia Baggio Ratti¹; Marilisa Berti de Azevedo Barros¹; Andréia Turmina Fontanella²; Sotero Serrate Mengue²; Karen Sarmento Costa¹

¹ State University of Campinas, Campinas, São Paulo, Brazil

² Federal University of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

Context: The accelerated aging process is characterized by changes in health indicators and increased life expectancy. The elderly are a population segment with a high prevalence of chronic degenerative diseases, and the set of these conditions results in an increased demand for health services and greater use of medicines. Epidemiological studies on the elderly have shown that access to medicines can be influenced by several factors, such as socioeconomic and demographic factors.

Objectives: To analyze aspects related to free access and medication use by the elderly in Campinas/SP.

Methods: This is a cross-sectional population-based study that used data from the health survey of Campinas (ISACAMP2014 2015). The sample consisted of 86 elderly aged 60 years or older.

Results: 88.6% of the elderly population used at least one drug. Free access was higher among women in the separated elderly compared to married, with lower monthly income, among them without health insurance and with fewer chronic diseases. The most used sources for obtaining medicines free of charge were the pharmacies of the unified health system (87.8%) and those accredited by the popular pharmacy program (5.4%). The medications most frequently used were the cardiovascular system (58.4%), such as captopril and enalapril, first-line for the treatment of hypertension, and drugs that act on the digestive tract and metabolism (16.9%), such as metformin and glibenclamide, used to treat diabetes. Those which were purchased on their own, most of them, were drugs for the treatment of stomach acid disorders and heart therapy. Where, 79% of elderly could not get a drug, and the reasons most often identified were not found in the health center and no resources to buy.

Conclusion: The results point out the need to establish new strategies that guarantee equal access to medicines for this population.

Funding: Project supported by Fundação de Amparo à Pesquisa do Estado de São Paulo – Fapesp (2012/23324-3).

Ethics statement: 409714.

Analysis of serious drug interactions in paracetamol poisoning in Brazil

Letícia Mariano Lagemann¹; Julia Hiromi Hori Okuyama¹; Marcus Tolentino Silva ¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Paracetamol poisoning has differences between age groups, intentionality, and use of isolated or compound substances. We have poor evidence about Brazilian poisoning estimates to perform health regulation.

Objectives: To describe paracetamol poisoning associated with other drugs and to analyze the factors associated with serious drug interactions.

Methods: The data come from the notifiable diseases information system (SINAN) registered in 2017. After selecting cases of paracetamol poisoning, records with interaction with pesticides, food, alcohol, and drugs of abuse were excluded. Medicines related to intoxications together with paracetamol were identified, coded and grouped by the anatomical therapeutic chemical classification. The intensity of the interaction was analyzed for severity according to Micromedex. The results were stratified by age group, sex, pharmacological group, federation state, quarter of the year, day of the week, type of interaction, and intentionality. For the analysis of the association between serious drug interactions and the variables of interest, Poisson regression adjusted for estimating the prevalence ratios was calculated.

Results: 813 notifications of poisoning by paracetamol and other drugs were identified in 2017. The prevalence was higher in the population aged 18 to 36 years old (53.9%), woman (76.9%), predominantly in the south/southeast regions in the country. There was a higher frequency in the last quarter (2.9 cases/day) and on weekdays (2.3 cases/day). There were 5.6 cases of attempted suicide for an unintended case. Paracetamol poisoning related to analgesics indicates a 2.6 (95% CI 1.5 - 4.3) greater risk of drug interaction than in its absence and 4.0 (95% CI 2.4 – 6.8) greater in the presence of non-steroidal anti-inflammatory drugs.

Conclusion: Serious drug interactions in cases of paracetamol poisoning have been associated with analgesics, anti-inflammatory drugs, increasing age, and suicide attempts.

Funding: No funding to declare.

Ethics statement: Not applicable.

Promotion of rational use of medicine through social media

Júlia de Souza Faria¹; Camila Pereira de Araújo¹; Maysa da Silva Gonçalves¹; Lavínea Silva de Lima¹; Pamela Souza Almeida Silva Gerheim¹

¹ Federal University of Juiz de Fora, Juiz de Fora, Minas Gerais, Brazil.

Context: Social media are widely used in Brazil and are considered important tools for communication and dissemination of health information.

Objectives: To implement an Instagram social media profile to disseminate content on the rational use of medicines (RUM).

[Methods: The acronym Prosum (promotion of safety in the use of medicines), a logo, and an Instagram account (@prosumufjf) was developed. The posts were made from April 2020 to January 2021, 3 to 4 times a week, conducted by four undergraduate students under the supervision of a pharmaceutical professor. All publications were based on scientific evidence and executed in four stages, being: (1) survey of topics related to rum; with priority for the most current themes, those suggested by followers and content of public utility; (2) research and synthesis of data; (3) preparation and review of the material and (4) posting on Instagram. The posts were made taking into account the need to create attractive material, with short texts, topics, diagrams, and good quality figures.

Results: Prosum had on Instagram 509 followers, being the most prevalent age group 18 to 24 years old, feminine gender, with the location in the city of Juiz de Fora. 104 Posts were made, totaling 5022 likes, 692 comments, 750 shares, and 463 saved. The publication with the theme “ivermectin for the treatment of COVID-19?” Had the highest number of likes, shares and reached 551 accounts. The most reproduced reels, with 1876 views, was “why does antimicrobial resistance increase in the world?” and the most successful igtv, with 629 views was “correct disposal of medicines”. The analysis of the followers’ spontaneous comments was considered as one of the impact indicators of the project, being highlighted by the public the importance of the proposed themes and the didactic and creative format of the publications.

Conclusion: The use of Instagram was an effective communication strategy for the promotion of rum, especially in this context of pandemic and social distancing. It is necessary to plan and publish scientific content clearly and attractively, thus reducing misinformation and the irrational use of medicines.

Funding: Project supported by PROEX/UFJF.

Ethics statement: Not applicable.

Profile of antimicrobial dispensing in primary health care in the context of the COVID-19 pandemic

Antonio Joaquim Bonfim¹; Alane Andreino Ribeiro¹; Clariça Rodrigues Soares¹; Wenderson Walla Andrade¹; Sandra de Castro Barros¹

¹ Ministry of Health, Brasilia, Brazil

Context: The so-called O'Neill report cautions that if the use of antimicrobials in human and veterinary medicine is not reduced, by 2050 increased antimicrobial resistance could take the lives of 10 million people per year worldwide. Furthermore, pathogens with a high level of antimicrobial resistance may have their scope amplified due to the irrational use of antimicrobials during the ongoing coronavirus (COVID-19) pandemic. The use of access, watch, reserve ("aware") classification of antibiotics in the national essential medicines list could significantly contribute to promoting the rational use of antimicrobials and to reducing antimicrobial resistance. Brazil relies on its unified public health system's national database for pharmaceutical care (BNAFAR), which includes inventory status, distribution, and dispensing of medicines as well as strategic health supplies.

Objectives: To analyze the variation in the dispensing of antimicrobials that have been used in the Primary Health Care (PHC) before and during the pandemic in relation to COVID-19.

Methods: A descriptive analysis of dispensing that occurred in 2019 and 2020, using information from the national database.

Results: Drugs that showed a higher increase in the quantity dispensed PHC were cephalexin (29%), azithromycin (100.3%), amoxicillin (15%), ciprofloxacin (32.3%), and clindamycin (30.7%). Three of these medicines are in the access category. Azithromycin and ciprofloxacin are in the watch category in the "aware" classification, which includes antibiotics that have higher resistance potential and include most of the highest priority agents among the critically important antimicrobials for human medicine and/or antibiotics that are at relatively high risk of selection of bacterial resistance. With regard to the number of users treated, what stands out is that the users of azithromycin increased by 86.9% in 2020 when compared to the previous year.

Conclusion: We can highlight the importance of adequately registering to dispense in all of Brazil's states and municipalities in order to ensure access to reliable information on the use of medicines at the national level. The increase in the consumption of these antimicrobials may lead to an increase in the risk of antimicrobial resistance therefore there is a need to carry out monitoring in coordination with other preventive actions and control.

Funding: No funding to declare.

Ethics statement: Not applicable.

Monitoring actions for promoting the rational use of antimicrobials contained in the national action plan for the prevention and control of antimicrobial resistance 2018- 2022 (PAN-BR)

Alane Andreilino Ribeiro¹; Antonio Joaquim Bonfim¹; Clariça Rodrigues Soares¹; Wenderson Walla Andrade¹; Sandra de Castro Barros¹

¹ Ministry of Health, Brasilia, Brazil

Context: Resistance to antimicrobials is one of the greatest challenges to public health, with a significant impact on the health of animals and humans. The promotion of the rational use of antimicrobials requires efforts on various fronts in light of the five strategic objectives of the global action plan. In this regard, the national action plan for the prevention and control of antimicrobial resistance 2018- 2022 (PAN-BR) was developed. This plan converges with the “one health” approach, consolidating human and animal health with environmental issues. With regard to the rational use of antimicrobials, there are activities related to objectives 9 and 11 in the plan.

Objectives: To present the progress of activities promoting the rational use of antimicrobials related to the PAN-BR.

Methods: Descriptive analysis of the activities related to the rational use of medicines as contained in the PAN-BR operational monitoring quarterly plan.

Results: The federal scope developed incentive actions for the creation of a committee for the rational use of medicines at the municipal level and to strengthen the strategies of the pharmaceutical care national policy related to promoting the rational use of medicine, with a focus on the use of antimicrobials, as well incentives for the establishment of pharmaceutical clinical services and the enhancement of existing information systems to monitor the prescription, dispensing and use of antimicrobials. Among the activities in progress, national directives on the rational use of antimicrobials will be elaborated as well as efforts to enhance processes for the selection, distribution programming, and acquisition of antimicrobial, considering health requirements, the epidemiological profile and history of consumption, as well as the improvement of procedures for the distribution and stocking of antimicrobials according to local conditions.

Conclusion: Of the seven aforementioned activities, all have been initiated and four have already been completed. We can highlight that inter-federal coordination is fundamental for implementing activities and that the rational use of antimicrobials should be the priority of states and municipalities governments and society.

Funding: No funding to declare.

Ethics statement: Not applicable.

Analysis of the competence of written communication of prescriptions carried out by Brazilian pharmacy students in the last year of the graduation course

Nathane Stéfanie de Queiroz¹; Fernanda Coelho Vilela¹; Karina Aparecida Resende¹; Ana Elisa Melo¹; Bruna Mundim Cardoso¹; Angelita Cristina Melo¹

¹ Federal University of São João Del-Rei, Divinópolis, Minas Gerais, Brazil

Context: In the field of health service provision is essential that the pharmacist has efficient written communication that ensures the understanding of information and the correct and safe use of medicines.

Objectives: To analyze the competence of Brazilian pharmacy students in the last undergraduate year to carry out written communication in prescription.

Methods: A cross-sectional and multicenter study was carried out, with a simulation of the scenario of a 28-year-old female patient who, considering the pharmacist as presenting with mild allergic rhinitis and without other medical conditions, carried out in five federal universities, one from each Brazilian region, 168 prescriptions of 84 pharmacy students were evaluated. The written communication was double-blind evaluated using the parameters: presence without erasure of signature, contact, professional register, date of appointment, dose, treatment timescale, pharmaceutical form, frequency of drug administration, legible patient name, medicine name, and administration route.

Results: Only 1 student met all the requirements for good written communication. The most attended items were: patient name 77 students (91.6%), Followed by 71 students (84.52%) Included name of the medication (but only 7 students (8.3%) Via administration) and own signature; 64 students (76.20%) Informed the patient's contact and 61 of the prescriptions (73.80%) Were legible. Unfortunately, the time scale of the administration frequency criteria with only 14 students (16.6%), The professional registration number - CRF was the least attended with information from 9 students (10.70%).

Conclusions: The pharmacy students in their final year of graduation showed a lack of competencies to meet the minimum requirements for written communication in prescriptions. We recommend further studies using the same method on larger samples, as well as an in-depth analysis of the educational process and its impact on the development of this competency.

Funding: Project supported by the Coordination for the Improvement of Higher Education Personnel - Brazil (CAPES and the Federal University of São João Del-Rei).

Ethics statement: 2853052.

Impact of polypharmacy on the nutritional status of institutionalized elderly

Mayara Priscilla Dantas Araújo¹

¹ Federal University of Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil

Context: Polypharmacy, defined as the concomitant consumption of five or more medications, and malnutrition are two important health problems that affect institutionalized elderly people.

Objectives: To evaluate the impacts of polypharmacy on the nutritional status of institutionalized elderly.

Methods: This was an integrative review of the literature that sought to answer the question: what are the impacts of polypharmacy on the nutritional status of institutionalized elderly? A search was performed in Medline, PubMed, Web of Science, and Scopus. The selection of studies was made considering complete articles available online, in English, Spanish and Portuguese, published between 2011 and 2021. The following mesh terms and operators were used: aged and institutionalization and polypharmacy and nutritional status. The authors also reviewed the reference lists from the review articles reported to identify possible additional articles for inclusion.

Results: The searches resulted in 7 articles, of which 3 were duplicates. After reading the title and abstract, 2 articles were excluded, 1 for not evaluating institutionalized elderly and 1 for not addressing polypharmacy. The final sample consisted of 2 articles. Polypharmacy is very prevalent among institutionalized elderly and is correlated with the development of malnutrition in these individuals, especially excessive polypharmacy (> 10 drugs). However, moderate use of medications (1-2 drugs) seems to protect older women from malnutrition. Its side effects deserve attention because they indirectly lead to the occurrence of malnutrition due to loss of appetite, nausea and constipation. In addition, institutionalization also contributes to the occurrence of malnutrition, contributing to increased nutritional risk in these individuals.

Conclusion: Polypharmacy negatively impacts the nutritional status of institutionalized elderly, who may also present nutritional deficit stemming from institutionalization itself. The importance of promoting the rational use of medicines in long-term care institutions is highlighted in order to avoid excessive polypharmacy in these individuals and promote the maintenance of adequate nutritional status.

Funding: No funding to declare.

Ethics statement: Not applicable.

Potential drug-drug interactions of clinical importance between antihypertensives: prevalence and risks for patients in primary healthcare

Bruna Mundim Cardoso¹; Ana Elisa Melo¹, Nathane Stéfanie de Queiroz¹; Karina Aparecida Resende¹; Fernanda Coelho Vilela¹; Angelita Cristine de Melo¹

¹ Federal University of São João Del-Rei, Divinópolis Minas Gerais, Brazil

Context: Drug-drug interactions are more prevalent in the frail elderly. Those interactions can increase the chance risk of adverse events.

Objectives: To evaluate drug interactions in primary care and investigates factors combinations of antihypertensives drugs, antihypertensive drugs potentially inappropriate for the frail elderly and use for adults.

Methods: A sectional study performed in the central-western portion of the Brazilian state of Minas Gerais, to analyze the potential drug interactions of clinical importance between antihypertensive agents in adults or frail elderly (65 years old), using statistical learning theory, exhaustive-CHAID algorithm, Pearson chi-square test with Bonferroni adjustment, 10% significance for inclusion in the model and 5% for the final model. The stopping criteria for multivariate analyses of sociodemographic and pharmacotherapy characteristics, anticholinergic syndrome, potentially inappropriate medications were 100 cases in the parent node and 50 cases in the child node.

Results: 6596 patients had antihypertensive medicines on primary health care. The presence of potential drug interactions among antihypertensive drugs of clinical importance occurred in 5.7%. The multivariate analysis showed in the first hierarchical level for explaining has or has not at least one interaction was a total anticholinergic burden. In patients with an anticholinergic burden greater than two has drug-drug interaction around 20%. In the second hierarchical level for explain has or has not at least one interaction was the number of dispensed medicines in the month. The patients have two medicines dispensed has 11% of drug-drug interaction, and was potentially inappropriate for the older adult, has increased around 10% of drug-drug interaction. In the last hierarchical level, the patients who used between 1-3 medicines and were potentially inappropriate, has around 14% of drug-drug interaction.

Conclusion: There seems to be a relationship between the increase anticholinergic syndrome with a chance of being frail elderly using medicine of potentially inappropriate medicine list.

Funding: No funding to declare.

Ethics statement: 1520218.

Rethinking HIV/AIDS prevention public policies for the elderly based on the discourses of elders living with the virus

Marília Borborema Rodrigues Cerqueira¹

¹ State University of Montes Claros, Montes Claros, Minas Gerais, Brazil

Context: The proportion of elderly people in the population is growing, as is the number of cases of HIV/AIDS in this population group.

Objectives: The objective of this study was to analyze the discourses of elderly people living with HIV/AIDS, seeking elements that may subsidize the development of public policies for the promotion of sexual health and prevention of sexually transmitted diseases aimed at the elderly.

Methods: The method adopted was the rereading of interviews with elderly people living with HIV/AIDS, adopting content analysis. Twenty elderly people assisted in a public hospital in a Brazilian metropolis were approached. The inclusion criteria were being 60 years old or older, positive diagnosis for HIV, and agreeing to participate in the research; as exclusion criteria, it was the non-acceptance of the elderly or the difficulty in participating for some reason. The study was approved by the Research Ethics Committee of the Federal University of Minas Gerais and the Hospital itself. The interviews have information about the sexual life of the elderly, the discovery of their HIV status, their knowledge about HIV/ AIDS, and related topics.

Results: The main results show that the elderly do not have elaborate knowledge about the virus and do not know public policies in the sexual area, as well as health professionals, do not talk about sex, and the anti-HIV test was only done in the Hospital, the last health service where they were seen, causing the late diagnosis of HIV/AIDS.

Conclusion: Therefore, we conclude that the sexual invisibility of the elderly still exists, that is, there is a myth that the elderly do not have sex.

Funding: Project supported by Fapemig.

Ethics statement: 536872.

Use of the wise application as a tool in the therapeutic conduct of the elderly with chronic pain

Ana Karinna Silveira Pereira Caracas¹; Roberto Augusto Caracas Neto²; Tecia Vieira Carvalho²

¹ São Leopoldo Mandic College, Campinas, São Paulo, Brazil

² Federal Institute of Education, Science and Technology of Ceará, Fortaleza, Ceará, Brazil

Context: Chronic pains is multidisciplinary, subjective experience, associated with a psychological state and responsible for the decrease in an individual's quality of life. In this context, a web-based interdisciplinary system assessment (wise) was developed in Zurich

- Switzerland, translated into Brazilian Portuguese, containing objective and subjective questions about pain complaints and the use of drug therapies performed by the patient. The tool includes options for body design, biopsychosocial, and treatment assessment, including drug therapies which the patient is undergoing.

Objectives: The objective of this study was to evaluate the prevalence and effectiveness of drug treatment for chronic pain in elderly people living in the Praia do Futuro II community, attended at the primary care unit Frei Tito de Alencar Lima, in Fortaleza – CE, due to a deficiency of published academic works related to this subject.

Methods: The research using the web-based wise application went cross-sectional. The sample consisted of 250 elderly people, over 60 years old of both genders, with cognitive ability to answer the questionnaire.

Results: Of the interviewees, 72% are female and 28% male. Despite various comorbidities, the prevalence of elderly people who frequently use pain medication was 22.5%, While 50% did not undergo any pain treatment, 12.5% Underwent dental treatment, 7.5% Psychological treatment, 5% surgical and 2.5% Physiotherapy treatment. Regarding the effectiveness of the treatment, on a scale of 0 to 10, 50% reported low effectiveness, 10%effectiveness medium and 40% good or excellent effectiveness.

Conclusion: These results lead us to reflect on the need for a multi-professional team with cognitive behavioral therapy, a good referral, and counter-referral system at all levels of health care, looking for the origin of the suffering associated with chronic pain.

Funding: No funding to declare.

Ethics statement: 3706607.

Indiscriminate use of ascorbic acid during the COVID-19 pandemic in Brazil

Carmem Lúcia de Arroxelas Silva¹; Carlos Antônio de Arroxelas Silva²; Olagide Wagner de Castro¹

¹ Federal University of Alagoas, Maceió, Alagoas, Brazil

² Federal University of Acre, Rio Branco, Acre, Brazil

Context: The COVID-19 pandemic is characterized by a contagious infection caused by the Sars-cov-2 virus, which can compromise the respiratory system. This disease has become a challenge for global health, due to the rapid contagion and the high number of deaths. Current reports indicated an increase in the consumption of ascorbic acid (vitamin C) in an attempt, scientifically unfounded, to prevent and/or cure the COVID-19. This type of consumption corresponds to the practice of non-rational use of medicines, characterized by the use of medicines without guidance or prescription from health professionals. The use of medications in these circumstances can mitigate the aggravation of diseases, promoting reactions of intoxication and drug ineffectiveness.

Objectives: The objective of the present study was to evaluate the commercialization records of vitamin C to the Brazilian population during 2019 and 2020.

Methods: A descriptive observational study was designed based on secondary data obtained from the survey released by the federal pharmacy council of Brazil of vitamin C sales in 2019 and 2020. The data were analyzed in the form of relative frequency.

Results: The results showed a 180.01% increase in the number of vitamin C sales in Brazilian states in 2020 when compared to 2019. Additionally, the sales ratio between states and regions in Brazil, showed that the state of Bahia was the highest increase (316.6%), Followed by the southeast region (176.1%).

Conclusion: Our data indicate an existing culture in Brazil of self-medication and lack of knowledge about the harmful effects of the use of vitamin C. The reported negative effects can range from gastrointestinal and kidney problems, electrolyte disturbances, and increased iron overload absorption. The correct medic indication, following the precepts of science, will contribute to demystifying alternative therapies against COVID-19 that can lead to serious side effects.

Funding: No funding to declare.

Ethics statement: Not applicable.

COVID-19 pandemic induces increase in clonazepam consumption in the state of Alagoas, Brazil, 2020

Carlos Antônio de Arroxelas Silva¹; Carmem Lúcia de Arroxelas Silva²; Olagide Wagner de Castro²

¹ Federal University of Acre, Rio Branco, Acre, Brazil

² Federal University of Alagoas, Maceió, Alagoas, Brazil

Context: COVID-19 pandemic increased the frequency of neuropsychopharmacological disorders in the world population. Social isolation and a high rate of mortality may be a trigger for the development of these conditions. In this scenario, clonazepam is prescribed with frequency.

Objectives: To evaluate commercial records of sales of clonazepam in Alagoas state, Brazil, during the year 2020.

Methods: A descriptive observational study was designed from secondary data obtained from the National System for the Management of Controlled Products (SNGPC) of the National Health Surveillance Agency (ANVISA).

Results: Our results showed that clonazepam is the most sold controlled drug in the state of Alagoas. The months of May (20.467 units) and July (21.302 units) were the biggest sales when compared to other months of the year.

Conclusion: The expressive increase in the number of prescriptions raises the question of the real need for use of clonazepam. The chronic use of clonazepam is associated with a high risk of dependence, withdrawal syndrome, impairment of motor and mental functions. Additionally, there is a lack of public policies that provide appropriate clinical assistance for the correct use of clonazepam.

Funding: No funding to declare.

Ethics statement: Not applicable.

Eating habits and physical exercise in patients with type 1 *Diabetes mellitus*: is there a difference before and after pharmacotherapeutic follow-up?

Thays Santos Mendonça¹; William Neves Oliveira¹; Vinícius Silva Belo¹; Eduardo Sérgio Silva¹; Mariana Linhares Pereira¹; Paulo Roque Obreli-Neto²; André De Oliveira Baldoni¹

¹ Federal University of São João Del-Rei, Divinópolis, Minas Gerais, Brazil

² Faculdades Integradas de Ourinhos College, Minas Gerais, Brazil

Context: Pharmacotherapeutic follow-up has been shown to be beneficial for better control of *Diabetes mellitus* (DM). Currently, insulin analogues have been in high demand through the judicialization of health. However, there is a lack of studies that assess the effectiveness of pharmacotherapeutic follow-up on eating habits and physical exercise in these patients.

Objectives: To evaluate the effectiveness of pharmacotherapeutic follow-up on eating habits and physical exercise in patients with type 1 *Diabetes mellitus* (T1DM) who receive insulin analogues by judicial decision in a Brazilian municipality.

Methods: A single-arm intervention study was conducted through pharmacotherapeutic follow-up in patients with t1dm between April 2019 and August 2020. Participating patients underwent five pharmaceutical consultations. The consultations were conducted following the method adapted from the pharmacotherapy workup (PW), with the approach of aspects related to the general characteristics of DM, eating habits and physical exercises, blood glucose monitoring, training for self-care, and prevention of complications acute and chronic diseases through validated booklets. In relation to non-pharmacological treatment, an instrument was developed to collect data regarding the frequency of food, a diet with restricted sugars and carbohydrates, and physical exercises with at least 150 minutes per week. To compare the variables before and after the intervention, the Wilcoxon and McNemar tests were used, with a significance level of 5%.

Results: Of the 93 patients who received insulin analogues through judicial decision in 2019, 41 (44.1%) Agreed to participate. However, only 28 (68.3%) Participated in all stages. Of these, the majority were female (n = 15; 53.6%), with an average age of 32.8 ± 11.6 years. After the intervention, there was an improvement in eating habits {frequency of feeding (p = 0.012) And diet (p = 0.006)}. There was an increase in the number of patients who started to exercise, however, this did not present statistical significance.

Conclusion: The pharmacotherapeutic follow-up promoted behavioral benefits, with improvement in the eating habits of patients with T1DM who receive insulin analogues through judicial decision. This shows that this type of service can be a great ally in minimizing the negative impacts of the illness of judicialization of health.

Funding: Project supported by the Brazilian National Council for Scientific and Technological Development (CNPQ) (405145/2018-7) and by the Coordination for the Improvement of Higher Education Personnel – Brazil (CAPES).

Ethics statement: 87590518.9.000

Evaluation of medication use by public school teachers in Pelotas-RS

Paulo Maximiliano Correa¹; Patrícia Gomes Machado¹; Juliane Fernandes Monks da Silva¹

¹ Federal University of Pelotas, Pelotas, Rio Grande do Sul, Brazil

Context: Studies demonstrate that teachers have been suggested as one of the professional categories more predisposed to stress due to the work environment, to relieve their tension end up using medicines to solve the pain and illness acquired during the exercise of teaching.

Objectives: The present work aims to characterize the profile of the use of medicines by teachers of kindergarten, elementary and secondary education of three public schools in the city of Pelotas/RS.

Methods: A cross-sectional, descriptive, analytical study with a qualitative and quantitative approach was conducted, between June and September 2019, using a structured questionnaire.

Results: The sample consisted of 23 teachers, all women with ages between 33 to 63 years old. Participant's income ranges from three to five minimum wage. Regarding academic training all participants have university degrees, 69% have a specialization, 13% master's degrees, and 4% doctorates. 61% Of the interviewed teachers had some chronic disease,

91.3% of teachers used some medication in the last 15 days before the survey. 64% Stated that teaching affected their health. The most reported reasons for using the medication were hypertension (39.2%), depression and pain relief (26% each), anxiety (17.4%). 82.6% Of respondents believed that medications do not pose a health risk. Regarding self-medication, 39.1% do it once a month, 21.7% use self-medication more of them once a month, 17.4% self-medicated at least once a week.

Conclusion: significant use of medicines was found when compared to other studies carried out with teachers from different regions of Brazil. The high prevalence of self-medication indicates indiscriminate use of medication by this class of professionals. The lack of knowledge about the risk of self-medication is a concern for the health of teachers. Besides, we can observe that for this group of people, the level of education does not seem to have influenced the knowledge toward the risks of self-medication.

Funding: No funding to declare.

Ethics statement: 1787268.

Association between anticholinergic burden of medications and emergency care in the elderly assisted in primary health care, Sorocaba, SP, Brazil

Rodrigo Sugimoto Iwami¹; Fabiane Raquel Motter¹; Vera Maria Vieira Paniz²; Anna Luiza De Crescenzo Brotto¹; Luciane Cruz Lopes¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² Univerity of Vale do Rio dos Sinos, São Leopoldo, Rio Grande do Sul, Brazil

Context: Anticholinergic drugs can reduce physical and cognitive function, especially in the older population due to their reduced acetylcholine physiological reserve. Since these effects are dose-related and cumulative, it's critical to evaluate this population exposure to this type of drug and assess its influence on their health status.

Objectives: Evaluate the association between anticholinergic burden of medications (ABM) and the need of emergency care in the elderly assisted in Primary Health Care (PHC) in Sorocaba, SP.

Methods: Cross-sectional study involving people with age 65 years assisted in PHC Maria do Carmo between January and March of 2019. Data was collected through interviews with the elderly and/or caregivers. ABM was assessed using the Brazilian anticholinergic activity drug scale and dichotomized into 0-1 and >1. Emergency care was evaluated through the question "In the last 12 months, how many times have you been in an emergency room without being hospitalized?". Poisson regression was used to evaluate the association between ABM and emergency care, adjusted for confounders (sex, age, ethnicity, falls and polypharmacy).

Results: Of 125 elders interviewed, 74 (59.2%) were women; median age 72 (65-95), with <75 years comprehending 84 (68.3%); 50 (40.0%) who had fell in the last 12 months; median ABM 1 (0-7) and mean ABM 1.1 (SD±1.54), of which 27 (22.0%) had ABM >1; and 65 (52.0%) used at least one anticholinergic drug from a total of 92 identified. The most used drugs were Atenolol (n=15; 12.0%), Furosemide (n=8; 6.4%) and Captopril (n=7; 5.6%). The association between ABM>1 and emergency care 2 in the last 12 months was significant (p=0.016; RP=3.4; CI95%=1.27-9.02), even after adjusting for confounders.

Conclusions: Even in a small sample the association was significant, highlighting the impact of ABM in the health care service of the elderly and the need of more studies with bigger sample sizes.

Funding: Project supported by CNPq (number: 426720/2016-4) and FRM is supported by CAPES.

Ethics statement: 3773012.

Self-medication in Pernambuco: a preliminary analysis of SINAN data from 2017 to 2019

João Lucas Silva Pinheiro¹; Bruna Kristyer Lima de Paula¹, Camila Gomes de Melo¹

¹ Maurício de Nassau College, Aracaju, Sergipe, Brazil

Context: Self-medication represents a worldwide health complexity. The practice can promote several problems such as allergic reactions, drug resistance, dependence, and death. The rational use of medicines is directly related to proven and acceptable efficacy and safety, thus avoiding potential problems arising from incorrect use.

Objectives: To analyze and systematize the data regarding the cases of self-medication notifications in Pernambuco.

Methods: This is a study of data collection from the Notifiable Diseases Information System (SINAN) and Department of Informatics of the Unified Health System (DATASUS), linked to the Ministry of Health. The notifications of self-medication cases in the period from 2017 to 2019 were analyzed, with the following variables: gender, age group, city of notification, ethnicity, and type of exposure. After compilation, the data were organized in a systematic way.

Results: During the analyzed period, 1302 notifications were recorded, occurring an increase in consecutive years. 33.2% were male and 66.7% female. The most affected age group was 20 to 39 years - 38.9%, followed by 10 to 19 years - 28.3%. The highest incidence per city of notification occurred in Recife - 34.2%. It was observed that browns were more evident throughout the study - 60.8%. Moreover, the acute-only exposure was about 80.7%.

Conclusion: We identified a higher incidence in the female and young population, proposing causes such as socioeconomic independence and precariousness of information by the health system. Thus, it is necessary to extend the services of qualified pharmaceutical care for these groups through campaigns and educational activities, especially in health facilities and schools, in order to promote the rational use of medicines, addressing the main impacts of self-medication.

Funding: No funding to declare.

Ethics statement: Not applicable.

Pharmaceutical care in critical patients and the impact on the use of financial resources with medicines

Maurício Fábio Gomes¹; Camilo Molino Guidoni²

¹ State University of Maringá, Maringá, Paraná, Brazil

² State University of Londrina, Londrina, Paraná, Brazil

Context: Patients admitted to intensive care units have critical health conditions and generally need treatment with numerous medications with the potential to cause damage to health. In this way, the clinical pharmacist, by providing pharmaceutical care to critically ill patients, is able to optimize therapy in terms of needs, effectiveness, safety and costs.

Objectives: Analyze pharmaceutical interventions and the impact on the use of financial resources that the pharmaceutical care service promoted among critically ill patients admitted to the intensive care unit of a public hospital in Paraná.

Methods: This is a retrospective cross-sectional study which assessed pharmacotherapeutic follow-up sheets of patients assisted between July/2019 and April/2020 in the intensive care unit. Descriptive statistics were used to quantify the drugs introduced or suspended from therapy after the intervention of the clinical pharmacist. The measured values were obtained in the notices of the last bid in force during the study period. The measured values were obtained in the notices of the last bid in force during the study period. The cost figures took into account the number of therapeutic units saved/spent (ampoules, vials, tablets, others), not considering costs with materials, thinners, human resources, others.

Results: Of the 350 pharmacotherapeutic problems, 80% led to pharmaceutical interventions that impacted the consumption of therapeutic resources both when including and when suspending a drug and, these provided savings of R\$ 14,107.29. Of the proposed interventions, 81.4% indicated the need to suspend or reduce the dosage of certain therapeutic agents for reasons related to the need and safety of treatment, such as adjustment to renal function as well as counting doses of antimicrobials.

Conclusion: Thus, we conclude that pharmaceutical care had a positive impact on the therapy provided to critical patients, reducing costs to the public purse by optimizing the use of therapeutic resources.

Funding: No funding to declare.

Ethics statement: 347424.

Risk of infections during treatment of juvenile idiopathic arthritis with adalimumab versus other indications

Nayara Aparecida de Oliveira Silva¹; Daniela Oliveira de Melo¹

¹ Federal University of São Paulo, São Paulo, Brazil

Context: Juvenile idiopathic arthritis (JIA) is a chronic autoimmune rheumatic disease affecting individuals up to 16 years of age. The immunosuppressive monoclonal antibodies (mAB), such as adalimumab, are widely used in the management of JIA. Despite their good therapeutic results, these mABs may increase the risk of serious infections, requiring pharmacovigilance monitoring, especially in pediatric patients.

Objectives: This study aimed to compare the infectious profile of adalimumab in patients with JIA versus other indications more frequent in adults (e.g., psoriatic arthropathy rheumatoid arthritis, ankylosing spondylitis, Cronh's disease and psoriasis).

Methods: Reports of adalimumab registered by June 30, 2020 from the Food and Drug Administration Adverse Event Reporting System (FAERS) Public Dashboard were extracted. The analysis of the risk of developing infection events with adalimumab for JIA vs other indications was performed by calculating the Reporting Odds Ratios (ROR) with 95% confidence interval (CI). A potential signal was defined when ROR > 2.0 and minimal CI > 1.0.

Results: We identified 472,575 reports for adalimumab, of which JIA indication was present in 3,934 (0.8%). The JIA indication versus other indications had the highest risk of systemic infections such as septic shock (ROR 39.61 [33.36-47.03]) disseminated tuberculosis (ROR 3.45 [1.63-7.32]) and frequent infections in children and adolescents with emphasis on tonsillitis (ROR 4.99 [3.02-8.24]), infectious mononucleosis (ROR 5.93 [2.91-12.07]) and varicella (ROR 16.18 [9.06-28.90]).

Conclusion: Patients using adalimumab for JIA seem to be at increased risk of systemic infections when compared to other indications. Despite the limitations of the study, such as the absence of information in spontaneous reports of suspected adverse events about concomitant diseases and medications, the disproportionality analysis is large enough to indicate the need for further investigation.

Funding: No funding to declare.

Ethics statement: Not applicable.

Safety in the disposal of medicines: perspectives on rational use

Elba Lúcia Cavalcanti De Amorim¹; Jorge Veras Filho¹; Isa Renata De Souza Santos¹; Marcos Antônio Da Silva Araújo¹; Jéssica Wedna Da Silva Alves¹; Josenildo Pessoa Sena¹; Alex De Lima Dias¹; Maria Érika Da Silva Vilela¹; José Carlos Ferreira Júnior¹; Ítalo Rafael Alves De Souza¹; Filipe José Plácido De Oliveira¹; Juliana Martins De Faria¹

¹ Federal University of Pernambuco, Recife, Pernambuco, Brazil

Context: Medications have important relevance in the population's health and are present in every sphere of healthcare, with the potential of relieving symptoms and curing diseases. On another hand, the lack of information and professional guidance leads people to self-medicate causing a problem to the public health.

Objectives: To do a triage, by therapeutic class, of medications disposed at the collection point of the Pharmaceutical Science Department, at UFPE, between August and December of 2020, to evaluate the classes mostly discarded and their impact on the irrational use of medications.

Methods: Initially, the medications were collected as part of the university extension project "Safety in the Disposal of Medications: Our Responsibility". The medicines were separated from their packaging and leaflets which were sent to be recycled and then cataloged by name, pharmacological classification, pharmaceutical identification, and expiration date. After that, each class was quantified based on weight and statistical analysis was done with the help of the Excel Office 16 program.

Results: 312 medications were triaged, separated into 57 therapeutic classes. Five of them prevailed: antimicrobials (7.69%), NSAIDS (10.89%), antihypertensives (7.69%), vitamins (7.69%) and antihistamines (4.8%). From this analysis, we can notice that the highest percentage was of the antimicrobials. Besides, they are classes that do not require medical prescription.

Conclusion: Given the presented results, it is possible to observe that there is a great prevalence of medications that do not require a prescription, these are allowed by the sanitary authorities to treat minor ailments, but that still deserve attention because the indiscriminate use still brings risks to the individual. Another important data shows that the most disposed medication in this study, the antimicrobials when used and disposed of irrationally, can generate microbial resistance, a public healthcare issue that worries our society.

Funding: Project supported by PROEXC/UFPE.

Ethics statement: Not applicable.

Insulin therapy among users of a primary health care unit

Ruaan Oliveira Carvalho¹; Ana Caroline Silva Santos²; Izabel Cristina Pereira Rocha¹; Chiara Erminia Da Rocha¹

¹ Federal University of Sergipe, Aracaju, Sergipe, Brazil

² São Paulo State University, São Paulo, Brazil

Context: Diabetes mellitus (DM) is a disease whose main characteristic is hyperglycemia. Considered a public health problem, DM has a therapeutic option which is insulin treatment. However, for correct use, is required for the handler to follows essential steps for glycemic control from preparation to application.

Objectives: The study aims to assess the level of knowledge about insulin application.

Methods: This is a cross-sectional study, carried out from November to February 2021, in a Basic Health Unit in the city of Lagarto/SE. The sample was convenient and non-probabilistic. The collection instrument included information about sociodemographic data, types of insulin, time of treatment, usage (preparation and application), and insulin storage. The data were analyzed through descriptive statistics using Excel.

Results: The sample consisted of 73 users. All participants (100%) reported understanding the questions asked by the interviewer. The analysis revealed that 67.1% (n=49) were female and 49.3% (n=36) had incomplete elementary education level. Regarding to the insulin type, 56.1% (n=41) used only NPH insulin and 65.7% (n=48) applied it through a syringe. The level of insulin application knowledge was considered satisfactory for 72% (n=18) and 58% (n=24) of those who applied with the pen and syringe, respectively. Non-compliance with the steps for the correct and safe use of insulin therapy was observed concerning the cleaning of the rubber of the insulin bottle (42.5%; n=31), the introduction of the syringe needle into the insulin bottle and injection of air (56.2%; n=41), cleaning of the application area with alcohol (46.6%; n=34), compression of the application area with cotton (65.7%; n=48), and rubbing of the administration site (64.4%; n=47). The medicine storage was on the middle shelf of the refrigerator (58.1%; n= 43).

Conclusion: There is a lack of information ranging from preparation to the administration of insulin. In this manner, the provision of insulin therapy information to patients fulfills the perspective of promoting rational use of medicine and meets the actions of health promotion and disease prevention in primary health care.

Funding: No funding to declare.

Ethics statement: 16979.

Evaluation of a control system for the prescription of restricted antibiotics in a large Brazilian hospital

Isaltino Pereira de Andrade Junior¹; Fernando de Sá Del Fiol¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Antibiotics are used unnecessarily without controls and standardization in their prescription. These factors have led to the emergence of increasingly resistant microorganisms, which have no known means of treatment. Restriction policies for the use of antibiotics in hospital settings in Brazil are those of a restrictive nature (i.e., the professional must justify the prescription of certain antibiotics, called restricted antibiotics, for HICC to approve the prescription). The HICC should be based on epidemiological data and protocols for the treatment of the particular infection, approve or not approve the prescribed antibiotic, justifying its decision.

Objectives: This study aimed to evaluate patterns related to the request for restricted antibiotics by the physicians of a Brazilian public hospital in order to examine the reasons for authorization and non-authorization.

Methods: This study evaluated a system for the prescription of restricted antibiotics in a large hospital in Brazil. We evaluated 26,612 requests of 22 restricted antibiotics using a computerized system to request and approve these antibiotics.

Results: Results showed that 95% of restricted antibiotic applications were approved, while only 0.52% were not approved. The lack of microbiological data to support approval decisions and the lack of patient clinical data may be responsible for the frequency of approval of these drugs. The lack of standardization of treatments was also evident in the large discrepancy in the doses and durations of treatments with these antibiotics. It is essential that the hospital's management ensures technical and structural conditions for effective antibiotic stewardship to be implemented, generating financial savings and reducing instances of antimicrobial resistance.

Conclusion: These data show that the simple control of antibiotics is not sufficient for the establishment of an antibiotic stewardship system. It is necessary that all multidisciplinary teams are involved, and that the hospital management offers all the technical and infrastructure conditions necessary to implement rational antibiotic prescription and approval control.

Funding: No funding to declare.

Ethics statement: Not applicable.

Trend in the use of atypical antipsychotics in the Brazilian Public Health System: a cross-sectional study

Izabela Fulone¹; Marcus Tolentino Silva¹; Luciane Cruz Lopes¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Atypical antipsychotics are considered the first-line treatment for schizophrenia and their use has increased substantially in all age groups worldwide. Although the exposure to potential adverse effects, as well as the high cost they represent for the health system, few studies have explored the trend of using these drugs in the Brazilian National Health System (SUS).

Objectives: To investigate the use of atypical antipsychotics in patients with schizophrenia or schizoaffective disorder assisted by SUS.

Methods: A descriptive cross-sectional study developed through the records of the Authorizations of High Complexity Procedures (APAC) from the Ambulatory Information System (SIA/SUS), from 2008 to 2017.

Results: Of the 759,654 identified patients, 50.5% were women, aged 44 years (± 0.02), from the southeastern region (60.2%), and diagnosed with paranoid schizophrenia (77.6%). About 8.6% children/adolescents (up to 18 years), 65.6% adults (19 to 59 years) and 21.2% elderly (60 years or more). There was a predominance of males among children/adolescents and adults. In children/adolescents, the use of risperidone stood out (63.3%); in adults, olanzapine (34.0%) and, in the elderly, quetiapine (47.4%). The switch of atypical antipsychotics occurred more frequently in adults (14.5%). The time-mean of use was 32 months (± 0.4) and the abandonment to the assistance program in six months was 24.8%.

Conclusion: The pattern of use of atypical antipsychotics differed according to age. The findings may guide public policies as well as the planning of actions, the offer of the drugs with high unit value, and the improvement of the treatment of schizophrenia in SUS."

Funding: Project supported by FAPESP (2017/20668-7).

Ethics statement: Not applicable.

Is home blood pressure monitoring an opportunity to deprescribing antihypertensive drugs?

Pablo Maciel Brasil Moreira¹; Priscila Ribeiro de Castro²; Brunna Santos Oliveira²; Erlan Canguçu Aguiar²; Halanna Rocha Ferraz¹; Kleiton Coelho de Almeida¹; Márcio Galvão Oliveira²

¹ Vitória da Conquista Health Department, Vitória da Conquista, Bahia Brazil

² Federal University of Bahia, Vitória da Conquista, Bahia, Brazil

Context: Home blood pressure monitoring (HBPM) permits better follow-up of hypertensive patients and helps the identification of possible transient uncontrolled blood pressure (BP). As a result, patients with the white-coat effect using HBPM are not subject to overtreatment with antihypertensive drugs.

Objectives: To describe preliminary data of an HBPM study on the identification of possible hypertension overtreatment in primary care.

Methods: This is a cohort study developed in a public pharmacy in a Brazilian municipality. The study's participants were followed up by the clinical pharmacy service. All had been diagnosed with hypertension and presented uncontrolled BP, according to the goals established by the Brazilian Guidelines for Arterial Hypertension. Each patient received an automated blood pressure measuring device, Afib model (Microlife®), to be used at home for a period of seven consecutive days, with two measurements per day, one in the morning and one at night, totaling 42 measurements. They were all trained on how to use the equipment which conducts three BP measurements and calculates the mean. It is also possible for these measurements to be transferred to a database through software developed specifically for this project. Participants' data were also collected during appointments. We used the t-test and chi-square to compare mean BP and proportion of patients with BP measurement off target.

Results: 100 patients participated in the study, of which 66% were female, aged 61.4 on mean. In relation to the results of the HBPM, 65% of patients presented mean BP values above the proposed target whereas 91% had presented altered measurements in the pharmacist's office ($p < 0.01$). The mean difference between systolic blood pressure values collected in the office and the ones obtained using HBPM was 13.4 mmHg; for diastolic blood pressure, this difference was 6.9 mmHg ($p < 0.01$).

Conclusion: Preliminary data show that the use of HBPM contributed to the identification of a significant number of patients with possible hypertension overtreatment who are candidates for deprescribing antihypertensive drugs or dosage reduction. A pharmacotherapy review conducted by pharmacists in partnership with family physicians and patients may contribute to the deprescribing of antihypertensive drugs in this study's population.

Funding: Project supported by Ministry of Health (PROADI-SUS 25000.028646/2018-10) and Medtronic Foundation.

Ethics statement: 1861073.

Opioids in Brazilian federal hospitals

Patricia Krauze de Almeida¹; Elaine Miranda²; Claudia Osorio de Castro¹

¹ Sergio Arouca National School of Public Health, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

² Fluminense Federal University – UFF, Niterói, Rio de Janeiro, Brazil

Context: Global consumption and deaths related to inappropriate opioid use have increased. While countries in the northern hemisphere debate about an opioid epidemic, countries in the south, such as Brazil, still need studies to measure differences in their use and investigate how inpatient consumption is contributing to this increase.

Objectives: To analyze public purchases of opioids in Brazilian federal hospitals.

Methods: A longitudinal study of consumption was performed, using purchases as a proxy for consumption. Opioids were selected based on national and international consumption trends and relevance of use, from 2010 to 2019, in Brazilian federal hospitals. The bidding data was collected from the Integrated System of General Services Administration (SIASG) and hospital data from the national registry of health establishments (CNES). Hospitals were stratified by region, the number of beds, complexity, and service characteristics. Consumption was expressed in mg / 100 beds (since not all opioids had DDD or s-DDD and occupancy rates were not available for calculating bed-days).

Results: The general consumption of opioids in 85 federal hospitals increased 159% over the ten-years, from 3.4 Million mg/100 beds in 2010 to 8.8 in 2019. The highest values were recorded in 2014 (36.8 thousand mg/100 beds) in 7 hospitals in the Midwest region, which consume 5.5 Times more than the southeast region, with 17 hospitals. In 2019, 25 hospitals in the southeast region consumed 3.5 Times the number of opioids compared to the other four Brazilian regions. Tramadol and morphine were the most consumed opioids, corresponding to more than 95% of purchases during the study period.

Conclusion: Despite the limitations regarding the availability of data for calculating the NDDD/100 bed-days, there was a substantial increase in consumption and regional disparities in federal hospitals. As federal hospitals concentrate a considerable amount of highly complex care in the public sector, measuring opioid consumption by these facilities is important to keep up with general trends in Brazil.

Funding: No funding to declare.

Ethics statement: Not applicable.

Falls among the elderly assisted in primary health care: prevalence and associated factors

Nathalia Margarida Cantuaria¹; Fabiane Raquel Motter¹; Vera Maria Vieira Paniz²; Anna Luiza de Crescenzo Brotto¹; Luciane Cruz Lopes¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² Univerity of Vale do Rio dos Sinos, São Leopoldo, Rio Grande do Sul, Brazil

Context: Falls among the elderly represent a serious public health problem in Brazil and worldwide, being one of the leading causes of hospitalization and death in this population. Identifying the factors associated with the occurrence of this event can aid in the development of prevention strategies in order to reduce their impact on the health system and the quality of life of this population.

Objectives: To determine the prevalence of falls in the last 12 months and associated factors among elderly patients assisted in a Primary Health Care (PHC) facility, Sorocaba, SP, Brazil.

Methods: A cross-sectional study was conducted with elderly individuals 65 years or more), who were assisted in the PHC facility Maria do Carmo between January and March of 2020. An interviewer-administered questionnaire was used to collect the data. Falls were evaluated through the question "Have you had any falls in the last twelve months?". The association between falls and independent variables (sex, age, education, comorbidities, polypharmacy, and hospitalization) was analyzed by Poisson regression.

Results: From the 125 participants, 59.2% were female, and the average age was of 73.2 ± 6.7 years. The prevalence of falls among older adults in the last year was 40.0% (95%CI 31.3-48.7), Among the older adults who fell (n=50), 56.0% had a single fall, 16.3% had a fracture as a consequence and 10.2% were hospitalized. After adjustment, female gender (PR = 1.95; 95%CI 1.03-3.67; p=0.040), less education (PR=2.07; 95%CI 1.01-4.30; p=0.050), and hospitalization in the previous year (PR= 2.04; 95%CI 1.01- 4.14; p=0.048) were associated with the occurrence of falls.

Conclusion: In this sample, a high prevalence of falls among older patients was observed, demonstrating the need for educational actions within the context of primary health care, especially with a focus on women with less education."

Funding: Project supported by CNPq (Process: 426720/2016-4); CAPES supported fellowship for Fabiane Raquel Motter.

Ethics statement: 377301.

The role of pharmaceutical intervention in the management of drug shortage in an epidemic context

Matheus William Becker¹; Gustavo Aristimunha de Almeida²; Carine Raquel Blatt²

¹ Hospital de Pronto Socorro de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil

² Federal University of Health and Science of Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil

Context: The COVID-19 pandemic directly impacted the shortage of drugs, including antibiotics. In this context, the pharmacist plays a fundamental role in solving problems of shortages and rational use of medicines.

Objectives: To evaluate the results of the pharmacist's intervention in relation to the management of the shortage of injectable antibiotics Clindamycin and Amoxicillin/Clavulanate.

Methods: Retrospective data collection study and descriptive data analysis. Through a real problem of the risk of a shortage of injectable drugs Clindamycin and Amoxicillin/Clavulanate, all prescriptions of these drugs were evaluated in November and December, 2020. The month of October was considered as a baseline to compare the influence of consumption by interventions. The interventions were performed on the first prescription day: suggestion to substitute the pharmaceutical form, substitution with a medication with a similar spectrum, or decrease in treatment time. Changes made up to 48 hours after the intervention was considered accepted.

Results: Between November and December, 29 prescriptions were evaluated, with 10/17 (58%) interventions for Amoxicillin/Clavulanate and 7/12 (58%) interventions for Clindamycin being accepted. The reduction in consumption of Clindamycin was 75% and 63%, and of Amoxicillin/Clavulanate by 69% and 90%, respectively. The reduction in consumption prevented shortages.

Conclusion: Despite a median percentage of acceptance, the intervention prevented the shortage of medicines. It was observed that the prescription of these drugs decreased, therefore the hypothesis is raised that the intervention could have an educational effect and influence the prescription through awareness about the rationing of medicines. With the decrease in consumption, it was possible to avoid the lack of these drugs until resupply. The participation of the clinical pharmacist in the management of pharmacotherapy has the potential to improve the efficiency of hospital service resources, especially in a pandemic scenario.

Funding: No funding to declare.

Ethics statement: Not applicable.

Pharmaceutical Services in the perspective of the users of public pharmacies in a municipality of Rio de Janeiro state: gaps that can impact access

Fernanda Lacerda da Silva Machado¹; Danielle Maria de Souza Sérgio dos Santos¹; Vanessa Pinheiro da Silva Fernandes Gervasio¹; Renato Rodrigues de Figueiredo Cruz¹

¹ Universidade Federal do Rio de Janeiro, Macaé, Rio de Janeiro, Brazil

Context: Community engagement is a key element to improve health systems services and in Brazil, it is one of the principles of the Unified Health System (SUS). However, few studies have addressed the perception and knowledge of the users about pharmaceutical services in SUS.

Objectives: To evaluate the perception and knowledge of users about essential medicines lists and the barriers to access treatments in SUS.

Methods: Cross-sectional study conducted using semi-structured interviews with users of public pharmacies managed by the Municipal Secretary of Health in a municipality in the Rio de Janeiro state. The study included adults aged 18 years or more who visited public pharmacies to obtain medicines. The data collection was held during September, December, and March (2019-2020). The study intended to conduct 10 interviews per month at each of the six public pharmacies, with an estimated sample of 180 users. The study protocol was approved by the Research Ethics Committee of the Federal University of Rio de Janeiro- Macaé.

Results: In total, 142 users were interviewed. When asked about medicines from SUS, 71 of them (50.0%) declared they knew the municipal list, however, only 19 (13.4%) answered where it could be consulted. In regard to the barriers to access medicines, 56 users (39.4%) responded that shortage was the main challenge. Additionally, 33 users (23.4%) declared that, in their view, physicians did not mind prescribing treatments available in the public health system and 22 (15.7%) do not feel comfortable asking for that.

Conclusion: Most of the users do not receive proper information about the medicines available in SUS. In their perception, the shortage was the main barrier. However, this may be influenced by the fact that some prescribers and users may not discuss the therapeutic options supplied by the public system.

Funding: Project supported by PROFAEX-UFRJ.

Ethics statement: 4187153.

Use of medicines by elderly people in primary care of the Unified Health System (SUS) in Brazil

Viviane Félix Silveira¹; Renata Cristina Rezende Macedo do Nascimento¹

¹ Federal University Of Ouro Preto, Ouro Preto, Minas Gerais, Brazil

Context: Elderly people often have chronic health problems, which leads to greater use of health services and high consumption of medicines. However, few population-based studies present data on the prevalence of problems related to medicines use in the elderly.

Objectives: This study aimed to assess the profile of medicines use and aspects related to therapeutic safety in the elderly assisted by SUS Primary Health Care (PHC).

Methods: This is a cross-sectional study, part of the National Survey on Access, Use and Promotion of Rational Use of Medicines (PNAUM – Services), with a representative sample of the Brazilian population. The sample consisted of people aged 65 or over, with data collection performed in PHC services, from July / 2014 to May / 2015. Medicines were classified according to Anatomical Therapeutic Chemical, and drug-drug interactions were identified by the Micromedex®. For the classification of potentially inappropriate medication (PIM), the Beers Criteria 2019 was adopted. Polypharmacy was defined as the concomitant use of five or more medicines. Data analysis was performed using SPSS® software.

Results: The study population consisted of 1,157 elderly, with 63.4% (95% CI 59.9-66.7) being female; 58.7% (95%CI 54.1-63.2) married; and 58.8% (95%CI 50.4-66.7) with incomplete elementary education. Most of the elderly (93.86% 95%CI 92.32-95.17) used at least one medicine in the 30 days before to the interview, an average of 2.87 ± 2.12 medicines/elderly. The most used drugs were hydrochlorothiazide (10.55%), losartan (7.96%) and captopril (7.01%). Of the total of identifiable drugs, 22.24% were classified by the Beers Criterion 2019 as PIM for the majority of the elderly, being acetylsalicylic acid (23.39%), omeprazole (18.06%), and glibenclamide (9.84%) most used. 1,226 potential drug interactions were identified, and the prevalence of polypharmacy was 18.78% (95% CI 16.50-21.24), with an association between the use of medicines and worse health indicators ($p < 0.05$).

Conclusion: A high prevalence of polypharmacy, drug-drug interactions, and use of PIM was identified among the elderly assisted by PHC in Brazil, indicating the need to improve health care to ensure quality and safety in medicines use.

Funding: Project supported by Department for Pharmaceutical Services and Strategic Health Supplies and Department of Science and Technology of the Secretariat of Science, Technology and Strategic Supplies of the Brazilian Ministry of Health (SCTIE/MS – Process 25000.111834/2, Decentralization of FNS Resources).

Ethics statement: 398131.

Drug therapeutic prescribed according to initial complaint and sign/symptom to users of the infanto-juvenile psychosocial care center.

Ana Beatriz dos Santos¹; Nevoni Goretti Damo¹

¹ Regional University of Blumenau, Blumenau, Santa Catarina, Brazil

Context: The prescription and consumption of psychiatric drugs by children and adolescents is one of the concerns of health professionals and Brazilian society. Medicines for the treatment of mental disorders and psychological distress are therapeutic resources frequently prescribed to children and adolescents with psychological distress assisted at the Psychosocial Care Centers for children and adolescents (CAPSi).

Objectives: This study sought to describe the therapy prescribed according to the initial complaint and signs / symptoms of children and adolescents at CAPSi; identify the main drugs prescribed; relate the drugs to the initial complaints and/or signs/symptoms; analyze the percentage of prescription drugs present in the National Drug List (RENAME) and the Municipal Drug List (REMUME).

Methods: Descriptive, retrospective study with a quantitative approach with data obtained from physical and electronic medical records of children and adolescents of both sexes, with prescription of medicines during the last medical consultation in the year 2018.

Results: 48 children and adolescents aged between 13 and 18 years, 28 male and 20 female. 85 initial complaints, 97 signs/symptoms were reported. Of the prescription drugs, 68.42% are present in RENAME and/or REMUME. The main medication prescribed as a form of treatment was Risperidone 22.10%. The most common initial complaints were aggression and anxiety 13.40%. The active substances Risperidone, Fluoxetine Hydrochloride, and Escitalopram Oxalate are the main substances prescribed to children and adolescents of CAPSi, which when used in a rational way can benefit the treatment of complaints and signs/symptoms presented in the study. It is essential to highlight the importance of expanding the discussion on prescriptions for drugs not available, such as Risperidone and Escitalopram Oxalate in SUS, as these drugs are prescribed, but are not included in the list of essential drugs, both National and Municipal.

Funding: No funding to declare.

Ethics statement: 2731093.

Pharmacists' role in patient safety: evaluation of dispensing errors in a regional hospital

Isabela Rufo Cordeiro Vieira¹; Amanda Teixeira de Araújo¹; Carolina Silveira Franco¹; Lucas Duarte Silva¹; Lilianne de Fatima Pereira²; Renata Cristina Rezende Macedo do Nascimento¹

¹ Federal University of Ouro Preto, Ouro Preto, Minas Gerais, Brazil

² Santa Casa da Misericórdia de Ouro Preto, Ouro Preto, Minas Gerais, Brazil

Context: In the hospital environment, the medication process is highly complex and can generate circumstances that lead to errors, which in turn, can cause permanent or lethal harm to patients. Understanding the processes of medicines use in hospitals and the determinants of medication errors are essential strategies to mitigate risks.

Objectives: To analyze the medication dispensing errors identified by the pharmacy of a regional hospital.

Methods: This is a cross-sectional study carried out in a medium-sized regional hospital (118 beds) in Minas Gerais state/ Brazil, from July 2019 to June 2020. The dispensing error was defined as a discrepancy between a prescription and the medicines delivered by the pharmacy to the patient or distributed to the ward-based on of this prescription. Data were collected at the time of drug dispensing and immediately recorded in a Microsoft Excel® spreadsheet. Dispensing errors were presented as absolute and relative numbers.

Results: Of the 19,655 prescriptions included in the study period, 380 dispensing errors were identified, an error rate of 1.9%. The period from September 2019 to February 2020 presented the highest rates (3.6% per month). A high prevalence of errors was identified in women (n = 165; 53.4%) and in the elderly aged 65 years or older (n = 140; 36.9%). The most prevalent dispensing errors were dose omission and/or medication omission (n = 165, 42.8%). In relation to therapeutic groups, antibacterial for systemic use (n = 38, 12.6%) was the main pharmacological group related to errors. Considering the hospital sector, the prescription of generalist medications had the highest rate of dispensing errors (n = 115; 30.3%).

Conclusion: Dispensing errors are still frequent in the hospital environment and may indicate quality deviations in the service provided by the pharmacy. It is the responsibility of the pharmacist to minimize them, based on the implementation of barriers and actions that lead to the reduction of the occurrence of flaws in the medication system, in order to ensure patient safety throughout the health-care process.

Funding: No funding to declare.

Ethics statement: Not applicable.

Use of medicines in child mental health: a description

Sandna Larissa Freitas dos Santos¹; Felipe Moreira de Paiva¹; Marinara Fonseca Freire; Maria Eduarda da Silva Costa¹; Ana Paula Soares Gondim¹

¹ Federal University of Ceará, Fortaleza, Ceará, Brazil

Context: Drug utilization research act as a tool to assess the prescription, dispensation and administration of drugs in a given population, providing an overview of therapeutic problems, helping to identify the needs for interventions. In children, the use of psychotropics has become a public health problem, as it is a practice-based mainly on adaptations of use in adults.

Objectives: To describe the use of psychotropic drugs by children in a community mental health service.

Methods: A descriptive and retrospective study, with a quantitative approach, carried out in a child and adolescent psychosocial care center with secondary data from the year 2017. Data collection was carried out by surveying the medical records of the children attended in 2017. The data were stored in the program epi-info 7.0. A simple statistical analysis of the following variables was performed: age, diagnostic hypothesis, amount of psychotropic per child and name of the psychotropic. Study approved by the research ethics committee (number 1,684,517).

Results: Of the total of 395 children who participated in the study, the most frequent age group was 5 to 7 years old with (47.8%), Followed by the 8- to 10-year-old group (35.4%) And the 11 and 12 years (16.7%). 482 Diagnostic hypotheses were identified, including childhood autism (70.5%) And moderate mental retardation (58.9%), With more than one hypothesis. All children used psychotropic drugs, with an average of 1.5 Drugs per child. Among the drugs described, risperidone (51.9%), Valproic acid (34.2%) And methylphenidate hydrochloride (14%) stands out.

Conclusions: Finally, this study shows that it is necessary to develop strategies that promote safe use in the care process with children with possible disorders, in order to minimize the risks inherent to pharmacotherapy in the field of child mental health.

Funding: Project supported by Cearense Foundation to Support Scientific and Technological Development (FUNCAP) (process number: 44976-87).

Ethics statement: 1684517.

Psychotropic medication usage by children referred to centers for psychosocial child and adolescent care

Felipe Moreira de Paiva¹; Sandna Larissa Freitas dos Santos¹; Nívia Tavares Pessoa¹; Maria Eduarda da Silva Costa¹; Marinara Fonseca Freire¹; Ana Paula Soares Gondim¹

¹ Federal University of Ceará, Fortaleza, Ceará, Brazil

Context: The mental health network represents a network of integrating care for mental health in Brazilian population, which center for psychosocial child and adolescent care is responsible for reception and care of children and adolescents with moderate and severe mental disorders. It is a specialized service, which supports multidimensional care, and the medication usage may be a problem when users are referred to these mental services already using psychotropic medication from other services.

Objectives: To describe the frequency of psychotropic medication usage by children referred to centers for psychosocial child and adolescent care of a metropolitan region of the northeast, in Brazil.

Methods: Descriptive and retrospective study, quantitative approach, using secondary data (medical records) of children of 0 to 12 who were referred to two centers for psychosocial child and adolescent care in 2017. Variables analyzed were age, place of origin of referral to the service, psychotropic medication usage before getting in centers, and drug name, with anatomical therapeutic chemical classification (ATC) code. Data were analyzed according to simple descriptive statistics and were stored in the Epi info program, 7th version.

Results: From a total of 752 medical records of children who got in two centers, the average age was 7 (SD 2 for minus and more). The majority local of referrals was basic health unit, 23.3% (149) of referrals, followed by 20.0% (128) from spontaneous demand, 18.4% (117) from hospitals and 15.7% (100) from schools. Over half of the children, 56.8% (364), were referred to centers with psychotropic medication prescribed. Of those medications prescribed, risperidone, ATC code N05AX08, was the most frequently, 56.3% (205), followed by periciazine, ATC code N05AC01, 10.9% (40), and valproic acid, ATC code N03AG01, 7.7% (28).

Conclusion: Even though most children are referred from basic health units, it is worrying the data above more than half of children getting into specialized service already using psychotropic medication, mainly risperidone. According to this information, we may discuss the necessity of premature exposure of children by psychotropic medication even before going to the specialized services.

Funding: Project supported by Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP).

Ethics statement: 168417.

Profile of use of ceftazidime/avibactam in ICU patients in a tertiary hospital

Alice Silva¹, Luis Phillpe Nagem¹; Camila Oliveira¹; Diamantino Ribeiro Salgado²; Elisangela da Costa Lima¹

¹ Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

² Clementino Fraga Filho University Hospital, Rio de Janeiro, Brazil

Context: Ceftazidime/avibactam is a new antibiotic for treating specific hospital-acquired infections, registered in Brazil in 2019. Its empirical use and for community infections is not recommended.

Objectives: To assess ceftazidime/avibactam use in the Intensive Care Unit (ICU) of a private hospital in Rio de Janeiro.

Methods: A cross-sectional descriptive study was conducted (May 2019-May 2020), which included all patients admitted to the ICU who used systemic antimicrobials for more than 48 hours. Exclusion criteria were not applied. The use was classified as empirical (without proven sensitivity in vitro) or guided (based on sensitivity analysis). Dose, treatment duration, pathogen, and infectious focus were assessed. The observed use was compared to that described in the package insert. Ethics and research committee approved this research.

Results: 578 patients were included, of which 21 (3.6%) had 27 infections treated with ceftazidime/avibactam: pneumonia (n=21; 77.8%) urinary tract (n=2; 7.4%), primary bloodstream Infections (BSI) (n=2; 7.4%), cellulitis (n=1; 3.7%) and osteomyelitis (n=1; 3.7%). There was empirical use in six cases (22.3%) to ventilator-associated pneumonia. In culture-guided treatments, all bacteria identified were multidrug-resistant. In 17 cases, *Pseudomonas aeruginosa* was only sensitive to ceftazidime/avibactam. In four, KPC carbapenemase-producing *Klebsiella pneumoniae*, sensitive to ceftazidime/avibactam and aminoglycosides, were identified. Median (interquartile range) of antibiotic use time was 7 (6-13) days and doses equal to 2.5G every eight hours. Use was guided by cultures for BSI, cellulite, and osteomyelitis. However, there was no indication (14.8%) for these cases. No deaths or adverse reactions were observed in 28 follow-up days.

Conclusion: This real-life study showed the infrequent use of ceftazidime/avibactam in the investigated hospital, mainly guided by culture and indications consistent with previous clinical studies. Nevertheless, empirical and off-label use was observed for one-third of cases. It suggests the need for new studies and discussion of strategies to rationalize new antimicrobials.

Funding: Federal University of Rio de Janeiro supported a scientific initiation scholarship.

Ethics statement: 3924840.

Investigation of anti-infectives adverse reactions in hospitalized children in a university institute in the city of Rio de Janeiro

Thais de Barros Fernandes¹; Elisangela da Costa Lima²

¹ Sergio Arouca National School of Public Health, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

² Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

Context: Children are more susceptible to adverse drug reactions (ADR) comparing to adults. Anti-infectives are one of the most prescribed classes and more related to the occurrence of these ADR in pediatrics. Analyzes of causality, avoidability, and intensity of ADR are important to establish the likelihood that a suspicious drug will cause a reaction, develop actions to prevent it, and conduct appropriate clinical management.

Objectives: An observational study was developed with prospective data collection (May-October 2019) on the suspicions of ADR to anti-infectives in children aged 0 to 17 years hospitalized at a teaching hospital. The cases were assessed for causality and avoidability (Liverpool ADR assessment tools and intensity with the Hartwig scale. The data were also used to calculate the overall incidence and risk factors. Descriptive statistics were used and values with $p < 0.05$ were considered significant.

Results: 171 patients were included in the study. The children with more ADR belong to the age group between 0 and 6 years, are female, used more than one anti-infectious, and remained hospitalized for more than a week. The incidence of ADR was 5.46% and 5.84%, considering the numbers of patients and admissions. The drugs most related to these reactions were cefepime and oxacillin. The gastrointestinal tract and the skin were the most affected systems. 26.9% reactions are defined and 15.4% are likely as causality. 81.8% of the reactions were possibly preventable and 18.2% were unavoidable. The reactions were considered mild in 54.5% of the cases, moderate in 36.4%, and severe in 9.1%. Half of the ADR required the administration of other medications to contain the symptoms of the reaction.

Conclusion: The study suggested children who use more drugs and children who use drugs for a long time are more likely to develop ADR.

Funding: No funding to declare.

Ethics statement: 3264238.

Anticoagulant and antiplatelet therapy in older patients: a Brazilian consensus of potentially inappropriate prescribing indicators

Fabiane Raquel Motter¹; Sarah Nicole Hilmer²; Luciane Cruz Lopes¹; Vera Maria Vieira Paniz³

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² University of Sydney, Australia

³ Univerity of Vale do Rio dos Sinos, São Leopoldo, Rio Grande do Sul, Brazil

Context: Potentially Inappropriate prescribing (PIP) of antiplatelets and anticoagulants among older patients increases the risk of adverse outcomes such as bleeding and drug-related hospitalizations, and death in this population. Definition of PIP indicators in the Brazilian geriatric population could be extremely useful for healthcare professionals in order to improve the management of these patients and to avoid preventable adverse events.

Objectives: To develop and validate Potentially Inappropriate prescribing (PIP) indicators for use of antiplatelets and anticoagulants medications in older people adapted to the Brazilian context.

Methods: A preliminary PIP indicators suitable for the Brazilian market was developed on the basis of three published international PIP indicators [Beers 2015, Screening Tool of Older People's Potentially Inappropriate Prescriptions - 2015, European Union (7) PIM list]. We used the modified Delphi technique (three-round) to validate concerns of use of antiplatelet and anticoagulant medications in older adults aged 65 years or more in Brazil. The panel involved seven Brazilian experts in geriatric pharmacotherapy. All items with mean Likert scale score ≥ 4.0 (agree) and the lower limit of 95% confidence interval ≥ 4.0 were considered validated in this study.

Results: At the end of the consensus process, 57 (65.3%) items of 61 were validated. In total, consensus was reached for 8/8 (100.0%) concerns about drugs that should be avoided in older patients regardless of diagnosis, for 16/17 (95.7%) concerns about drugs that should be avoided in older patients with specific conditions or diseases, and for 15/15 (100.0%) with special considerations of use.

Conclusion: Although these indicators are not designed to replace clinical judgement, PIP indicators can be useful to inform prescribers, pharmacists, and health care planners and may serve as a starting point for safe and effective use of medications in older people.

Funding: Project supported by CNPq (426720/2016-4).

Ethics Statement: 13773012.

Interventions for depression and anxiety in diabetes mellitus patients: overview of systematic reviews and meta-analyses

Reginaldo Tavares Franquez¹; Isabela Muniz de Souza¹; Cristiane de Cássia Bergamaschi¹

¹ University of Sorocaba. Sorocaba, São Paulo, Brazil

Context: Diabetes mellitus (DM) and depression are common public health problems in the world that are causing a significant negative impact on physical, psychological, social, and occupational functioning as well as increasing economic burdens.

Objectives: This overview of systematic reviews (SR) summarized the available scientific evidence regarding the effectiveness and safety of interventions to treat depression and other psychological symptoms in patients with diabetes mellitus type 1 (T1) and 2 (T2).

Methods: The following sources of information were searched without restriction on language and year of publication: Cochrane library, Medline, Embase, Web of Science, and Lilacs. Interventions to treat depression and/or anxiety in patients with DM were compared with placebo or standard interventions. The primary outcomes were an improvement or clinical remission of depression and anxiety and improvement in the quality of life. Secondary outcomes were reduction in HbA1c values, treatment costs, death from any cause, adherence to treatment for diabetes, and complications of this disease. Reviewers, in pairs and independently, selected the studies, extracted their data, and assessed the quality of the studies using the AMSTAR 2. The narrative synthesis of the findings was carried out.

Results: This study found 15 SR with 55,213 participants. Pharmacological interventions, collaborative care, and health education improved the depression and glycemic control in adults with T1 and T2 DM compared to the controls. Cognitive behavior therapy improved depression and glycemic control in adults and elderly with T1 and T2 DM. Health education was superior to usual care in remission of depression and glycemic control in adults with T1 and T2 DM, but not for web-based health education. Structured education improved psychological distress in adults with T1DM, but not symptoms of depression. Diet and/or physical activity decreased depression scores in adults with T2DM. Most of the studies had methodological problems.

Conclusion: The interventions discussed in this study seemed to be effective and safe. However, the quality of the evidence obtained from these findings indicated that future randomized controlled trials are needed to confirm this. This study can guide patients and health professionals in making safe and effective decisions concerning the technologies used to care for patients.

Funding: Project funded by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Prevalence of *Enterococcus faecium* resistant in poultry farming: systematic review

Stephanie Valerio Silva¹; Anna Luiza Crescenzo Brotto¹; Felipe Bernardini Antoniazzi¹; Júlia Módolo Stievano¹; Renata Lopes Simões da Silva¹; Cristiane de Cássia Bergamaschi¹; Marcus Tolentino Silva¹; Silvio Barberato-Filho¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Antimicrobials are used in poultry farming to prevent and treat diseases caused by bacteria and to reduce the effects caused by confinement stress. However, the indiscriminate use of antimicrobials may be related to increased bacterial resistance, not just in animals, but in humans too.

Objectives: This study determined the prevalence of resistant *Enterococcus faecium* in poultry.

Methods: This systematic review sought eligible studies from the following sources of information: Scopus, Web of Science, Medline, and Embase, without language restrictions. The outcome of interest was the prevalence of resistant enterococcus faecium. The geographic distribution of resistant microorganisms and the time course of resistance trends over the past ten years were described. The prevalence of resistance was grouped by the random model of DerSimonian and Laird.

Results: Twenty-four studies were included, mainly carried out in European countries (n=9), in production (n=11), slaughter (n=4) and retail (n=9) environments. The prevalence of resistant *Enterococcus faecium* in poultries was 18.4% (95% CI: 17.0 to 19.4%), being higher in Africa (40.6%; 95% CI: 34.6 to 46.8%). There was higher prevalence in samples from production environments (24.8%; 95% CI: 23.3 to 26.4%), and in poultry feces (24.2%; 95% CI: 22.7 to 25.6%). There was no significant variation in prevalence in the temporal course ($p>0.525$). In the resistance investigation the most tested antibiotics were vancomycin (in 54% of the evaluated sample it presents resistance), erythromycin (84%), ciprofloxacin (95%), tetracycline (95%), and gentamicin (50%).

Conclusion: The prevalence of resistant *Enterococcus faecium* in poultry was higher in Africa, with no significant variation in prevalence over time. A higher percentage of resistance was observed with tetracycline and ciprofloxacin, antibiotics also routinely used in human medicine. The results of this study reinforce the need to plan interventions that can reduce the use of antimicrobials in poultries for food production.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

Prevalence of *Enterococcus faecium* resistant in commercial cattle: systematic review

Anna Luiza de Crescenzo Brotto¹; Stephanie Valerio Silva¹; Renata Lopes Simões da Silva¹; Felipe Bernardini Antoniazzi¹; Julia Módolo Stievano¹; Silvio Barberato-Filho¹; Marcus Tolentino Silva¹; Cristiane de Cássia Bergamaschi¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: The use of antimicrobials in animals is broader compared to humans, which has been identified as one of the main causes of the emergence of bacterial resistance. Data on the resistance profile in cattle are scarce in the literature.

Objectives: This study determined the prevalence of resistant *Enterococcus faecium* in commercial cattle.

Methods: This systematic review sought eligible studies from the following sources of information: Scopus, Web of Science, Medline and Embase, without language restrictions. The outcome of interest was the prevalence of *E. faecium* resistant. The geographic distribution of resistant microorganisms and the time course of resistance trends over the past ten years were described. The prevalence of resistance was grouped by the random model of DerSimonian and Laird.

Results: Eighteen studies were included, mainly carried out in European countries (n= 9) and in production (n= 11) and retail (n= 7) environments. The main material used in the detection of the microorganism was the milk. The prevalence of *E. faecium* resistance in cattle was 4.3% (95% CI= 2.8–5.0%) being higher in Asia [25.4% (95% CI= 20.5-30.6%)]. There was higher prevalence in samples from retail [13.7% (95% CI= 11.5-16.1%)], collected mainly from equipment and surfaces [12.5% (95% CI= 5.5-26.1%)]. There was no significant variation in prevalence in the temporal course (p> 0.05). The antibiotics most tested were vancomycin, tetracycline, ciprofloxacin, and erythromycin, with resistance of 50%, 59%, 79%, 94%, respectively.

Conclusion: The prevalence of *E. faecium* resistant in cattle was higher in Asia, with no significant variation in prevalence over time. A higher percentage of resistance was observed with erythromycin and ciprofloxacin, antibiotics also routinely used in human medicine. The results of this study reinforce the need to plan interventions that can reduce the use of antimicrobials in cattle for food production.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

Efficacy and effectiveness of mobile health technologies to reduce alcohol or tobacco consumption during pregnancy: scoping review

Laura Inês Gomes Maricato¹; Alanis Fernanda Proença Montrezol dos Santos¹; Maria Raquel Gomes Fernandes²; Livia Luize Marengo¹; Silvio Barberato-Filho¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² São Paulo State University, Botucatu, São Paulo, Brazil

Context: More than 20% of pregnant women worldwide consume alcohol and 1.7% smoke during pregnancy. Mobile health technologies help to improve the monitoring of health conditions and allow patients to take a more active role in their care, in addition to ease of use, acceptance, and accessibility.

Objectives: This study summarized the evidence on the efficacy and effectiveness of mobile health technologies to reduce alcohol or tobacco consumption during pregnancy.

Methods: This scoping review sought eligible studies from the following sources of information: Medline; Excerpta Medica Database; Cumulative Index to Nursing and Allied Health Literature; the Cochrane library; Web of Science; Scopus; PsycInfo and Virtual Health Library, without language restrictions. Randomized controlled trials examining the effectiveness of mobile health technologies for reducing consumption of alcohol or tobacco among pregnant women were included. The extracted data were analyzed for the type of mobile technology, population, intervention, comparators, and outcomes.

Results: Thirteen studies were included: ten addressing smoking alone and three simultaneously addressing the reduction of alcohol and tobacco consumption during pregnancy. In primary studies, 3,603 volunteers were randomized. Among the included studies, ten were carried out in the United States (77%) and three in Europe (23%). The short message service was the most frequent intervention (77%). Only three studies (23%) were able to support the benefits of mobile technologies, and all involved the same intervention.

Conclusion: Large heterogeneity in interventions, comparators, and outcomes, in addition to methodological weaknesses limited our ability to make assertive conclusions relative to the overall effectiveness of mobile technologies. While promising, further research is necessary to determine mobile technologies' benefits in reducing alcohol and tobacco consumption during pregnancy.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

Wearable devices designed to improve medication adherence: a systematic map

Livia Luize Marengo¹; Julia Hiromi Hori Okuyama¹; Alan Martinez Kozyreff¹; Maria Raquel Gomes Fernandes²; Reginaldo Tavares Franquez¹; Isaltino Pereira de Andrade Junior¹; Marcus Tolentino Silva¹; Silvio Barberato Filho¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² São Paulo State University, Botucatu, São Paulo, Brazil

Context: Poor medication adherence is relatively common. About 20 to 30 percent of drug prescriptions are not compliant, reaching 50 percent for chronic diseases. Wearable devices were designed to improve medication adherence and alert patients to noncompliance. These devices explore everything from detecting the opening of a pill bottle to smart necklaces with sensors to identify whether the drug has been swallowed, issuing audible and vibrating alerts, or sending text messages to users.

Objectives: In this systematic map, wearable devices designed to improve medication adherence were investigated.

Methods: Studies that evaluated wearable devices or described the development of these devices to improve medication adherence were included. Wearable devices that did not allow patients' autonomy in decision-making regarding medication adherence (or not) were excluded. The search was completed in March 2020 in Medline, Embase, Scopus, CINAHL, PsycInfo, and Web of Science databases. Reviewers, in pairs and independently, screened the titles and abstracts and applied the eligibility criteria to the full texts. The extracted data were recorded in an electronic spreadsheet and the results were presented in horizon scanning maps and narrative description.

Results: A total of 7,087 records were identified in the databases and 27 studies were included, of which 24 were published between 2013 and 2020. The studies involved smartwatches (n=9), patches (n=6), wristbands (n=3), another isolated wearable technology (n=5) and technology applicable to various wearable devices (n=4). Most studies (n=18) addressed specifications for the technical development, six of which related to smartwatches. Thirteen studies evaluated wearable devices from the users' perspective, assessing acceptability (n=5), usability (n=3), device adherence (n=3), and satisfaction (n=2). In 12 studies the results were favorable to the wearable device. Only one study on patient satisfaction showed a negative outcome.

Conclusion: The growing number of studies published in recent years demonstrates advances in the development and evaluation of wearable devices. While promising, further research is necessary to determine wearable technologies' benefits in medication adherence.

Funding: Project funded by the University of Sorocaba and by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Efficacy and safety of repellents marketed in Brazil against bites from *Aedes aegypti* and *Aedes albopictus*: a systematic review

Maria Raquel Gomes Fernandes¹; Luciane Cruz Lopes²; Rodrigo Sugimoto Iwami²; Mariana Del Grossi Paglia²; Bruna Mateus de Castilho Soleder³; Alan Maicon de Oliveira⁴; Izabela Fulone²; Ricardo Silveira Leite²; Cristiane de Cássia Bergamaschi²

¹ São Paulo State University, Botucatu, São Paulo, Brazil

² University of Sorocaba, Sorocaba, São Paulo, Brazil

³ São Paulo State University, Araraquara, São Paulo, Brazil

⁴ University of São Paulo, Ribeirão Preto, São Paulo, Brazil

Context: Dengue, Zika and Chikungunya viruses represent a serious public health problem. No evidence is available on the efficacy of repellents commercially available in Brazil.

Objectives: This systematic review assessed the efficacy and safety of products containing repellents commercially available in Brazil for protection against bites from *Aedes aegypti* and *Aedes albopictus*.

Methods: We performed a systematic review using the Central, Medline, Embase, Cinahl, Web of Science, Amed, Lilacs and Scopus databases. Randomized clinical trials (RCTs) and non-randomized clinical trials (NRCTs) comparing topical repellent products registered with the Brazilian Health Surveillance Agency (ANVISA) were included. Outcomes of interest included: protection time (PT), percentage of repellence (%R), and adverse effects (AE). Pairs of reviewers selected the studies, extracted the relevant data, and evaluated the risk of bias.

Results: Sixteen studies (5 RCTs and 11 NRCTs) evaluating the outcomes PT, %R and AE were retrieved. Against *A. aegypti*: IR3535 (15%-spray and 10%-solution) provided similar %R than Icaridin (20%-spray and 10%-solution). Considering PT outcome, DEET (10% and 20%-spray) was similar to IR3535 (10% and 20%-spray) and longer than Citronella (5%-spray); DEET (25%-solution) had longer PT than Eucalyptus (25%-solution), while DEET (20%-lotion) had longer PT than Citronella (10%-lotion). There was no difference considering PT between Eucalyptus (10%-oil) and Citronella (10%-oil) or Citronella+Eucalyptus (5%-oils). Against *A. albopictus*: DEET (15%-spray) was similar to Icaridin (20%-spray) and longer than Citronella (10%-spray). No AE were reported (n=1).

Conclusion: These results suggest that DEET has greater efficacy than other synthetic and natural repellents marketed in Brazil.

Funding: Project funded by the São Paulo Research Foundation (FAPESP), grant n° 2017/07813-8.

Ethics statement: Not applicable.

Antimicrobial resistance in periodontal infections (MICROBE-DENT): systematic review

Flávia Casale Abe¹; Cristiane de Cássia Bergamaschi Motta¹; Silvio Barberato Filho¹; Katia Kodaira¹; Luciane Cruz Lopes¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: The lack of knowledge about the microbiota present in periodontitis might lead to the selection of an antibiotic for intrinsically resistant pathogens, compromising the effectiveness of the treatment and increasing the probability of a clinic developing antimicrobial resistance. The antibiotic resistance crisis is occurring worldwide, and bacterial infections are once again a threat.

Objectives: To identify and to characterize the circulating bacteria isolated from periodontal infections as well as their antibiotic resistance profile.

Methods: This systematic review included case series. Inclusion criteria were as follow: patients with untreated periodontal infections. The protocol was registered in PROSPERO (CRD42018077810). The bibliographic search included all publications without language restrictions from January 2013 to June 2020 in the following electronic databases: Ovid Medline, Ovid Embase, BVS, CINAHL, Web of Science, Cochrane central register of controlled trials. The search for gray literature was also included. “The Joanna Briggs Institute” tool was used to assess the risk of bias in the included studies. The findings were summarized in descriptive analysis.

Results: Eight studies were included (n=1670 patients, 870 with periodontal disease) that evaluated the antibiotic resistance of 41 bacteria species into aggressive or severe periodontitis and chronic or unspecific periodontitis. Among the reported infections the antibiotics that showed the highest resistance rates (100%) were ampicillin, cefotaxime, dicloxacillin, metronidazole and penicillin in the presence of *Aggregatibacter actinomycetemcomitans* and *Staphylococcus aureus*. The resistance genes found were *blacpa*, *blacfxa*, *blashv*, *blatem*, *ermb*, and *ermf*. In both infection, virulence factors that contribute to increase antimicrobial resistance have also been reported. The level of confidence in the quality of the studies was critically low. The inconsistency of the data extracted from the articles did not allow meta-analysis pooled estimate.

Conclusion: All bacteria found were resistant to at least one of the tested antibiotics. Studies with better design and robust methodology including genes identification are required to check the prevalence of antimicrobial resistance in periodontal infections in Latin American countries.

Funding: Project funded by a scholarship from the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Critical appraisal of clinical practice guidelines for the treatment of urinary incontinence: a systematic review

Flávia Blaseck Sorrilha¹, Lauren Giusti Mazzei¹, Luciane Cruz Lopes¹, Silvio Barberato-Filho¹, Juliana Castro¹, Anaura Castro¹, Claudia Marcela Vélez², Cristiane de Cássia Bergamaschi¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² University of Antioquia, Medellín, Antioquia, Colombia

Context: All around the world, urinary incontinence is a common complaint, a cause of distress, as well as a source of significant costs for individuals and society.

Objectives: This systematic review evaluated the rigor of the development of clinical practice guidelines (CPG).

Methods: The searched sources were Medline, Embase, Web of Science, Scopus, and specific databases of CPG. Reviewers, organized in triplicate and independently, selected the studies and assessed the quality of the guidelines using the Appraisal of Guidelines Research and Evaluation (AGREE II) instrument, which contains six domains for classification. The classification of the CPG prioritized the domain 3 (Developmental Rigor) considering: high (score >60%), moderate (score 30-60%) or low quality (score <30%). The results were checked for discrepancies and decided by consensus. The interventions were described. Descriptive statistics presented the results.

Results: Of the ten CPG evaluated, five were of high methodological quality, three were of moderate quality, and two were of low quality. Three documents were not recommended for use. The domains with the highest scores were Scope and Purpose (mean=90.1%), and Clarity of Presentation (mean=88.9%). The domains of Editorial Independence (mean=41.4%) and Applicability (mean=29.3%) were those with the lowest score. The most cited interventions in CPG were the non-pharmacological, such as lifestyle interventions, bladder training, or re-education and pelvic floor muscle training.

Conclusion: Most guidelines showed rigor in development and were recommended for use; however, editorial independence and applicability were domains that needed to be improved in these documents. Our findings can guide the choosing of CPG for the treatment of urinary incontinence.

Funding: No funding to declare.

Ethics Statement: Not applicable.

Development of a nursing care guideline for patients with external ventricular drainage

Tainara Wink Vieira¹, Victória Tiyoko Moraes Sakamoto¹, Luiza Casais de Moraes¹, Carine Raquel Blatt¹, Rita Catalina Aquino Caregnato¹

¹ Federal University of Health Sciences of Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil

Context: The External Ventricular Drainage (EVD) is a diagnostic and therapeutic technology frequently used in critical neurosurgical patients, which requires knowledge from the nursing team. Guidelines are used to qualify assistance and communication in the work process, and to promote safer care.

Objectives: To develop a nursing care guideline to promote the safety of adult patients undergoing external ventricular drain assisted in intensive care units.

Methods: This is a technological development in health research carried out in three stages: (a) scoping review; (b) evaluation of the quality of the evidence; and (c) development of the nursing care guideline according to the recommendations of Pimenta et al. and the Appraisal of Guidelines Research and Evaluation (AGREE II).

Results: The scoping review stage resulted in the selection of 54 articles from different information resources that served as a basis for determining the main nursing care. Each of them was evaluated according to the Grading of Recommendations Assessment, Development and Evaluation (GRADE), resulting in 2 studies with high quality, 13 with moderate quality, 34 with low quality and 5 with very low quality. The guideline contains flowcharts that help in decision-making and presents 20 nursing interventions with their respective justifications. Such care was subdivided into five categories, namely: positioning and mobilization in bed; drainage system; catheter care; monitoring of intracranial pressure; and medication administration.

Conclusion: Assistance protocols are health technologies whose complexity varies according to the different demands related to care. Such complexity encourages nursing professionals to develop and improve their critical thinking, promoting increasingly qualified and safe care. Nurses are directly involved in this process, and they are essential to promote best care practices with the external ventricular drainage, therefore, tools must be developed and implemented to support and strengthen safe practices.

Funding: No funding to declare.

Ethics statement: Not applicable.

Prevalence of *Enterococcus faecium* resistant in commercial aquaculture: systematic review

Renata Lopes Simões da Silva¹; Anna Luiza de Crescenzo Brotto¹; Felipe Bernardini Antoniazzi¹; Júlia Módolo Stievano¹; Stephanie Valério Silva¹; Cristiane de Cássia Bergamaschi¹; Silvio Barberato-Filho¹; Marcus Tolentino Silva¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: The use of antibiotics is a routine in livestock. With it, the control of bacterial diseases that affected production became possible, but this has increased antimicrobial resistance in these environments in recent years. Among the microorganisms with interest in public health, *Enterococcus faecium* stands out, associated with bacteremia, endocarditis, urinary tract infections, and wounds. Data on antimicrobial resistance related to *Enterococcus faecium* in aquaculture are scarce.

Objectives: To estimate the prevalence of resistance of *Enterococcus faecium* in aquaculture.

Methods: A systematic review, in which the sources of information consulted were Scopus, Medline, Embase, and Web of Science. The reviewers, independently and in pairs, selected the studies according to the eligibility criteria and performed the data extraction. The time course of resistance trends, potential sources of contamination, and the geographic distribution of resistant microorganisms were described. The prevalence of antimicrobial resistance was grouped by the random model of DerSimonian and Laird and the double arcsene transformation proposed by Freeman-Tukey.

Results: 12 articles were selected. One in every ten products showed *Enterococcus faecium* resistance (8.7%, 95% CI: 7.6-9.9%), being the European continent (13.7%) and the breeding environments (15.8%) with the highest prevalence rates and with a slightly grow in the last 10 years. It was also noted that the bacterium was resistant to many important antibiotics, such as kanamycin, clindamycin, and streptomycin.

Conclusion: *Enterococcus faecium* presented a high prevalence of resistance to antibiotics that are used by human and animal medicine being mainly found in European continent and in breeding environments. This result suggests that regulatory measure of the use of antibiotics in aquaculture, aiming to reduce the risk of the development and dissemination of superbugs are required.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

The incorporation of biological medicines into the Brazilian Unified Health System (SUS), 2000-2019

Kaite Cristiane Peres¹; Fabíola Bagatini Buendgens²; Claudia Marcela Vargas-Pelaez³; Mareni Rocha Farias¹

¹ Federal University of Santa Catarina, Florianópolis, Santa Catarina, Brazil

² Santa Catarina State Health Secretariat, Florianópolis, Santa Catarina, Brazil

³ National University of Colombia, Bogotá D. C., Colombia

Context: Brazil is among the ten largest consumers of pharmaceutical products in the world, mainly medicines. Among the drug policy strategies adopted by the World Health Organization, we place emphasis on the list of essential medicines, called the National List of Essential Medicines (RENAME). Among the categories of essential medicines, we draw special attention to biological medicines, which over the last decades have gained space in the market with high prices and have created a dilemma in public policy making and management in terms of the sustainability of health systems.

Objectives: To analyze the chronological evolution of the incorporation of biological medicines into the Brazilian Unified Health System (SUS) from 2000 to 2019.

Methods: We categorized the biological medicines according to Resolution RDC55/2010 and associated the presence of these medicines with the RENAME list and the List of Exceptional Medicines. To facilitate the analysis, all medicines were classified according to Pharmaceutical Assistance components (namely CBAF, CEAF, CESAF).

Results: In 2000, 43 active ingredients had been incorporated into the SUS, and 70 in 2010. Regarding medicine availability, it was observed that in the CBAF, no other biological medicines were included after 2010, being noteworthy that, in this component, synthetic medicines with low acquisition values predominate compared to biological medicines. Conversely, in the CESAF and CEAF there was an increased number of incorporated medicines. In the CESAF, this situation has been ongoing since 2000, probably because it encompasses the medicines provided for the National Immunization Policy. Vaccines, serums, and immunoglobulins represent 46.1% (n=53) of the category of biological products available. As for the medicines available within the 2000-2019 period, most technologies remained on the medicines list, evincing the predominant inclusion of technologies into the health system. One limitation of this study is that it did not consider clinical protocols.

Conclusion: Since 2000, the incorporation of biological medicines into the SUS has been increasing with the objective of covering the predominant diseases of the population, and after 2015, it has shown significant importance for the treatment of rare diseases by covering the different care service lines.

Funding: Project funded by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Equity in health and COVID-19: bibliometric approach of impact publications

Renata Maria Santos de Freitas¹; Rílva Lopes de Sousa Muñoz¹; Eduardo Sérgio Soares Sousa¹

¹ Federal University of Paraíba, João Pessoa, Paraíba, Brazil

Context: The 2019 new coronavirus disease pandemic (COVID-19) revealed the close interconnections between socioeconomic inequality and health inequities in the global population. COVID-19 is disproportionately affecting the poor, minorities, and a wide range of vulnerable populations due to its uneven spread and disparities in health access.

Objectives: The aim of this study was to map scientific output on health equity during the COVID-19 pandemic using a bibliometric approach to impact publications in 2020.

Methods: This was a bibliometric and text mining study with a quantitative approach. A search was carried out on the Web of Science database (Web of Science Core Collection) to include high impact articles using the keywords “Health Equity” OR “Vulnerable Populations” AND “Health Economics Agents” AND “COVID-19”, being refined by period (January 2020 to January 2021) and fields (title, summary, keywords).

Results: After exclusion by title and abstract, 100 articles were retrieved, published in 53 journals, and written by 454 authors linked to 253 institutions, mainly Harvard University, Johns Hopkins University and University of California, with 98 publications in English, 59 original articles, mainly from the research areas of public environmental occupational health (49) health care sciences services (16) and general internal medicine (13). The sum of the number of citations was 365, with an average of 3.65 citations per publication and h-index 11. The publications were from 25 countries, 71 in the USA, followed in descending order, England (10), Australia (7), Canada (7), China (5), Belgium (4), Brazil (4) and France (4) and 12 other countries with lower frequencies. The analysis of the keywords showed the following profile: Health, COVID-19, equity, social disparities pandemic, coronavirus, global policy, community determinants public, race, racism, disease, justice, populations research, ethics, infectious, lockdown, primary care access and behavior.

Conclusion: Most of the articles are empirical studies with a general tendency for themes of public, environmental, and occupational health, ethnic diversity, and social disparities in health access. Research in this field can shed light on the current health situation and future health challenges. There was a lack of theoretical studies on equity and the pandemic. The USA had a strong influence in the polls, with absolute leadership. Bibliometric analyzes on COVID-19 and equity are new and vital, as they will provide guidance on public health, health management, health economics and for future publications.

Funding: No funding to declare.

Ethics statement: Not applicable.

Access to innovative biological medicines in Brazil: an analysis of FDA and ANVISA drug approvals

Rafael Augusto Mantovani Silva¹; Daniela Oliveira de Melo¹

¹ Federal University of São Paulo, São Paulo, São Paulo, Brazil

Context: The registration process for a new drug takes time due to its complexity. Most drugs are registered first by the Food and Drug Administration (FDA). The FDA, since 1992, has instituted an accelerated approval process, prioritizing the evaluation of medicines to treat unmet and/or rare medical conditions. The National Health Surveillance Agency (ANVISA) has followed this trend since 2017. However, ANVISA is still criticized for its long registration process.

Objectives: To evaluate if Brazilians have access to innovative biological medicines and whether there has been a change in the approval time of these medicines by ANVISA.

Methods: Innovative biological medicines registered by FDA between 2000-2020 were extracted. Then, their registrations were checked at ANVISA, the time between the registration application and its approval, as well as between the registration by both agencies. Biosimilar drugs were excluded from the analysis.

Results: 128 drugs registered by FDA were identified. Of these, 83 were registered by ANVISA. The other 45 drugs, 26 were registered by FDA between 2018-2020. The mean time between FDA and ANVISA registrations for 126 drugs was 26 ± 33 months (ranging from less than one month to 182 months). The average number of registration approvals at ANVISA was 4/year, peaking in 2017 with 14 approvals. The average time between registration application and approval was 13 months (± 7 months), ranging from 1 month (pegfilgrastim and laronidase) to 36 months (romiplostim). Since 2013, the time between application and approval at ANVISA has been decreasing, with an average of 10 months in the last 3 years.

Conclusion: Brazilians have had access to most innovative biological medicines registered by FDA even though the time between registration in the US and Brazil varies widely. The approvals number has been increasing and its process time has decreased over time, particularly in the last decade.

Funding: No funding to declare.

Ethics statement: Not applicable.

Health technology assessment study on effectiveness and safety of OK-432 (picibanil) in cervical lymphatic malformations

Hérica Núbia Cardoso Cirilo¹; Marcela Maria Faria Peres Cavalcante¹; Marta Curado Carvalho Franco Finotti¹

¹ Federal University of Goiás, Goiânia, Goiás, Brazil

Context: OK-432 (Picibanil) is a biological preparation with sclerosant properties produced in Japan. A HTA report was required to support the use of OK-432 in a pediatric inpatient with cervical lymphatic malformations (LM) in a Brazilian teaching public hospital.

Objectives: To conduct a HTA study to evaluate the effectiveness and safety of OK-432 in cervical LM, under the perspective of a public hospital.

Methods: Evidence synthesis studies were searched in Medline, Cochrane and Epistemonikos. Recommendation reports were consulted at CONITEC, NICE and EUnetHTA. The search took place in January/2021 and considered all published studies. Inclusion criteria consisted of studies relating to cervical LM.

Results: Six systematic reviews and one case report were included. Data analysis demonstrated that most patients achieved complete/excellent lesion reduction. In studies that differentiated the type of lesion, macrocystic ones showed superior results. Recurrence rate has been assessed in some studies, ranging from 2 to 25%. Rates of complications and adverse reactions were low. In most studies, there was no comparison between OK-432 and other technologies. Shock and anaphylactoid symptoms may occur according to manufacturer. OK-432 is not registered by ANVISA, however, its import and use in Brazil is authorized. Cost analysis was not carried out, due to the lack of information regarding the direct cost of OK-432 in official government price banks.

Conclusion: Sclerotherapy with OK-432 in LM proved to be potentially safe and effective, with better response in macrocystic lesions. The results represent the current state of the literature and is a summary of the evidence available for therapeutic decision making in clinical practice. Consideration should be given to the low quality of the primary evidence in the included studies. Until stronger evidence is available, the recommendation is in favor of using OK-432 as a first-line, effective and safe treatment for cervical LM.

Funding: No funding to declare.

Ethics statement: Not applicable.

Expectation for the development of health technology assessment in Brazil

Julia Simoes Correa Galendi¹; Carlos Antonio Caramori²; Clarissa Lemmens¹; Dirk Müller¹; Stephanie Stock¹

¹ University of Cologne, Cologne, North Rhine-Westphalia, Germany

² São Paulo State University, Botucatu, São Paulo, Brazil

Context: In Brazil, Health Technology Assessment (HTA) is mandatory for incorporating new technologies into the benefits catalogue of the national healthcare system (SUS). Compared with international practice, there are still challenges to the application of the HTA processes.

Objectives: To investigate (i) experts' expectations for future development of HTA, (ii) to derive measures to strengthen HTA in Brazil and its impact on decision-making.

Methods: An online survey was conducted among Brazilian experts. Based on a scoping literature review and expert interviews, a questionnaire was developed, including seven domains: (i) Capacity building, (ii) Public involvement, (iii) Role of cost-effectiveness analysis (CEA), iv. Institutional framework, v. Scope of HTA studies, vi. Methodology of HTA, and VII. HTA as basis for jurisdiction. Results of Likert scales were presented as means and mode.

Results: Thirty experts completed the questionnaire. Experts agreed with several actions to promote HTA within the scope of each domain and to five of the eight theses proposed. Main challenges relate to low funding availability and shortage of capacity. Experts differed regarding the theses on the domains 'public involvement' and 'institutional framework'.

Conclusion: The implementation and expansion of HTA processes must be shaped to the characteristics of the SUS and the availability of resources. Capacity building related to HTA was identified as a central domain to develop HTA, because it influences the methodological quality and timeliness of reports, as well as promotes reliability of the HTA process to be used as basis for decision-making and jurisdiction.

Funding: Project funded by the German Chancellor Fellowship - Alexander von Humboldt Foundation.

Ethics statement: Not applicable.

Risk of osteonecrosis in bisphosphonate users submitted to dental procedures: systematic review and meta-analysis

Lúcio Henrique Ives Martins¹; Delaine Cristina Ferreira¹; Marcus Tolentino Silva¹; Rogério Heládio Lopes Motta¹; Reginaldo Tavares Franquez¹; Cristiane de Cássia Bergamaschi¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Although maxillary osteonecrosis be considered a rare condition, it can be refractory to available treatment therapies and clinically difficult to manage. Some dental procedures can represent a risk factor for triggering the condition. No recent data on the risk of this adverse effect after dental interventions in bisphosphonate users were found in the literature.

Objectives: This study determined the risk of osteonecrosis of the jaw in bisphosphonate users undergoing dental interventions.

Methods: This systematic review searched the following information sources: Medline (via Ovid), Embase (via Ovid), Web of Science, Scopus, Virtual Health Library, and the Thesis Repository of the Coordination of Superior Level Staff Improvement, with no restriction on language or publication date. Reviewers, in pairs and independently, selected the studies, extracted their data, and assessed the risk of bias.

Results: The risk of osteonecrosis in relation to bisphosphonate treatment duration was determined. The meta-analyses were pooled using the DerSimonian and Laird random effects model. A total of 27 studies (4,865 adult patients) were included, with samples comprising predominantly females and older adults. The most used bisphosphonates were intravenous zoledronate (n=17 studies) and oral alendronate (n=19) for treating cancers (n=11) or osteoporosis (n=16), respectively. Dental extraction was the most common procedure (n=21). Most of the studies were of low methodological quality (n=21). The risk of osteonecrosis was 0.8% (95%CI: 0.5-1.2%) proving higher for intravenous bisphosphonate [8.9% (95% CI: 6.5 to 11.5%)] than overall [0.0% (0.0-0.1%)]. There was no association between longer treatment duration and greater risk of osteonecrosis.

Conclusion: There was a higher risk of oral osteonecrosis in intravenous bisphosphonate users submitted to dental extraction. Further studies collecting more detailed information on the bisphosphonates used and with greater methodological rigor are suggested. The present findings can help inform prescribers, dental surgeons, and other health professionals on the risks of bisphosphonate use in patients undergoing dental interventions.

Funding: No funding to declare.

Ethics statement: Not applicable.

Safety and efficacy of treatment with zoledronate compared with pamidronate in children with osteogenesis imperfecta: rapid review

Cristiane Rocha de Oliveira¹; Grasiela Martins da Silva¹; Kelly Carneiro de Freitas Nakata²; Quenia Cristina Dias Morais¹, Verônica Clemente¹

¹ National Institute of Traumatology and Orthopedic, Rio de Janeiro, Rio de Janeiro, Brazil

² Mato Grosso State Health Secretariat, Cuiabá, Mato Grosso, Brazil

Context: Osteogenesis imperfecta (OI) is a genetic disease characterized by bone fragility, increased risk of fractures and skeletal deformities. Its prevalence is estimated at 1:20.000 born, being considered a rare disease. The aim of the treatment is to reduce fracture frequency. Bisphosphonates inhibit bone resorption. Intravenous pamidronate is the standard bisphosphonate treatment in children with OI. Its infusion must be within 4h for safety reasons. Zoledronate is third-generation bisphosphonate with higher potency and convenient administration because of shorter infusion time, but its efficacy and safety has not been established for OI.

Objectives: To review studies that compare zoledronate and pamidronate in children with OI.

Methods: It was used a rapid review methodology. Comparative studies were searched in PubMed, Cochrane Library and BVS. Of the 68 studies found, 2 were included by peers of reviewers. They were open, prospective, randomized and a retrospective observational study.

Results: Controlled study: 23 children, 11 used pamidronate 1mg/kg/day, for 2 days, every 3-4 months and 12 used zoledronate 0.025-0.05Mg/kg/day, for 2 days, in the same period. Respectively, after 1 year the groups increased lumbar spine (ls) bone mineral density (BMD) mean by 51.8% (P=0.053) and 67.6% (P=0.003); The fracture rate decreased (p=0.025 And p=0.048); there was no significant difference in CA metabolism parameters and side effects were similar in both groups. No serious clinical symptoms were reported. Observational study: 40 patients, 20 used pamidronate 1.5Mg/kg/day in 2 days, every 3 months and 20 used zoledronate 0.05Mg/kg/day, every 6 months. After 1 year, BMD LS Z scores increased significantly (p<0.001) +1.67(1.46–2.21) And +1.75(1.46–2.00), respectively. After 2 years, there was a similar incidence of long bone fracture rates between groups without new vertebral fractures. There were no serious adverse reactions.

Conclusion: Treatment with pamidronate and zoledronate showed similarity in safety and efficacy in children with OI in two comparative studies. Cost-utility studies are needed to consider preferences of patients and support ideal bisphosphonates protocol.

Funding: Project funded by Center for Studies, Research and Technological Development in Health Collective (CEPESC), grant n° SCON2020-00259.

Ethics statement: Not applicable.

Integration of dental attention to the multi-professional work process in the basic health unit of vila do engenho located in the municipality of Itacoatiara

Tawan Buás¹

¹ São Leopoldo Mandic College

Context: “Expedita Holanda” is a community health center, located in Itacoatiara, the state of Amazonas, Brazil. It local health center faced a problem; dental care was not included in the work processes by the health team in that location reality. This fact hindered the dental work process making the assistance provided to that population a merely curative activity in face of the complex health process and prevention process. In view of this reality, strategies were implemented for the integration of dentistry schedules, focusing on the development of work multidisciplinary care. This study was based on the recommendations of the national oral health policy on interdisciplinary and multiprofessionalism strategy and on the Brazil family health strategy which constitutes technological innovation on the way of management.

Objectives: To integrate dentistry schedule care into the routine of activities through a multi-professional work process in primary health care.

Methods: This was a case report study, with an application of observation techniques, daily monitoring, analysis of health indicators, and analyses of work processes, to identify practices that hinder integrated work. Meetings, team building, multi-professional strategies execution plan, awareness-raising activities through lectures, guidance on referrals, survey of demands for specialties, organization of dental records were used as strategies. It was supported by the involvement of the medical team, nursing and the relevance of the local Community Health Workers (CHWS) using technological and expedient resources by own financial resources, mainly used, for the displacement till the villages on the riverside areas and maintenance of the health team during these visits.

Results: Results were measured by the consolidated data got from the ministry of health, comparing the annual worker's production and health indicators of the previous years and the results obtained after the research. Significant changes were noticed in the whole work process, awareness-raising, and participation of professional health team and users about the integral and multi-professional assistance. It resulted in an increase of 58% in programed/ scheduled assistance in relation to the free demand. Improvement in the reception of pregnant women with 75% coverage by the first dental consultation. Strengthening prevention campaigns, unified and integral care (doctor / dentist / nurse). Guaranteed access to the riverside population was by targeted agenda, easy access, and continual assistance respecting their particularities and specific needs. These were the most significant and relevant points of management changes.

Conclusion: Public health requires from its workers, that it go beyond technical knowledge and skills, it's necessary to add capacities that involve collective responsibility for the results of the work, a humanized and welcoming view, in addition to knowing and experiencing the reality of the population of these locations, reaching then better results.

Funding: No funding to declare.

Ethics statement: Not applicable.

Avoidable public expenditures with selected antineoplastic drugs for breast cancer in Brazil

Ranailla Lima Bandeira dos Santos¹; Vera Lúcia Edais Pepe¹; Claudia Osorio de Castro¹

¹ Sergio Arouca National School of Public Health, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

Context: Breast cancer is the most incident type of among Brazilian women. Among the treatments for breast cancer, chemotherapy is used in 60% of patients. Timely treatment can cure, prevent metastasis and improve quality of life. Scaling up access for oncology drugs depends on effective public procurement.

Objectives: The study analyzes the Brazilian Federal procurement profile for selected antineoplastic drugs for breast cancer, and the behavior of acquisitions in relation to the quantities and prices practiced between 2013-2019, pointing out avoidable expenditures with these drugs.

Methods: Antineoplastic agents, category I of the ATC classification system, were selected according to their presence in breast carcinoma Brazilian guidelines either for adjuvant or neoadjuvant chemotherapy. Fixed combined treatment regimens were excluded. Regular purchases registered in the integrated system for administration of general services were analyzed, considering quantity, unit price, date, and purchasing agency. These agencies were linked to the three most important buyers in the public sphere: the ministry of health, the ministry of education and the ministry of defense. Any other public agencies were added in the “other” category. Prices were deflated to December 2019 by the Brazilian pricing index.

Results: Ten antineoplastics drugs were selected: anastrozole (1mg), docetaxel (40mg), exemestane (25mg), letrozole (2,5mg), paclitaxel (6mg), tamoxifen (20mg), trastuzumab (440mg), vinorelbine (10, 20 and 30mg). The “other” category presented greatest total expenditure. Trastuzumab and tamoxifen were responsible for the highest expenditure and volume, respectively. The ministry of education carried out the largest volume of purchases, except mentt for trastuzumab, which was most purchased by the “other” category. The ministry of defense presented the highest purchase prices, standing out for the highest overspending in the period. The total avoidable overspending, considering all categories and medicines was over BRL 99 million.

Conclusion: The analysis of government purchases can bring essential information for the understanding of availability and access to antineoplastic agents. The differences observed in the performance of different categories of buyers as to amounts purchased and prices practiced for antineoplastics drugs could be reduced by employing strategies to expand the centralization of purchases, resulting in expanded access to breast cancer medicines in the public sector.

Funding: Project supported by CNPq (Project number: 304975 / 2016-8) and; CAPES supported fellowship for RLBS.

Ethics statement: Not applicable.

Right to free health care and high-cost medications: brief notes

Anderson Miller Silva Varelo¹

¹ Paraíba State University, João Pessoa, Paraíba, Brazil

Context: In a still peripheral society like Brazil, the right to health is still exercised unequally. Thus, access to high-cost medications by the poorest people who depend on the Unified Health System (SUS) is not always equal and the citizen is forced to resort to the judiciary.

Objectives: This research aims to review the existing literature on these drugs and to understand the jurisprudential understanding of the subject.

Methods: The methodology employed was bibliographic, deductive, and qualitative research. The Scielo platform was used, with the descriptors: “high-cost drugs” and “judicialization of health care”, and 10 scientific articles were found.

Results: From the readings, it was observed that the reason why drugs are the object of litigation is due to a policy of medicine structuring that does not meet the demands arising from inequalities created by the asymmetrization of marginalized social classes and by existing budget limitations. Moreover, in view of the fact that certain medications are not regularly dispensed by SUS, the only way out is judicialization. The STF has established jurisprudential understanding that not just any medication can be demanded from the state for supply, and that certain requirements must be met, such as the existence of a registration request for the medication and the inability of the current treatments available to meet the demand, in addition to observing the parameters of reasonableness and the impacts on the budget, causing the granting of the request to be analyzed from a multifactorial standpoint.

Conclusion: Thus, we conclude that the effectiveness of the right to health in Brazil is conditioned to structuring parameters that are not always sustained in social equity and that the state cannot cover all the petitions of its citizens. However, such situation can be changed with better administrative organization.

Funding: No funding to declare.

Ethics statement: Not applicable.

Saber-SUS: Evidence based health and recommendations for SUS

Daniela Oliveira de Melo¹; Sandro Aparecido Tonin¹

¹ Federal University of São Paulo, São Paulo, Brazil

Context: The Saber-SUS is a project that intends to compile recommendations from guidelines of high methodological quality in order to develop a website and a distance learning course for sus professionals. Here we will present the results of the first step of the project, which included the identification, eligibility, and quality assessment of the guidelines.

Objectives: Select high-quality guidelines to serve as a basis for the development of support materials for sus professionals.

Methods: The most prevalent Chronic Diseases (CD) among older adults with multi- morbidity were listed: hypertension, chronic heart failure, stable angina, atrial fibrillation, coronary artery disease, hypercholesterolemia, diabetes mellitus, osteoarthritis, osteoporosis, chronic obstructive pulmonary disease, asthma, gastroesophageal reflux, dementia, depression, and benign prostatic hyperplasia. For each of the DCS, a process of identification and eligibility of the CPGS was carried out. The identification of CPGS was based on a systematic search of the literature (Medline (via PubMed), Embase and Cochrane library databases and in CPG repositories, updated on 09/10/2020. All cpgs developed from systematic review, published in English, Spanish or Portuguese, between 2015 and 2020 were included. Methodological quality assessment was performed using the agree-II tool. For each CD, the CPGS were selected considering a score equal to or greater than 60% in domains 3 and 6 of agree ii.

Results: After the identification and eligibility process, 356 CPGS were included for quality assessment. Of these, 85 (23%) were rated as high quality, although the mean scores for domains 3 and 6 were only $49\% \pm 22\%$ and $49\% \pm 24\%$, respectively. As in other studies, the applicability domain had the lowest score ($34\% \pm 21\%$). It was possible to identify at least 1 high quality CPG for each CD.

Conclusion: Even with wide variation in the mean score of the AGREE II domains, it was possible to list at least 1 high quality CPG to be adapted to the Brazilian context.

Funding: Project supported by FAPESP (PPSUS - 2019/03883-7).

Ethics statement: 4162164.

Analysis of financial viability in the participation of the municipality of Maringá in the Paraná health consortium

Larissa de Souza Zanolli¹; Marco Antônio Costa¹

¹ State University of Maringá, Maringá, Paraná, Brazil

Context: The present study analyzed the purchase of medicines from the Basic Component of Pharmaceutical Assistance (CBAF) through the Paraná health consortium and its bidding process in the city of Maringá.

Objectives: To assess whether there is a significant economic difference between the forms of face-to-face or electronic acquisition and whether there is economic viability in the municipality's participation in the consortium by quantifying the difference in total expenditure.

Methods: The documentary research considered the unit values of the drugs acquired through the bidding processes carried out by the municipality itself through face-to-face auctions and unit values practiced by the Paraná health consortium in 2019, by electronic auction, as well as the quantities of medicines purchased through the consortium, assigning to them the unit values of the municipal bidding and the quantity acquired by the municipality through its own bidding assigning the unit values practiced in the consortium.

Results: In terms of economy, we conclude that the two forms, face-to-face and electronic auctions, are similar since the average annual economy of the municipality compared to the consortium, reached a value equal to $9.95 \pm 1.52\%$, While the average of consortium was $9.81 \pm 0.73\%$. After analyzing the quantities acquired in medicines, it was found that in 2019 the total amount of R\$ 922,458.70 Was saved by the municipality, of which R\$ 468,275.61 was due to the acquisition through a consortium, already reducing the amount the annuity fee (R\$ 111,463.04) And R\$ 454,183.19 For the acquisition by municipal bidding.

Conclusion: Thus, we concluded that there is no statistically significant difference between face-to-face and electronic auctions and that there is financial viability in maintaining the two forms of acquisition, consortium and municipal bidding, considering the large savings generated for the municipality in 2019.

Funding: No funding to declare.

Ethics statement: Not applicable.

Profile of health judicialization involving drugs from specialized component of pharmaceutical assistance in the state of Pará, Brazil

Jocileide de Sousa Gomes¹; André Vilhena Da Silva², Amanda Thamires Cardoso Monteiro³

¹ Pará College, Belém, Pará, Brazil

² Federal University of Pará, Belém, Pará, Brazil

³ Amazônia College, Manaus, Amazônia, Brazil

Context: The financing of pharmaceutical assistance is the responsibility of the three management realms of Brazil's Unique Health System and the federal resources are transferred by pharmaceutical assistance financing blocks, which are made up of basic, strategic and specialized component. The specialized component is divided into three groups: Group 1, which is subdivided into 1A and 1B, group 2 and group 3. The drugs in group 1 are the responsibility of the federal government, those in group 2 must be supplied by the State and the Federal District, the last group, under the responsibility of Municipalities and the Federal District. In Brazil, the judicialization of health has compromised the stability and governance of the three realms of government over their financial resources for access to medicines.

Objectives: To identify the profile of the judicialization of health involving the specialized component of pharmaceutical Assistance in the state of Pará, in 2018.

Methods: Descriptive, quantitative research conducted from March to October 2019, involving spreadsheets of lawsuits organized by the State Department of Pharmaceutical Assistance and Central Supply and Logistics of the State Department of Health of Pará.

Results: The information from 30 lawsuits with complete registration of purchase of specialized component of pharmaceutical Assistance medicines was analyzed in 2018. The State Public Ministry was the main legal representation of them (70%), with most cases from the city of Belém of the Pará (33%) and with public prescriptions (70%). It was found that half of the cases demanded drugs from subgroup 1B, especially hormones and anti- psychotics. However, in subgroup 1A, although less in demand was the one that had the greatest financial impact, accounting for 73% of public spending for specialized component of pharmaceutical Assistance medicines requested by health judicialization (R\$ 94.754,37).

Conclusion: The health judicialization involving specialized component of pharmaceutical Assistance medicines, in the State of Pará, deserves to be highlighted as an important public management discussion agenda, especially in the planning of resources to be allocated to the acquisition of high-cost medicines, demanded by the court.

Funding: No funding to declare.

Ethics statement: Not applicable.

Profile of administrative demands for off label use of mofetilla mycophenolate, in the state of Pará

Ana Paula Almeida Gomes¹; Jocileide de Sousa Gomes²; Adriana Almeida Gomes¹

¹ Fibra College, Belém, Pará, Brazil

² Pará State Health Secretariat, Belém, Pará, Brazil

Context: Increasing access to medicines by administrative demands has established challenges for the management of the Unified Health System. In the state of Pará, administrative demands must comply with Normative Instruction No. 01/2017, which guides the use of evidence scientific justification for requesting the prescription drugs demanded by this route, among them, those for off label use. The prescription of off label medications corresponds to indications, uses and purposes different from those recommended in package inserts approved by the country's health authority.

Objectives: This study aimed to identify the profile of administrative demands involving the offlabel use of the medication Mycophenolate of Mofetila, sent to Pará State Health Secretariat (SESPA).

Methods: Descriptive research involving the request for mycophenolate mofetil for off label use, recorded in spreadsheets of administrative demands organized by the State Department of Pharmaceutical Assistance of Pará, in the year 2019 to 2020. For data organization and exploratory analysis, used the Microsoft Excel program.

Results: The drug Mycophenolate Mofetila is registered with the National Health Surveillance Agency as an Immunosuppressant, indicated for prophylactic treatment of organ rejection and in the treatment of refractory rejection of kidney, heart or liver transplantation. Between January 2019 and December 2020, SESPA received 508 administrative processes aimed at serving various medicines, 147 (29%) of which were prescribed for off-label use of Mycophenolate Mofetila. Of these, 84 (57%) were used to treat patients diagnosed with systemic lupus erythematosus, 53 (36%) for systemic lupus erythematosus with lupus nephritis and 10 (6.8%) for the treatment of systemic sclerosis. All the processes presented technical justification for such indications, however, the available evidence was 42% of studies sponsored by the pharmaceutical industry.

Conclusion: Although the National Commission for the Incorporation of Technologies in SUS has a favorable opinion for incorporating mycophenolate mofetil in the Clinical Protocol and Therapeutic Guidelines for Systemic Lupus Erythematosus, the National Health Surveillance Agency has not yet authorized the registration of such indication in the package insert referred drug, which makes its dispensation for that purpose unfeasible and contributes to the increase in the number of administrative demands against the State of Pará.

Funding: No funding to declare.

Ethics statement: Not applicable.

A comparative analysis of capillary blood glucose with deproteinized glucose oxidase in relevant clinical glucose concentrations in diabetic Wistar rats after beheading

Thanise Pitelli de Nigro¹; Susan Webber de Souza¹; Louryana Padilha Campos¹; Mateus Santana Lopes¹; Graciele More Manica¹; Geraldo Picheth¹; Fabiane Gomes de Moraes Rego¹

¹ Federal University of Paraná, Curitiba, Paraná, Brazil

Context: The measurement of capillary blood glucose is a practical and reliable form for the monitoring of induced diabetes in rats. The differences in blood rheology and other characteristics in human and laboratory rats could affect the glucose measurements since capillary tests were designed for human blood.

Objectives: We compared blood glucose capillary measurements with the Accu-chek active glucometer (Roche) with the deproteinized glucose oxidase method to monitoring glycemia in Wistar rats.

Methods: To determine glycosimeter analytical coefficient of variation (CVA), 10 ml of human whole blood (EDTA k2, vacutainer) obtained from a healthy individual was divided into four aliquots (1) "Very low" blood was incubated at 37°C for 6 hours; (2) "low" incubated at 37°C for 3 hours; (3) "normal" kept on ice; and (4) "high" glucose was added. Blood glucose was quantified in 20 independent determinations of each aliquot (15 µl of whole blood added to the test strip) in concentrations simulating hypo-, normo- and hyperglycemia. Additionally, the glucose levels were measured in 24 streptozotocin (STZ) diabetic Wistar rat tail blood with a glycosimeter and with deproteinized (trichloroacetic acid) glucose oxidase method (trinder god/pod labtest sa).

Results: The CVA in different glucose concentrations (mean glycemia) were 9.6% (0.90 Mmol/l); 4.6% (2.4 Mmol/l); 1.8% (4.6 Mmol/l) and 1.3% (16.4 Mmol/l). The comparison between the glycosimeter and deproteinized glucose oxidase showed a median (interquartile range) of 26.8 Mmol/l (13.3-36.6) And 20.1 Mmol/l (9.5-26.7), $P=0.020$ (Mann-Whitney U test). Therefore, a difference of about 25% was identified between the two methods. The two methods showed a high correlation (Pearson, $r=0.897$; <0.001). The capillary blood glucose CVA is higher in hypoglycemia (9.6%) and decreases in the presence of hyperglycemia (1.3%).

Conclusion: Both methods showed significant differences (~25% average) and high correlation ($r=0.89$), thus, we were able to identify the transition and monitoring of normal to diabetic rats.

Funding: No funding to declare.

Ethics statement: 23075.022975/2019-39.

Use of limonene as a therapeutic agent against *Escherichia coli* infections

Carlos Alberto Mendes da Silva Filho¹; Henrique John Pereira Neves²

¹ Tabosa de Almeida College, Caruaru, Pernambuco, Brazil

² Federal University of Pernambuco, Recife, Pernambuco, Brazil

Context: Limonene, 4-isoprenyl-1-methyl-cyclohexene, is a monocyclic monoterpene present in the composition of the essential oils of several aromatic plant species. It is found in citrus fruits and has a variety of industrial and microbiological applications. *Escherichia coli* is a gram-negative bacillus that inhabits the human intestine in large quantities. To diagnose it one must seed the collected material on blood agar and MacConkey agar to then perform biochemical tests such as production of indole through tryptophanase, decarboxylation of lysine and use of citrate as the only carbon source. Also, when it is mobile, it does not ferment sorbitol which is an important criterion to differentiate it from other strains.

Objectives: To perform the antibiogram with limonene on *Escherichia coli* strain and determine if this agent has therapeutic efficacy.

Methods: We prepared the limonene by taking 200g of orange peel and 500 ml of distilled water, processed in a blender and placed in the cleverer, this equipment separates the oil and water from the peel and the other components. Then we have put the disks impregnated with limonene on the plates and then in the oven, visualizing the result in the next day. We replicated the *Escherichia coli* strain supplied from the UFPE laboratory onto 50 blood agar plates.

Results: 90% of the strains showed sensitivity to limonene and 10% showed resistance to limonene.

Conclusion: In view of the results obtained, limonene can be used in the therapeutics for *Escherichia coli*.

Funding: No funding to declare.

Ethics statement: Not applicable.

Effects of *Lentinula edodes* cereal bar consumption on antioxidant profile: a double blind randomized clinical trial

Sara Rosicler Vieira Spim¹; Ana Maria Holtz Pistila¹; Thaisa Borim Pickler¹, Marcus Tolentino Silva¹; Denise Grotto¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Shiitake (*Lentinula edodes*) is considered a nutritionally complete food, high amount of protein, vitamins, minerals, and bioactive compounds, such as glucans and phenols. These compounds have antioxidant and anti-inflammatory properties.

Objectives: The objective of this study was to evaluate antioxidant status in borderline high cholesterol individuals consuming cereal bars with shiitake. This is a phase II, double-blind randomized trial.

Methods: Individuals aged from 20 to 65 years old, of both genders, who had one of these biochemical markers at the borderline high level (total cholesterol 180 to 239 mg/dl; low density lipoproteins (LDL) 130 to 159 mg/dl; triglyceride (Tg) of 150 to 200 mg/dl) were recruited. 68 Individuals were randomized into 2 groups: (I) shiitake food bar (n=32); (II) food bar without shiitake (n=36). Blood collection was performed at 0.33 and 66 days after bar consumption. Reduced glutathione, catalase and thiobarbituric acid reactive substances were evaluated using spectrophotometric methods. Data were expressed as mean $\pm$ standard deviation. Differences between shiitake and placebo groups were analyzed by Student's t-test, differences among the three times were performed using ANOVA test, and data were normalized to logarithmic scale. $P < 0.05$ Was accepted as statistically significant.

Results: 70% of individuals completed participation. Glutathione levels increased significantly after ingesting bars with shiitake compared to the placebo group ($p=0.0058$). Substance reactive to thiobarbituric acid showed a significant difference ($p=0.0384$) In the intervention group in relation to placebo, showing a reduction in lipid peroxidation. There was no difference in catalase levels over time.

Conclusion: Thus, cereal bars with shiitake are a functional food alternative, well-accepted by population and they demonstrated beneficial action and improvement in the antioxidant status of individuals.

Funding: Project funded by Yuri Mushrooms, JP Pereira Ltda and University of Sorocaba.

Ethics statement: 2824297.

Indicators for safe medication use: systematic review

Gledsen Garcia de Paiva Melo¹; Luciana Contijo¹; Flavio Marques Lopes¹

¹ Federal University of Goiás, Goiânia, Goiás, Brazil

Context: The Patient Global Safety challenge proposed to reduce 50% of serious and preventable damage related to medications in the next five years. In Brazil, this challenge is extremely difficult since there is not much knowledge about medication errors in hospitals. These factors make us question: how to improve something that we have not yet measured? Hence the importance of searching for safe medication indicators (SMI).

Objectives: Identify and classify (according to Donabedian) the indicators for safe use of medicines applied in hospital environments.

Methods: A methodological survey was carried out in the Medline, PubMed, Lilacs, Scielo, and gray literature databases, through Scholar Google. There was no language restriction or time frame. All studies that dissipated at least one medication-related quality indicator were included, whether numerators and denominators were defined, or could be interpreted, and whether consensus methods were developed. Studies that did not dissipate SMI in hospitals were excluded, as well as letters, editorials, news, comments from professionals, case studies, and articles without an abstract. Two independent evaluators selected the studies.

Results: The search identified 6,621 studies, of which 18 met the inclusion criteria. Most studies were conducted in the Netherlands and published in English. 391 SMI were found, of which the majority were from the process. Most of their data location was identified in the patients' medical records. Out of the 391 SMI, 269 were medication indicators in general and 122 of antimicrobial medication. Most studies developed indicators using the literature review method associated with the Delphi technique with modifications.

Conclusion: 391 SMI were identified and classified according to Donabedian's proposal. Such indicators are essential for monitoring and analyzing medication errors.

Funding: No funding to declare.

Ethics statement: Not applicable.

Comparison of fungiculture and viniculture residues as biosorbents for lead contamination

Nathália Roberta Cardoso Mendes Castanho¹; Giovanna Caroline Floriano Dias¹; Renan Angrizani de Oliveira¹; Rodrigo Mendes Pereira²; Raquel Andrade Leite Vieira¹; Bruno Lemos Batista²; Angela Faustino Jozala¹; Denise Grotto¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² Federal University of ABC, Santo André, São Paulo, Brazil

Context: Pollution is a subject of extreme recurrence, because in addition to environmental damage, environmental contamination can be causally linked to public health problems. Heavy metals contaminants, such as Pb (lead), can damage the blood system, in addition to being neurotoxic, especially for children. According to the United Nations Children's Fund (UNICEF), about 1 in 3 children has low blood Pb levels but can silently wreak havoc on children's development and health. Searching for alternative materials for remediation of pollutants, fungiculture and viniculture residues were proposed to be studied mainly by the wide range of production in Brazil, in addition to easy post-use treatment, not increasing the cost of research.

Objectives: In view of these points, the work sought to evaluate and compare the biosorption capacity of fungiculture and viniculture residues, simulating Pb-contaminated water contamination in laboratory concentrations.

Methods: This is an experimental study. The fungiculture and viniculture residues were dried in an oven and ground in the knife and hammer mill, going through the granulometry process through sieves of different sizes. Samples of each residue were added separately in flasks with Pb solution. Flasks were kept under agitation, and each of them were removed at different times to analyze the sorption capacity of biosorbents. To analyze the remaining Pb in water, Inductively Coupled Plasma Mass Spectrometry (ICP-MS) apparatus was used. Results were presented in percentage and they were analyzed by the ANOVA test.

Results: The sorption potential for all biomaterials was of great relevance; the grape bagasse had 93.8% Pb removal in 60 minutes, while the shiitake substrate had 72% (30 minutes), the shiitake stalks 82% (30 minutes), and the champignon stalks 83% in 120 minutes.

Conclusion: The percentages of Pb removal were excellent, demonstrating there are possibilities to make use of fungiculture and viniculture residues, adding value to a waste that would be discarded.

Funding: Project funded by the São Paulo Research Foundation (FAPESP), grant n° 16/22873-4.

Ethics statement: Not applicable.

Doxycycline safety through hematological, hepatic, and renal profile in chronic pre-clinical exposure

Alex Sandro Brizolla Atoji Junior¹, Natália Tribuiani Silva¹, Yoko Oshima-Franco¹, Denise Grotto¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Doxycycline, a well-known antimicrobial agent, has been widely studied in several parameters for off-label use, mainly for its properties that have emerged in studies of neuroprotection, oxidative stress reduction and for Parkinson's disease treatment. Evaluation of its therapeutic safety in chronic exposure is necessary when the focus is Parkinson's disease, which requires continuous treatment, and the usual doxycycline treatment is used for short periods.

Objectives: To evaluate the safety of doxycycline on the hematological, hepatic, and renal parameters in chronic exposure.

Methods: *In vivo* study. Male Wistar rats were divided into two groups: control and experimental (n=8 animals/group). The dose of doxycycline was 10mg/kg/day, for 90 days. After euthanasia, blood was collected for hematological assessment, creatinine (Crea), urea (Ure), alanine aminotransferase (Alt), aspartate aminotransferase (Ast) dosages, and liver was histologically analyzed. Student's t test was applied to verify statistical differences between groups. Values of $p < 0.05$ were considered significant.

Results: There was a decrease in leukocytes and monocytes, and an increase in corpuscular hemoglobin in the doxycycline group, which may be linked to intrinsic mechanisms of the drug. Regarding hepatic and renal markers, no differences were observed comparing doxycycline to control in Crea (0.47 ± 0.08 And 0.52 ± 0.06 Mg/dl, respectively, $p=0.141$), Ure (51.7 ± 9.3 And 55.2 ± 0.06 Mg/dl, respectively; $p=0.246$), Ast (115.1 ± 18.9 And 125.3 ± 21.0 U/l, respectively; $p=0.342$), Alt (115.1 ± 18.9 And 73.0 ± 10.1 U/l, respectively; $p=0.320$). Liver also did not show any significant structural changes in cell morphology or tissue architecture.

Conclusion: Doxycycline, in the parameters of hepatic, hematological and renal profile, is safe for chronic administration.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: 114/2017.

Biogenic synthesis and characterization of silver nanoparticles based on *Beauveria bassiana*

Bianca Migliorini Ramos¹; Renata de Lima¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Studies indicate that biogenic metallic nanoparticles hold promise because of their high activity and low toxicity.

Objectives: The study aimed to synthesize silver nanoparticles using the fungus *Beuveria bassiana*, perform its physical-chemical characterization and its toxicity on cell lines through MTT and Trypan Blue.

Methods: The nanoparticles were synthesized using the *Beauveria bassiana* filtrate as a reducing and stabilizing agent, and silver nitrate (AgNO₃). For physical-chemical characterization, nanoparticle tracking and dynamic light scattering techniques were used. Toxicity tests were performed with HaCaT, 3T3 and V79 strains.

Results: The synthesis of the nanoparticles was successful and the physical-chemical characterization showed a hydrodynamic diameter of 135.4 ± 4.0 Mn, polydispersion index of 0.366 ± 0.036 , Zeta potential of -14.2 ± 1.3 , size of 114.4 ± 7.5 , and concentration of $2.87E + 011$. By the MTT assay, it was observed that only the HaCaT strain showed IC₅₀, while by the Trypan Blue assay, a reduction in the viability of all cell lines was observed in the three concentrations tested.

Conclusion: Thus, the results obtained indicate that the use of nanotechnology can be a promising path.

Funding: Project funded by the National Council for Scientific and Technological Development (CNPq).

Ethics statement: Not applicable.

Analysis in laboratory-scale of cadmium removal in aqueous solution utilizing bacterial nanocellulose

Gustavo Alexandre dos Santos¹; Angelica Martiniano Dias¹; Nathalia Roberta Cardoso Mendes Castanho¹; Victoria Soares Soeiro¹; Rodrigo Mendes Pereira¹; Bruno Lemos Batista¹; Denise Grotto¹; Angela Faustino Jozala¹.

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Bacterial nanocellulose has several applications for allowing strategies to use its micro and nano elastic fibers for a dynamic design, such as filters.

Objectives: Therefore, this study used bacterial nanocellulose to remove cadmium in aqueous media.

Methods: For that, samples of bacterial cellulose were produced by culturing *Gluconacetobacter xylinus* ATCC 53582 using standard medium Hestrin & Schramm (HS). After production, the membranes were kept whole and crushed. Subsequently, they were kept wet and were also lyophilized, to perform the adsorption kinetics test of the metallic solution of cadmium at 100ppm. The kinetics test was carried out in 50 ml flasks containing 20 ml of the metallic solution and 0.1 g of cut or not nanocellulose, lyophilized or not, was added. The flasks were kept shaking at 100 rpm, at 25°C, for up to 24 hours. At predetermined times, the vials were removed, and the solution was separated from the nanocellulose. The volume of the final solution, the final mass of the nanocellulose, and the efficiency of cadmium removal was analyzed.

Results: As a result, it was possible to verify that the entire lyophilized nanocellulose retains 2 times greater volume of solution than the cut one. However, wet shredding showed greater metal removal efficiency.

Conclusion: The use of nanocellulose can be attractive because it is biodegradable, has good molecular interactions, returns filtered water, and does not leave toxic residue and removes the polluting metal.

Funding: Project funded by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Antimicrobial evaluation of a bioproduct composed of bacterial nanocellulose, fibroin and nisin

Victoria Soares Soeiro¹, Carolina Fernanda Oliveira Santos Tallarico¹, Gabriel Pereira Machado¹, Erick Unterkircher¹, Nathália Roberta Cardoso Mendes Castanho¹, Denise Grotto¹, Norberto Aranha¹, Angela Faustino Jozala¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: There is a trend in research aimed at the production and application of bioproducts as dressings, as they promote patient comfort, above all, by pain relief.

Objectives: The aim of this study was to evaluate an antimicrobial activity of nisin when loaded on bacterial nanocellulose and fibroin.

Methods: This research is an *in vitro* pre-clinical experimental study. For this, nanocelluloses were incorporated at 0.01 g/ml of nisin by immersion at 100 rpm, for 4 hours. After this period, the material was immersed in a fibroin solution and maintained at 100 rpm, for 24 hours. After, membranes were dried at room temperature and evaluated for antimicrobial activity. The agar diffusion methodology was carried out using the microorganisms *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Streptococcus mutans* and *Streptococcus oralis*. In addition, scanning electron microscopy (SEM), tomography and mucoadhesiveness of the membranes were performed.

Results: As a result, no antimicrobial activity against gram-negative bacteria was observed, as expected. Nisin has a greater activity against gram-positive bacteria. For this reason, it inhibited the growth of *S. aureus*. The mucoadhesive strength of the lyophilized was performed and, analyzing the results, it was observed that the values of the force exerted to remove it from the mucin disc tended to decrease considerably with the presence of fibroin. The SEM with eds shows the presence of fibroin and nisin inside the nanocellulose.

Conclusion: In the near future, it will be possible to adjust nisin for greater antimicrobial activity.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

Phytol-based nanoemulsion: a promising anti-cancer action

Victoria Louise Pinto Freire¹; Mariana Farias Alves da Silva¹; Debora Caroline de Azevedo Pereira¹; Antonio Arnobio da Silva Junior¹

¹ Federal University of Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil

Context: Cancer is a multifactorial pathology characterized by uncontrolled cell proliferation and one of the main causes of death worldwide. In the past year, approximately 10 million deaths were reported. The toxicity and development of multidrug resistance to conventional chemotherapy support the need to find effective treatments. Phytol is a bioactive component of chlorophyll belonging to the diterpenes class. A wide range of biological activities are found for phytol, including cytotoxic effects on cancer cells. In this perspective, nanotechnology has become associated with medicine to improve the pharmacokinetic and pharmacodynamic properties through bioactive encapsulation.

Objectives: The aim of the study is to perform a pre-clinical experimental study *in vitro* to develop nanoemulsions containing phytol as a promising alternative for enhancing antiproliferative activity in tumor cells.

Methods: The nanoemulsion was prepared by the low energy phase inversion method. The type of co-solvent was varied to find the monodispersed nanometric formulation. The composition and method parameters were evaluated measurements of particle diameter, polydispersion index (PDI) and zeta potential (ZP) by Dynamic Light Scattering. All results are expressed as mean and standard deviation.

Results: The physico-chemical properties were analyzed, obtaining average particle size 212.77 ± 2.48 nm (PDI 0.38 ± 0.01) and negatively charged ZP -29.93 ± 0.15 mV for the formulation using glycerin as a co-solvent. For those of propylene, the particle size was 233.17 ± 11.47 nm (PDI 0.55 ± 0.10) and ZP -20.07 ± 0.76 mV. Phytol was incorporated in both formulations at a concentration of 5 mg/ml.

Conclusion: Experimental data showed that the surfactant variation was successfully optimized, and the system prepared with glycerin had a smaller diameter and polydispersity, proving to be a promising nanoemulsion for *in vivo* and *in vitro* tests for anti-cancer activity.

Funding: Project funded by the National Council for Scientific and Technological Development (CNPq), grant n° 479195/2018.

Ethics statement: Not applicable.

Clinical trials with hormonally active herbal substances on the reproductive health of men and women: systematic literature review

Rilva Lopes de Sousa Muñoz¹; Eduardo Sérgio Soares Sousa¹

¹ Federal University of Paraíba, João Pessoa, Paraíba, Brazil

Context: Hormone-active substances extracted from plants can be beneficial to human health. There are many plant compounds with this potential. The present study was delimited on the hormonal effects related to the hypothalamus-pituitary-gonadal axis of men and women.

Objectives: The aim of this study was to summarize and evaluate the evidence from clinical trials on the effects of natural bioactive substances extracted from plants with a presumed action on the hypothalamic-pituitary-gonadal axis of men and women using herbal preparations in scientific publications of the last 20 years.

Methods: This systematic review was conducted in accordance with the Reporting Guidelines for Systematic Reviews and Meta-Analyses (PRISMA). The PubMed, Scopus and Web of Science databases were searched using the following terms: “phytotherapy” or “medicinal plant” and “reproductive health” or “gonadal hormones” and “therapeutic action”. Abstracts of clinical studies involving humans were included, evaluating the effects of active plant ingredients as the main result. The intervention had to be an herbal medicine in a prospective clinical trial (not necessarily controlled) in the form of extract(s) from a single plant or preparation(s) containing several plants. The papers were screened by titles and abstracts by two reviewers, with independent data extraction.

Results: 92 clinical trials were identified that met the eligibility criteria, 59 involving women and 33 involving men. Among the phytotherapies used, there were 40 natural substances and seven generically called Chinese herbs in combination. The most frequent substances were soy isoflavones, fenugreek (*Trigonella foenumgraecum*) and puncture vine (*Tribulus terrestris*), both for postmenopausal women and men, indicated mainly for vasomotor symptoms, polycystic ovary syndrome and erectile dysfunction. Studies have shown that plant compounds are promising for the treatment of disorders of the reproductive sphere. The soy isoflavone extracts were supported by the best evidence.

Conclusion: Clinical trials are available on a wide range of herbal preparations prescribed for conditions in the sphere of reproductive health. Medicinal plants with effects on disorders of the hypothalamic-pituitary-gonadal axis of women and men at various stages of life, such as climacteric syndrome, polycystic ovaries and erectile dysfunction with therapeutic effects are valuable for modern systems for the discovery of natural medicines. More work needs to be done to better understand the most relevant approaches.

Funding: No funding to declare.

Ethics statement: Not applicable.

Validation of the x-ray fluorescence technique for the use in pharmaceutical sciences

Thais Hora Paulino Estanagel¹; José Martins de Oliveira Junior¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Elementary impurities in medicines can arise from several sources and therefore, there is a need to investigate the chemical composition of these drugs, since the presence in excessive concentrations of some elements, as well as the deficiency of others, can lead to a series of metabolic disorders. For this investigation to be carried out with quality it is essential to validate the technique used. The energy dispersion x-ray fluorescence technique (ED-XRF) can contribute to this end, because it is an elementary analysis technique capable of identifying and quantifying a range of chemical elements in different concentrations.

Objectives: Validate the ED-XRF technique for use in pharmaceutical sciences, whose focus is on elementary identification and quantification.

Methods: The samples are transformed into tablets using a press and then the fluorescence analyzes are performed with equipment sold by the company panalytical, model Epsilon-1, which uses an excitation x-ray source with variable voltage from 20kV to 50kV and silver target; silicon drift detector and different filter sets. The estimated reading time for each sample is variable to provide data with adequate statistics for the research objectives. The qualitative and quantitative results are obtained by equipment's software, which performs the fundamental analysis parameters.

Results: Our results are in accordance with the parameters: (a) selectivity for matrix in the presence of impurities; (b) linearity in the concentrations of 70ppm, 35ppm, 17.5ppm, 8ppm, 4ppm, 2ppm and 1ppm; (c) accuracy in the working range of 80% to 120% recovery and calculation of standard deviation; (d) precision below 4.5% for the relative standard deviation; (e) limit of detection <0,5 ppm; (f) limit of quantification <1,5 ppm and (g) robustness.

Conclusion: The increase in demands for reliable and certified analytical results imposes the need to implement a quality system and the technique proposed in this work proved to be effective and has been successfully validated for use in the various segments of the pharmaceutical sciences.

Funding: Project funded by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Anti-inflammatory *in vivo* evaluation by the air pouch model of the crude hydroalcoholic extract of *Myrcia pubipetala* (EBH-MP)

Pâmela Pacassa Borges¹; Aline Lana Schneider¹, Eduardo José Wagner¹, Ester Teresinha Schmitt¹, Michele Alberton Debiasi¹, Isabel Daufenback Machado¹

¹ Regional University of Blumenau, Blumenau, Santa Catarina, Brazil

Context: *Myrcia pubipetala* is an endemic plant present in Brazil, part of the *Myrtaceae* family, with important specimens in popular use, such as *jaboticaba*, guava and eucalyptus that have reported anti-inflammatory activity.

Objectives: To evaluate the innate immune response involved in the *in vivo* anti-inflammatory activity of the extract of *Myrcia pubipetala* miq.

Methods: Pre-clinical experimental study. We cordially received the EBH-MP from Dr. Michele Alberton Debiasi. The air pouch model consists in a space in the subcutaneous tissue on the dorsum of the mice induced by air injection for 7 days. On the seventh day the animals were treated in doses of 3, 30 and 300 mg/kg (v.O.) Of EBH-MP or indomethacin or vehicle, one hour later 1ml of 1% carrageenan was administered in the air pouch to induce inflammation acute, after 4h we collected the exudate and tissue. The leukocyte count was performed, and nitric oxide was measured on the exudate samples. Part of the tissue covering the pouch was collected. The statistical analysis used was ANOVA by graphpadprism, admitting $p < 0.05$.

Results: All doses tested reduced the total leukocyte count, and the volume of exudate was reduced at the doses 30 and 300 mg/kg. EBH-MP was effective as well as indomethacin with no statistical difference. The nitric oxide production on the exudate was reduced by EBH-MP at 30 and 300mg/kg. In the histological analysis was possible to visualize the reduction of leukocyte migration and exudate production.

Conclusion: The results obtained suggest that the EBH-MP has an anti-inflammatory effect *in vivo*, and more tests will be performed to elucidate the biological activity and the action mechanism.

Funding: Project funded by the Santa Catarina Research Foundation (FAPESC) and by the Regional University of Blumenau (FURB).

Ethics statement: 009/2020.

Use of citrus fruit-derived essential oil as a bactericidal agent on *Escherichia coli* strains

Gleidisson Soares da Silva¹; Kassy Jhons de Souza Silva¹; Henrique John Pereira Neves²

¹ Tabosa de Almeida College, Caruaru, Pernambuco, Brazil

² Federal University of Pernambuco, Recife, Pernambuco, Brazil

Context: *Escherichia coli* is found in coprocultures and is easily cultivated. To diagnose it, the collected material must be seeded on blood agar and MacConkey agar, and then biochemical tests must be performed, such as: production of indole by tryptone, decarboxylation of lysine, use of citrate as the only carbon source, and when it is mobile it does not ferment sorbitol, which is an important criterion to differentiate it from other strains. Essential oils are natural products formed by aromatic plants, characterized as complex mixtures, usually colorless or slightly yellowish, with a strong odor, soluble in organic solvents of low polarity, not very stable, especially in the presence of air, light, heat, moisture, and metals. Essential oils are found and formed in secretory structures located in different organs of plants, such as: leaves, flowers, stems, rhizomes, roots, fruits, or seeds.

Objectives: To determine the bactericidal action of essential oil against *Escherichia coli* strains.

Methods: *Escherichia coli* strains, supplied by the laboratory of the Federal University of Pernambuco, were placed in petri plates with the culture medium blood agar. We re-potted them to 50 petri plates with blood agar and after that we placed the disks impregnated with the essential oil on the plates, in five different positions with a distance of approximately 5 cm from the end of the plate. They were then placed in the oven at 35.5°C for 24 hours. The next day we read the results using the halo size conference, showing sensitivity or resistance to the essential oil used.

Results: After the antibiogram it was observed that in 50 plates, 10 showed resistant colonies and in the remaining 40 plates there was total sensitivity.

Conclusion: Therefore, the use of essential oil showed a bactericidal efficacy in the strains used in this study.

Funding: No funding to declare.

Ethics statement: Not applicable.

***In vitro* antidiabetic activity of medicinal plants used by communities in the city of Vitória de Santo Antão - PE, Brazil**

Patrícia Maria da Silva Neri Cruz¹; Jorge Veras Filho¹; Jenifer Rodrigues de Oliveira¹; Isa Renata de Souza Santos¹; Thiago Antônio de Souza Araújo²; Elba Lúcia Cavalcanti de Amorim¹

¹ Federal University of Pernambuco, Recife, Pernambuco, Brazil

² Maurício De Nassau College, Recife, Pernambuco, Brazil

Context: A lot of plants with popular indications to treat diabetes or its symptoms have been the subject of scientific studies to prove its therapeutic effects.

Objectives: Therefore, the purpose of this study was to evaluate inhibitory activity against the enzyme α -amylase from plant extracts of the most suitable species to treat diabetes, and its comorbidities.

Methods: A semi-structured questionnaire was made to obtain data about plant species, then plants with higher citation rates were subjected to extraction by maceration, and it was used 92,8° INPM ethanol as a solvent, with solvent renewal every 72 hours, until the exhaustion. The extracts were filtered and obtained by evaporation under reduced pressure, at a temperature of $40 \pm 5^\circ\text{C}$, until total dryness. Then, the enzyme inhibition assay of α -amylase was performed by using the colorimetric reagent 3,5 – dinitrosalicylic acid (DNS) of yellow color, and then converting it to 3-amino-5-nitrosalicylic of red-brown color in the presence of sugars reducers. The reaction is quantified spectrophotometrically at a wavelength of 540nm, and it can evaluate an enzymatic activity.

Results: The ethnospices that have the lowest number of citations were *Syzygium jambolanum* (java plum), *Cissus sicyoides* (possum grape), *Mentha x vilosa* (small mint), *Cymbopogon citratus* (lemongrass), *Morus nigra* (mulberry), *Lippia alba* (citronella), *Bauhinia cf. monandra* (cow's leg) and *Plenctranthus amboinicus* (large mint). The extract of *Bauhinia cf. monandra* presented the best IC₅₀ value (0.778 ± 0.014 mg/ml), and it does not present a change statistically different of the extracts of *Syzygium jambolanum* and *Lippia alba*, against a positive pattern of acarbose with an IC₅₀ of 0.967 mg/ml. The extracts of large mint, small mint and lemongrass did not dissipate enzyme inhibition and behaved as amylolytic.

Conclusion: The extracts that have the best inhibition values obtained an IC₅₀ lower than the positive standard that was used. It shows the potential of these species, and indicate a possible mechanism of action of these in the treatment of diabetes, corroborating to the popular use as well.

Funding: Project funded by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: 2006671.

Understanding the use of nanoparticles in drug delivery systems and its importance in the treatment and monitoring of breast cancer

Eliel Luiz de Oliveira Araujo¹; Adrielle Antonia da Silva¹; Tais Genuino da Silva¹; Priscila Magna do Nascimento Silva¹; Eva Caroline dos Santos¹; Pablo Ramon Gualberto Cardoso¹; Bárbara Virgínia Mendonça da Silva Correia¹; Auvani Antunes da Silva Júnior¹

² Maurício De Nassau College, Caruaru, Pernambuco, Brazil

Context: Having value and potential application in treatments, diagnostic and monitoring processes, nanoparticles allow drug delivery systems and act as biosensors due to their unique chemical and physical properties. Breast cancer treatment involves the action of chemotherapy, which administered directly can generate adverse effects. Nanoparticles comprise an important part of the development and improvement of technologies that assist the treatment, prognostic and supervision of patients affected by breast cancer.

Objectives: The aim of the literature review was to elucidate the use of nanoparticles in drug delivery systems and their ability to share the therapeutic process and monitor breast cancer.

Methods: Descriptors and articles used were in English, dated between 2016 and 2020 indexed in ScienceDirect and PubMed databases. Inclusion criteria was review articles that cite use of nanoparticles in combination with drugs for treatment of breast cancer. Exclusion criteria was articles that dealt with non-drug therapies and were directed at other types of cancer.

Results: Of 764 articles found, 10 were selected. In each article, aspects were analyzed that allowed to understand how the use of nanoparticles in combination with drugs is positive for the treatment and monitoring of breast cancer. The outcome was analyzed in relation to standard drug treatment involving chemotherapy. Therapeutic agents such as trastuzumab, neratinib and talazoparib, in association with nanoparticles are examples of those used in the treatment. The prospects are that nanocomponents allow greater accuracy when interacting with neoplastic cells, favor regulated distribution of medications, target cancer stem cells, and allow resistance to factors such as pH changes and tissue hypoxia, therefore, reducing the impact on healthy cells and minimizing adverse effects. Properties of organic and inorganic nanoparticles such as polarization and associated surface molecules also allow cancer monitoring.

Conclusion: The nanoparticles associated with drugs allowed good sensitivity and specificity in the treatment of breast cancer, the regulated and direct distribution of the drug reduced adverse effects, properties associated with the particles allowed its identification and monitoring of the disease.

Funding: No funding to declare.

Ethics statement: Not applicable.

Preliminary study of medicines used against androgenetic alopecia by mutagenicity parameter (Ames Test)

Ederson Constantino¹, Caroline Malavasi Barros¹, Anna Paula Farias de França¹, Jaqueline de Cássia Proença Assunção¹, Estrela D'Aurea Machado², Flávia Aparecida Resende Nogueira³, Yoko Oshima-Franco¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² São Paulo State University, Botucatu, São Paulo, Brazil

³ University of Araraquara, Araraquara, São Paulo, Brazil

Context: Androgenetic alopecia (AGA) is a condition that interferes with the quality of life of the person affected by the disease, besides the self-esteem and social life. The search for the best treatment to eradicate or minimize hair loss depends on the safety of its use. To measure safety to a compound or mixture, its toxicity, including genotoxicity, should be known and the Salmonella/microsome assay is recommended for this purpose.

Objectives: The objective of this study was to evaluate the safety of the combination finasteride with a blend containing growth factors, through the Salmonella/microsome parameter (Ames test).

Methods: Finasteride concentrations (50 mg/ml), blend of growth factors (1.2 mg/0.1 ml) and finasteride + blend (50 mg/ml each) were preliminary tested at the maximum limit recommended by the regulatory agencies (5 mg/plate) with the strain *S. typhimurium* TA100 for the toxicity test. Subsequently, the assay was performed in strains of TA97a, TA98, TA100 and TA102, in the absence (-S9) of exogenous metabolic activation. The results were statistically analyzed with controls (positive of each lineage and negative) and the mutagenicity indexes (IM 2.0) calculated. All trials were performed in triplicate.

Results: The maximum values of 5 mg/finasteride plate, blend and combination did not present toxicity to the TA100 strain. In the Ames Test, without metabolic activation, finasteride, blend, and association were not mutagenic in the tested strains.

Conclusion: The commercial blend studied did not show to be a direct mutagen, because it did not present mutagenicity in the absence of metabolic activation.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not Applicable.

Probiotic formulations for the treatment of intestinal constipation: development and evaluation

Fabiana Cristina Rosa Mitelmão¹; Cristiane de Cassia Bergamaschi¹; Marli Gerenutti²; Karin Häckel³; Marcus Tolentino Silva¹; Victor Manuel Cardoso Figueiredo Balcão¹; Marta Maria Duarte Carvalho Vila¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² Pontifical Catholic University of São Paulo, São Paulo, São Paulo, Brazil

³ Clinic of Gastroenterology Karin Häckel, Sorocaba, São Paulo

Context: Functional intestinal constipation is a clinical manifestation that affects about 20% of the world population. Various treatments, such as dietary interventions and the use of laxatives, are used. However, conventional treatment does not always provide significant improvement. In this regard, functional supplementation with probiotic microorganisms can be useful in reducing the disorder.

Objectives: The aim of this study was to assess the stability of probiotic mixtures for the use as food supplement.

Methods: Two formulations with probiotics were developed, one with *Lactobacillus acidophilus*, *Bifidobacterium bifidum* and *Lactobacillus rhamnosus* (3 billion Colony-Forming Units-CFU); and another with *Lactobacillus acidophilus*, *Bifidobacterium bifidum*, *Lactobacillus rhamnosus*, *Lactobacillus paracasei*, *Bifidobacterium longum*, *Bifidobacterium lactis*, *Lactobacillus defens*, *Bifidobacterium animalis* (8 billion CFU). Supplements were evaluated for their stability, resistance to gastric and intestinal juice at 0, 3, 6, 9, 12 and 24 months and evaluation of microbiological quality.

Results: The samples showed maintenance of physical-chemical parameters, with a characteristic aroma of vanilla, white color, pH between 5.5 to 7.0 and humidity 4.9%. The microbiological analysis indicated adequate sanitary conditions with the presence of fecal coliforms 5 CFU/g and absence of *Bacillus cereus*, coagulase-positive *Staphylococci*, and *Salmonella*. The average population of viable *Lactobacillus* and *Bifidobacteria* cells proved to be as expected, with a viability reduction of less than 20% in the tests performed at 0, 3 and 6 months. Regarding exposure to gastric and enteric fluids, the formulations also showed values of less than 20% in reducing bacterial viability.

Conclusion: Thus, from the results presented, the products developed showed stability and can be considered promising for the use as food supplement aimed at treating functional constipation.

Funding: No funding to declare.

Ethics statement: Not applicable.

Preliminary toxicity assessment of silver nanoparticles (AgNPs) synthesized with curcumin (Ames Test)

Jaqueline de Cássia Proença Assunção¹; Ederson Constantino¹; Caroline Malavasi Barros¹; Anna Paula Farias de França¹; Carolina Alves dos Santos¹; Flávia Aparecida Resende Nogueira²; Marco Vinicius Chaud¹; Yoko Oshima-Franco¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² University of Araraquara, Araraquara, São Paulo, Brazil

Context: Silver nanoparticles (AgNPs) are of great interest mainly for their antibacterial properties. Here we produced AgNPs by the curcumin-reduction on silver ions.

Objectives: The aim of this study was to evaluate the mutagenicity of AgNPs (0.081 mg/ ml, 130 nm) synthesized with curcumin from *Curcuma longa* (Zingiberaceae) through the Salmonella/microsome (Ames test) preliminary tests.

Methods: The design of this study was *in vitro*. These preliminary toxicity tests for AgNPs in TA98 and TA100 strains were performed without metabolic activation. The tests were performed in triplicate, in a controlled condition (positive, negative, and spontaneous controls), and the AgNPs were tested at concentrations of 0.0081; 0.0060; 0.0040; 0.0020; 0.0010 mg/plate. Mutagenicity indexes $MI \geq 2.0$, in any strain, indicate mutagenicity. The results were statistically analyzed and values with $p < 0.05$ were considered significant.

Results: In the absence of metabolic activation, AgNPs were not mutagenic in any of the concentrations evaluated, as the mutagenicity indexes were less than 2.0.

Conclusion: Under the parameters of the preliminary test Salmonella/microsome, the AgNPs did not present mutagenicity in the absence of metabolic activation, therefore, they are not classified as direct mutagens.

Funding: Project funded by the São Paulo Research Foundation (FAPESP), grant n° 2015/01420-9, and by Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

Development of curcumin gel and choline-geranate for psoriasis treatment

Rodrigo Boscariol¹; Marta Maria Duarte Carvalho Vila¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Psoriasis is a chronic inflammatory skin disease with a strong genetic predisposition, auto-immune pathogenic traits, with recurrent course and variable worldwide prevalence. Curcumin is the main bioactive extracted from the rhizome of *Curcuma longa* and has antioxidant, anti-inflammatory and antibacterial properties. Due to these properties, curcumin has been used to treat various skin conditions, such as psoriasis. However, permeation by the stratum corneum is hampered by its low solubility and absorption. To improve skin permeation, substances that facilitate topical permeation, such as ionic liquids, have been studied.

Objectives: To develop a curcumin gel and ionic liquid (choline-geranate) for the transmission of curcumin to treat psoriasis.

Methods: This is a pre-clinical experimental study *in vitro*. The synthesis of the ionic liquid was performed in the reaction between choline bicarbonate and geranic acid [1: 2]. Different proportions of choline-geranate (0%; 0.5%; 1%; 1.5%; 2% and 2.5% w/w) were evaluated to define the lowest concentration of ionic liquid capable of permeating the asset. *In vitro* skin permeation studies were performed using Franz diffusion cells and porcine ear skin as a membrane. The measurement of permeate curcumin was performed by spectrophotometry at $\lambda = 421\text{nm}$. Choline-geranate was synthesized in 45% yield.

Results: The product showed fluidity and transparency at room temperature. The curcumin dosage method showed a correction coefficient of 0.9983, with an equation ($A = 6.0571C + 0.0003$) showing to be linear and suitable for the intended purpose.

Conclusion: After the experiments, a greater permeation of curcumin for the sample with 2% of ionic liquid can be observed. This proportion being used for the next experiments.

Funding: Project funded by the Coordination of Superior Level Staff Improvement (CAPES).

Ethics statement: Not applicable.

Preliminary assay: safety assessment of doxycycline by mutagenicity assay in *Salmonella typhimurium* microsomes

Natália Tribuiani¹, Jocimar de Souza¹, Yoko Oshima-Franco¹.

¹ University of Sorocaba, Sorocaba, SP, Brazil.

Context: Doxycycline has shown promise for the treatment of chronic neurodegenerative diseases including Parkinson's Disease. The repurposing by regulatory agencies of certain medicines depends on the knowledge about the effects of chronic oral administration as suggested for Parkinson's disease treatment. The Salmonella/microsome mutagenicity assay can provide information on the mutagenic potential of doxycycline and at the same time establish safety limits for its use. However, it is only possible to perform the test after knowing the toxicity on *S. typhimurium* strains, in a preliminary toxicity test.

Objectives: To evaluate the safety of doxycycline by mutagenicity parameters.

Methods: This was an experimental pre-clinical *in vitro* study. Preliminary concentrations of doxycycline (ml) were tested to select those that were not toxic. The mutagenic activity was tested by the Salmonella/microsome assay, using strains of *S. typhimurium* TA 97a, TA98, TA100 and TA102, with and without metabolism by the pre-incubation method. The results were analyzed with the Salanal statistical software package. The data were evaluated by analysis of variance (ANOVA), followed by linear regression.

Results: From the chosen concentration of 3.125 µg/ml, the Ames test was performed with and without metabolic activation. Doxycycline did not show mutagenic changes in strains of *S. typhimurium* with and without metabolic activation.

Conclusion: Doxycycline proved to be safe under the mutagenicity parameter, which can stimulate long-term use in neurodegenerative diseases.

Funding: Project funded by the Institutional Scholarship Program of Scientific Initiation (PROBIC) of the University of Sorocaba.

Ethics statement: Not applicable.

The benefits of using the ethanolic extract of *Catharanthus roseus* L.

Isabele Caroline Pereira da Silva¹; Henrique Jhon Pereira Neves²; José João dos Santo Neto¹

¹ Tabosa De Almeida College, Caruaru, Pernambuco, Brazil

² Federal University of Pernambuco, Recife, Pernambuco, Brazil

Context: The *Catharanthus roseus* plant, popularly known as rose periwinkle, originally from Madagascar is abundantly found in the Brazilian territory due to the ease of its cultivation in tropical climate. There is an interest in deepening the studies with this plant, as research points to the presence of tannins and alkaloids in its flowers, substances with interesting pharmacological characteristics, which can have healing, antimicrobial, antifungal and other actions.

Objectives: To evaluate the pharmacological activities of the ethanolic extract of the *Catharanthus roseus* flowers.

Methods: This was an experimental, pre-clinical and in vitro study. Obtaining the plant extract by the dry material extraction method using ethanol as a water-miscible organic solvent, where the extract was dried and fractionated between aqueous phase and water immiscible phase, the alkaloids that were retained in the aqueous phase were alkalized and subsequently extracted with ethanol. Then, the extract was purified by column chromatography. Once the extract was obtained, it was used to perform the antibiogram with strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*, and the antifungi-gram with *Candida sp.*

Results: The flower extract presented activity for both strains.

Conclusion: It is necessary further studies to confirm such activity.

Funding: No funding to declare.

Ethics statement: Not applicable.

Biogenic synthesis of silver nanoparticles based on *Penicillium sp* to combating resistant pathogens

Vitória Aparecida Nobrega Antunes¹, Tais Germano-Costa¹, Natalia Bilesky-José¹, Mariana Guilger-Casagrande^{1,2}, Leonardo Fernandes Fraceto², Renata Lima¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

² São Paulo State University, Sorocaba, São Paulo, Brazil

Context: Studies involving silver metallic nanoparticles have aroused the interest of researchers in recent years, this is mainly due to their bactericidal potential. Among these nanoparticles, the biogenic silver nanoparticles (AgNP) seem to be the most highly rated for application in the most different areas, due to the fact that they present excellent bactericidal action combined with low toxicity.

Objectives: The aim of this study was to synthesize biogenic AgNPs using filtrate of the fungus *Penicillium citrinum*, to evaluate its activity and toxicity.

Methods: Pre-clinical *in vitro* experimental study. The AgNPs were synthesized using the filtrate of *Penicillium citrinum* and silver nitrate (AgNO₃) as a precursor. After synthesis, the characterization was performed to obtain size, polydispersity index (PDI), zeta potential and concentration. 3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium Bromide (MTT) analysis and comet assay were used to measure the viability and genotoxicity, respectively. The bactericidal activity was also evaluated by means of the agar diffusion test, as well as by the minimum inhibitory concentration (MIC) and determination of the minimum bactericidal concentration (CBM) in pathogenic microorganisms and resistant bacteria.

Results: The diameter of the AgNPs was 184.4 ± 4.1 nm, polydispersion 0.3 ± 0.06 , zeta potential -20.7 ± 1.4 mV, average size of 107.6 ± 5.1 nm and concentration of 2.55×10^{10} NPs/ml. IC₅₀ values per MTT were 5.75×10^9 NPs/ml (3T3) and 5.58×10^9 NPs/ml (V79). The results showed that AgNPs have bactericidal activity, controlling the growth of microorganisms in pure and diluted concentrations. The results of the comet assay showed that there are no genotoxic effects of AgNPs.

Conclusion: The results showed that AgNPs can be an option for the control of pathogens since they achieved the expected results without presenting toxicity.

Funding: Project funded by the National Council for Scientific and Technological Development (CNPq), and the University of Sorocaba.

Ethics statement: Not applicable.

***In vitro* study of direct inhibition of cholinesterase enzyme by the organophosphate chlorpyrifos**

Jocimar de Souza¹; Fernando de Sá Del Fiol¹; Yoko Oshima-Franco¹

¹ University of Sorocaba, Sorocaba, São Paulo, Brazil

Context: Organophosphate compounds appeared between the years 1945 and 1970 due to the toxicity of organochlorines, which were used until then, one of the main products being Chlorpyrifos registered in 1965. Organophosphates are responsible for more than 36% of the total global market for chemical pesticides due to its high effectiveness for the control of sucking and chewing insects. The insecticidal activity of the organophosphate occurs, in most cases, from its neurotoxic action, through the “irreversible” inhibition of the enzyme acetylcholinesterase.

Objectives: To investigate, *in vitro*, the direct inhibitory action of the organophosphate pesticide chlorpyrifos from two different sources: extracted from a commercial product (Pyrinex® 480 EC) and commercial standard Sigma®.

Methods: The measurement of plasma cholinesterase, performed in triplicate, adding to the samples the control, containing the enzyme cholinesterase in concentrations: 4.4 mg/ml; 5.4 mg/ml and 6.4 mg/ml. Neostigmine [1 mg/ml] was used as a positive control. DMSO, solvent for chlorpyrifos, was also dosed as a vehicle control. The technical procedure was performed according to the manufacturer’s instructions (Bioclin Química Básica Ltda. Santa Branca, Belo Horizonte, MG, Brazil). Cholinesterase activity was determined spectrophotometrically at a wavelength of 405 nm.

Results: Both tests were shown to be cholinesterase enzyme inhibitors, indicating that the non-metabolized form of the organophosphate under study has its neurotoxic action. The extracted compound showed a high inhibitory power, not indicating a significant difference when compared to the positive control (Neostigmine). Contrary to what occurs with the commercial standard, that is, it has not shown high efficacy in direct *in vitro* inhibition. The results were expressed as the mean $\pm$ SEM and statistically analyzed by one-way ANOVA with $p < 0.05$ indicating significance.

Conclusion: Therefore, the literature presents the metabolite of chlorpyrifos, chlorpyrifos-oxon, with neurotoxic action. However within this study, it was shown that the pesticide can be a direct inhibitor of the cholinesterase enzyme.

Funding: No funding to declare.

Ethics statement: Not applicable.

Regional analysis of the implementation of pharmaceutical care in the health care network and its perception by pharmaceutical professionals

Julio Eduardo Pereira de Souza¹; Marina Tiemi Shio²; Álvaro Avezum Junior²

¹ Family Health Association, São Paulo, São Paulo, Brazil

² Santo Amaro University, São Paulo, São Paulo, Brazil

Context: The research deals with the role of the pharmacist in Pharmaceutical Care (PC) in Primary and Specialized Health Care, after the regulation of Ordinance nº 1,918/2016 (Municipal Health Secretariat of São Paulo) that expanded adherence to drug treatment in the Unified Health System (SUS).

Objectives: The objectives were to assess the perception of pharmacists and analyze PC after the implementation of the Ordinance.

Methods: An online questionnaire was used before and after standardizing the Ordinance and data was collected in pharmaceutical consultations. Inclusion criteria were adult and elderly patients with chronic non-communicable diseases who had difficulty adhering to drug treatment. Children and adolescents were excluded. The outcome measures of the initial hypothesis, qualifying the PC, were defined before data collection. Statistical tests used were: C de Cochran, Fisher's exact test and Chi-square test.

Results: For the pharmaceutical perception, 32 pharmacists answered the questionnaire. No differences were observed before or after the standardization of the PC Ordinance, interventions, perception of the ability to intervene and improve results. They often participated in team meetings and pharmaceutical consultations. They mentioned that they felt more inserted in the team (81.25%), they needed to change their professional posture (87.50%) and make an academic update (96.88%). The consulted patients were 40 and 89 years old. They had comorbidities: Diabetes mellitus Type 1 and 2, Systemic Arterial Hypertension and Dyslipidemia (25%). The number of pharmaceutical consultations ranged from one (50%) to twelve consultations (2.78%). The problems related to pharmacotherapy were drug interaction (35.63%) and incorrect administration (19.59%). The classifications of drug interactions moderate severity (87.07%) to high (12.93%). The interventions used were specific guidelines on diseases/drug treatment (45.83%), Medication Organizing Box (69.89%), among others (2.78%).

Conclusion: The implementation of the CP Ordinance increased the pharmaceutical bond between patients and staff, reducing problems.

Funding: Not supported by funding.

Ethics statement: Not applicable.