Short Communication

Open Access

Unknown Mutation Detection via Restriction Hybridization Method Instead of Using Next Generation Sequencing Method

Umair Masood

Department of Biology, Pakistan

ABSTRACT

In biology mutation is a change in the nucleotide sequences of the DNA of an organism. Mainly there are three types of mutation: point mutation, deletion and insertions. Once the mutation has been defined allele specific oligonucleotide hybridization, amplification, heteroduplex formation method referred to as a diagnostic method. Some advanced technique like CRISPR cas9 system is used for selected mutagenesis. Using restriction method system we can detect a mutation. Let's say you have a DNA sample with fluorescent labeled from patient and you want to make sure that gene you are interested in is healthy. We can design different short fragment sequences to scan through DNA or find specific gene or mutation. The sequences scan the DNA; if the sequences do not find targeted gene, it does not bind to it. This means that no fluorescence color appears under UV-light. Each different short fragment sequence is labeled with different colors. If the different short fragments sequence does not bind to the DNA or specific gene or area, this means that there will be no color appear under UV light. This part or gene will be separated from the DNA by using restriction enzyme to do a Sanger sequencing gel electrophoresis. Result of the Sanger sequencing will provide the information about sequence of unknown part or gene of the DNA. This method is easier and cost economic method instead of Next generation sequencing method [1-7].

*Corresponding author

Umair Masood, Department of Biology, Pakistan. E-mail: umairawan0505@gmail.com

Received: October 26, 2021; **Accepted:** November 06, 2021; **Published:** November 15, 2021

Result and Observation

In the given diagram E,G,F does not show any color under UV light. This means that E,G,F part will be separated from the DNA in order to perform a Sanger gel electrophoresis. E,G,F have unknown variation or unknown SNP which cannot be available in any Database or data sections fig 1.1.

References

1. Mulepati S, Héroux A, Bailey S (2014) Crystal structure of a Caspr RNA-guided surveillance complex bound to a ssDNA target. *Science*. 345: 1479-1484.
2. Barrangou R (2015) The roles of CRISPR-Cas systems in adaptive immunity and beyond. *Current Opinion in Immunology*. 32: 36-41.
3. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis 285: 125-144.
4. Tytgat Olivier (2021) STRide probes: Single-labeled short tandem repeat identification probes. *Biosensors and Bioelectronics*. 180: 113135.
5. Olsen GJ, Lane DJ, Giovannoni SJ, Pace NR, Stahl DA (1986) Microbial ecology and evolution: a ribosomal RNA approach. *Annu. Rev. Microbiol.* 40: 337-365.
6. Chen J, Miller BF, Furano AV (2014) Repair of naturally occurring mismatches can induce mutations in flanking DNA. *eLife*. 3: e02001.

7. Amann R, Ludwig W (2000) Ribosomal RNA-targeted nucleic acid probes for studies in microbial ecology. FEMS Microbiology Reviews. 24: 555-565.

Copyright: ©2021 Umair Masood. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.