Case Report

Open Access

Coronally Repositioned Semilunar Flap for Root Coverage: A Case Report

Sunny Mavi

MDS, Periodontics and Implantology, Sudha Rustagi College of Dental Sciences & Research, Haryana, India

ABSTRACT

Gingival recession is one of the most commonly encountered periodontal problem. It is one of the most difficult to treat periodontal issue since it requires great amount of clinical skills and patient co-operation. There are plethora of techniques and their modifications available with different levels of predictability for root coverage. One of the highlighted technique amongst many is the coronally repositioned semilunar flap. It has various advantages like tension free flap during healing (without sutures), better predictability and minimal clinical skills required. This case report describes in detail a case treated with semilunar flap and discusses further about it.

*Corresponding author

Dr. Sunny Mavi, Ho. No. 1283, Sector 21D, Faridabad, Haryana, India. E-mail: snnymavi@gmail.com

Received: June 11, 2021; **Accepted:** June 17, 2021; **Published:** July 03, 2021

Keywords: Gingival recession, Semilunar Flap, Coronally Advanced Flap, Keratinized tissue, Full Thickness Flap

Introduction

One of the commonest and difficult to treat periodontal condition is the gingival recession. Gingival Recession is defined as location of the gingival margin apical to the cemento-enamel junction [1]. The main etiologic factors of gingival recession are:

Periodontal disease: - it is the interaction between bacterial infection and the host immune response causes bone resorption, and down-growth of the epithelium followed by gingival recession [2].

Mechanical forces: Faulty tooth brushing—A common cause is improper brushing technique with aggressive tooth brushing leading to gradual abrasion the gingival tissue. In these cases, gingiva appears to be free of inflammation, the apical shift of the marginal gingiva denudes root surfaces [3].

Iatrogenic factors: Orthodontic movement—the movement of the teeth may result in the loss of the alveolar plate, followed by gingival recession [4].

Restorative dentistry: Crown preparations extending subgingivally is one of the reason for gingival recession [5]. Poorly-designed dentures can lead to the gingival recession [6].

Anatomical factors: Many anatomical factors can be correlated to gingival recessions. These are alveolar bone dehiscence, high frenum pull, gingival biotype and tooth position [7-9].

Overview of Various Techniques for Root Coverage

Several techniques have been proposed for the treatment of isolated gingival recession defects. The main treatment goals are

to achieve root coverage and increasing keratinized tissue width. Some of the pedicle soft tissue grafts like lateral pedicle graft, double papilla flap and oblique rotational flap, advanced flaps like coronally advanced flap [13] and semilunar flap have shown successful root coverage [10-14].

Coronally advanced flap combined with additive treatments like non resorbable barrier, resorbable barrier, enamel matrix derivative, platelet rich plasma, acellular dermal matrix, growth factors, tissue engineering and low intensity laser therapy have also shown predictable root coverage [15-21]. In this case report we intend to discuss simple & predictable technique of root coverage i.e. Coronally Repositioned Semilunar flap.

Technique

To begin with, the case selection for any root coverage procedure is of utmost importance. The prognosis of any root coverage procedure in Miller's class 1 & 2 gingival recession is much better as compared to the Millers class 3 & 4 gingival recession cases. After appropriate case selection, the initial preparation which consists of plaque control instruction, scaling and root planing is done 2 weeks prior to surgery. On the day of surgery, adequate anesthesia is obtained at the surgical site. The exposed root surfaces need to be planed thoroughly. A no. 15 blade is used for the incision. A semilunar incision is made following the scalloped shape of the free gingival margin. It extends till the core of the papilla but does not disturb papillary tip. The incision can be extended into the alveolar mucosa if the width of Keratinized tissue is insufficient to cover the recession. The incision should curve apically mid facially so that the apical part of the flap rests on bone after it is displaced. A crevicular incision is given. Following this, it is continued as a full thickness flap till the apical semilunar incision. This completely frees the tissue from the underlying bone. The crescent shaped facial tissue is then displaced coronally

towards the cemento enamel junction so as to cover the denuded root surface. The tissue is compressed in place with a moist gauze for 5- 10 minutes [22].

Case Report

A forty-five-year-old male patient reported to the Department of Periodontics with the chief complaint of gums moving upwards in the left upper side region of the mouth since past 3 months. On examination, patient was found to be suffering from Chronic Periodontitis with generalized periodontal pockets and Miller's Class 1 gingival recession in relation to maxillary right and left canine (13 and 23). For treating the gingival recession, a semilunar flap procedure was decided. Adequate anesthesia was attained at the surgical site. A semilunar incision was made following the contour of free gingival margin extending till the core of the interdental papilla. It was followed by placing a crevicular incision. The tissue between the two incisions was reflected (full thickness) with the periosteal elevator. A through and through passage could be observed. The flap was coronally advanced with the digital pressure and moist gauze to completely cover the denuded root surface. The flap was compressed at the new site for 5 minutes using a moist gauze. A tinfoil was placed and a periodontal dressing was given over it for 1 week. The patient was given the necessary post operative instructions and medications. The patient was evaluated 1 week, 1 month and 6 months post operatively.

Figure 1: Pre-operative view

Figure 2: Semilunar incision given in gingival recession in relation to 23

Figure 3: Crevicular incision given & probe under the full thickness flap displaying the through and through passage

Figure 4: Semilunar flap repositioned coronally

Figure 5: Periodontal dressing placed

Figure 6: One week post-operative view

Figure 7: One month post-operative view

Figure 8: Six months post-operative view

Discussion

The coronally repositioned periodontal flap has been mentioned by numerous writers in the literature. Kalmi was one of the first to describe a type of coronal repositioned flap which was performed after gingivoplasty of the attached gingiva [23]. Similarly,

Nordenram and Harvey employed surgical techniques that covered the denuded roots by coronally repositioning the mucoperiosteal flaps [24-26]. Many others like Sumner and Ward introduced modifications of the coronal repositioned flap. Bernimoulin et al. described a two-step coronally repositioned periodontal flap wherein a free gingival graft placed as well [27-29]. These numerous publications had affirmed the clinical success of the coronally repositioned flap as a root coverage procedure.

In the late 1980s, Tarnow introduced the semilunar coronally repositioned flap [22]. This technique is advantageous over other coronally positioned flaps as no sutures are required as well as there is no tension on the flap whatsoever. The vestibule depth is not affected and most importantly it does not interfere with the existing papillary unit.

The semilunar flap is indicated in cases of Miller's Class I gingival recession with minimal labial sulcus depth. In case when adequate zone of keratinized tissue is not present, a free autogenous soft tissue graft can be attempted. This procedure can be used where esthetics is a major concern. It can also be used to treat gingival recession around previous full coverage restorations in the anteriors [22].

Many case reports such as those by Jahangirnezhad, Ramya & co workers have been previously published [30,31]. These also showed a similar efficacy in the root coverage and highlighted the superiority of the procedure as compared to other root coverage procedures.

Conclusion

This case report represented a particularly valuable technique for root coverage i.e the coronally repositioned semilunar flap. The procedure overshadows the other root coverage techniques in the sense of clinical ease, postoperative patient comfort and predictable outcomes.

References

- Newman MG, Takei HH, Klokkevold PR, Carranza FA (2006) Clinical Periodontology. 10th ed. Missouri: Elsevier Saunders; 2006.p.370.
- Hangorsky U, Bissada NB (1980) Clinical assessment of free gingival graft effectiveness on the maintenance of periodontal health. J Periodontol 51: 274-275.
- Hall WB (1977) Present status of soft tissue grafting. J Periodontol 48: 587-597.
- Dorfman HS (1978) Mucogingival changes resulting from mandibular incisor tooth movement. Am J Orthod 74: 286-297.
- Donaldson D (1973) Gingival recession associated with temporary crowns. J Periodontol 44: 691-696.
- Gartrell JR, Mathews DP (1976) Gingival recession. Dent Clin North Am 20: 199-224.
- Lang NP, LoeH (1972) The relationship between the width of keratinized gingiva and gingival health. J Periodontol 43: 623-627.
- Buckley LA (1981) The relationship between malocclusion, gingival inflammation, plaque and calculus. J Periodontol 52: 535-540.
- Wennstrom J, Lindhe J, Nyman S (1982) The role of keratinized gingiva in plaque associated gingivitis in dogs. J Clin Periodontol 9: 75-85.
- Grupe HE, Warren RF Jr. (1956) Repair of gingival defects by a sliding flap operation. J Periodontol 27: 92.
- Cohen ES, Ross S (1968) The double papilla flap in periodontal therapy. J Periodontol 39: 65-70.
- Pennel BM, Higginson JD, Towner TD (1956) Oblique rotated flap. J Periodontol 36: 305-309.
- Bernimoulin JP, Luscher B, Muhlemann HR (1975) coronally repositioned periodontal flap. Clinical evaluation after one year. J Clin Periodontol 2: 1-13.
- Tarnow DP (1986) Semilunar coronally repositioned flap. J Clin Periodontol 13: 182-185.
- Pini Prato G, Tinti C, Vincenzi G, Magnani C, Cortellini P, Clauser C (1992) Guided tissue regeneration versus mucogingival surgery in the treatment of human buccal gingival recession. J Periodontol 63: 919-928.
- Roccuzzo M, Lungo M, Corrente G, Gandolfo S (1996) Comparative study of a bioresorbable and a non-resorbable membrane in the treatment of human buccal gingival recessions. J Periodontol 67: 7-14.
- Castellanos AT, de la Rosa MR, de la Garza M, Caffesse RG (2006) Enamel matrix derivative and coronal flaps to cover marginal tissue recessions. J Periodontol 77: 7-14.
- Jankovic S, Klokkevold P, Dimitrijevic B, Kenney E, Camargo P (2012) Int J Periodontics Restorative Dent 32: 41-50
- Henderson RD, Greenwell H, Drisko C, Regennitter FJ, Lamb JW, et al. (2001) Predictable multiple site root coverage using an acellular dermal matrix allograft 72: 571-582.
- Kawase T, Okuda K, Wolf LF, Yoshie H (2003) Platelet-rich plasma-derived fibrin clot formation stimulates collagen synthesis in periodontal ligament and osteoblastic cells in vitro. J Periodontol 74: 858-864.
- Murata M, Okuda K, Momose M, Kubo K, Kuroyanagi Y, et al. (2008) Root coverage with cultured gingival dermal substitute composed of gingival fibroblasts and matrix: a case series 28: 461-467.
- Tarnow DP (1986) Semihinarcoronally repositioned flap. J. Clin. Periodontol 13: 182-185.
- Kalmi J, Moscor M, Goronov Z (1949) The solution of the aesthetic problem in the treatment of periodontal disease of anterior teeth: Gingivoplastic operation. Parodontologie 3: 53-55.
- Nordenram, A, Landt, H (1969) Evaluation of a surgical technique in the periodontal treatment of maxillary anterior teeth. Acta Odontologica Scandinavica 27: 283-286.
- Harvey P. M (1963) Management of advance periodontitis. Part 1. Preliminary report of a method of surgical reconstruction. New Zealand Dental Journal 61: 180-184.
- Harvey P, M (1970) surgical reconstruction of the gingiva. Part II, procedure. New Zealand Dental Journal 66: 42-46.
- Sumner C F (1969) Surgical repair of recession on the maxillary cuspid. Incisally repositioning the gingival tissues. J Periodontol 40: 119-122.
- Ward H, Simering M (1973) Manual of clinical periodontics. St. Louis: C. V. Mosby Co. 81-82.
- Bernimoulin JP, Luscher B, Muhlemann HR (1975) coronally repositioned periodontal flap. Clinical indication after one year. J Clin Periodontol 2: 1-3.
- Jahangirnezhad M (2006) Semilunar coronally repositioned flap for the treatment of gingival recession with and without tissue adhesives: A Pilot study. J Dent Tehran 30: 36-40.
- Ramya V, Bhuwaneshwarri, Padmanabhan P, Manisundar N (2014) Semilunar Coronally Positioned Flap for the Treatment with Bilateral Miller's Class I Buccal Gingival Recessions- A Case Report. Biosci Biotech Res Asia 11: 353-355.

Copyright: ©2021 Sunny Mavi. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.