Case Report

Open Access

A Family of Matched Parent-Child HLA Haplotypes: A Case Study from Bahrain

Mariam Fida*, Ferdinand N. Webnda and Ryan Ed Bangate

Department of Molecular Medicine, Al-Jawhara Center for Molecular Medicine and Inherited Disorders, College of Medicine and Medical Sciences, Arabian Gulf University, P.O. Box 26671, Manama, Kingdom of Bahrain

*Corresponding author

Dr. Mariam Fida, MD, PhD, P.O. Box 26671, Manama, Bahrain

Received: September 13, 2021; **Accepted:** September 20, 2021; **Published:** September 28, 2021

Keywords: HLA, Bone Marrow, Transplant, Donor, Genetics, Bahrain

Introduction

Bone marrow transplant (BMT) is a medical procedure performed to replace the soft, fatty bone marrow that has been damaged or destroyed by disease, infection, or chemotherapy. Most bone marrow transplants are usually performed on patients with certain blood cancers (leukemias) and other blood disorders. Cases of fully matched offspring(s) with parents are very rare and would occur when both parents share at least one haplotype which is commonly seen in several generations of consanguineous marriages [1].

The possibilities of having a perfect match is raised in family members especially siblings due to the shared alleles from the same parents. Cases of fully matched offspring(s) with parents are very rare and would occur when both parents share at least one haplotype which is commonly seen in several generations of consanguineous marriages [2].

Nuclear family members provide the best possibility of a perfectly matched donor. The probability of finding a matched parent-child is higher in consanguineous marriages [3].

The high probability is due to the cultural trend that is common in the Middle East, West Asia and North Africa where large families have high population growth, density and increased rates of consanguineous marriages. A study also in Saudi Arabia revealed that 60% of all patients had the possibility of finding a sibling being a match [4].

To check for a match, a small amount of blood is drawn and the five main HLA alleles are analyzed and compared with the other family members of with the recipient in need of the transplant.

Case Presentation

We present a case of a unique family from Bahrain with four children; two sons and two daughters. All six family members presented to the clinic to test for HLA tissue typing. It is a blood test that identifies antigens on the surface of cells and tissues to check for a match for an organ transplant. Results showed the two sons were a perfect match with the mother and a daughter who was a perfect match with the father.

The four siblings and their parents were tested for HLA as they had a family member who needed a bone marrow transplant and they were interested in being a donor. Initially, when asked, the mother stated that she and her husband are not related despite having the same last name.

The results showed that of the four children, both sons were 100% match with the mother, while one of the two daughters was a 100% match with the father. When the results of the parents-children were confirmed to be a 100% HLA match, a detailed family pedigree of the extended family was drawn which clearly showed a common ancestor.

Materials and Methods

Six DNA samples were extracted from peripheral blood using the Qiagen QIAamp DNA Mini Kit. According to manufacturer's instructions (Qiagen, Hilden, Germany). The Qubit® 3.0 Fluorometer was then used to quantify each of the extracted DNA sample.

A separate PCR protocol was set up for each locus to be amplified, and for each individual sample to be tested. Each run included appropriate positive control/s of known genotype, and at least one negative control for each locus being amplified. Successful amplifications were confirmed on a 1% agarose gel electrophoresis using 2µl of PCR product with 5µl loading buffer.

Sequencing of the amplified product was done using the Conexio Genomics SBT Resolver kits for High resolution Molecular HLA typing with group specific and locus specific primers.

The HLA Sequencing Based Typing (SBT) procedure used was originally developed by D. Sayer in 2001 and developed into a single tube assay in 2004 [5]. The procedure involves the initial amplification of the target sequence followed by enzymatic treatment to remove unincorporated primers and dNTPs. The amplicon is then used as a template for direct automated fluorescent DNA sequencing using customized sequencing primers and the Big Dye® Terminator sequencing kit V3.1 (Applied Biosystems, Foster City, California, USA).

The sequencing products were purified using the ethanol precipitation method and denatured with Hi-Di™ Formamide available from

Applied Biosystems™ by Life Technologies™. The purified fragments were sequenced using the ABI 3130XL Genetic Analyzer (Applied Biosystems). The data generated was then analyzed and the results were interpreted according to the manufacturer's guidelines using the CareDx Pty Ltd's ASSIGN™ SBT software from CareDx Pty Ltd3-5. (CareDx Pty Ltd's OLERUP SBT™ HLA SBT kits) used for typing of HLA Class I and Class II genes from genomic DNA. The sequencing products were purified using the ethanol precipitation method and denatured with Hi-Di™ Formamide available from Applied Biosystems™ by Life Technologies™. The purified fragments were sequenced using the ABI 3130XL Genetic Analyzer (Applied Biosystems).

The data generated was then analyzed and the results were interpreted according to the manufacturer's guidelines using the CareDx Pty Ltd's ASSIGN™ SBT software from CareDx Pty Ltd3-5. (CareDx Pty Ltd's OLERUP SBT™ HLA SBT kits) used for typing of HLA Class I and Class II genes from genomic DNA.

Results

Our experience of HLA typing show the known 50% match between parent and child. However, in this case, and due to the consanguineous marriages seen within generations in the family (figure 1), we observed a shared haplotype in both parents (figure 2) leading to children being a 100% match to either one of the parents.

Figure 1: Extended family pedigree showing consanguineous marriages seen within generations in the family

Figure 2: Son 1 – 50% matched with the father and 100% matched with the mother
 Daughter 1 – 50% matched with the mother and 100% matched with the father
 Son 2 – 50% matched with the father and 100% matched with the mother
 Daughter 2 – 50% matched with the mother and 50% matched with the father

There were very strikingly matched allele patterns seen in the children tested for HLA-A, HLA-B, HLA-C, HLA-DRB1, and HLA-DQB1 (Figure 2).

Due to the high number of shared alleles between the parents, both sons were a 100% match with the mother, while one of the two daughters (daughter 1) was a 100% match with the father. Daughter 2 inherited the two HLA haplotypes that were not shared by the parents. Son 1 was a 100% match with son 2. He was also a 50% match with both his sisters (daughter 1 & 2). Both daughters were a 50% match with each other.

Discussion

Following the normal inheritance pattern, a biologic child usually is a 50 % match with his or her parent because each child receives half of the HLA genes from each parent. Reported findings in a Saudi population study revealed that, patients in need of HCT have a greater chance of finding an HLA-matching sibling than is reported in most Western countries [4]. There was a 43% chance of finding a matching sibling in patients aged 0 to 5 years, this percentage increased to 68% in patients aged 20 and above. The change in pattern observed in contrast to that in the western world in the Saudi population is mainly because of the larger number of siblings in most Saudi families [4]. However, among communities where the parents are related, the probability of matched parent-child as well as matched siblings are higher [2].

It has been reported in the literature from previous studies in Saudi Arabia and Turkey that the chances of finding a parent-child match is relatively high [2]. In these countries the probability for all

patients to have a matched sibling due to consanguine marriages is reported to be 60% [4]. There has been no such incidences reported yet in Bahrain. This pattern of occurrence would be expected to be seen in significantly higher percentages due to the similar marriage culture that prevails in the gulf region and North Africa.

Though our case becomes uniquely different with very closely matched parent and children having a high degree of compatibility. Further studies are suggested to come to a broader conclusion whether this is an isolated case or more of such cases would be found to be a trend in the Bahraini population.

References

1. Sung Yoon Choo (2007) The HLA System Genetics, Immunology, Clinical Testing, and Clinical Implications. Yonsei Medical Journal 48: 11-13.
2. Hajeer A, Algattan M, Anizi A, Ahmed S Alaskar, Mohammad S Jarrar, et al. (2012) Chances of finding a matched parent-child in hematopoietic stem cell transplantation in Saudi Arabia. American journal of blood research 2: 201-202.
3. Hamamy, Hanan (2012) Consanguineous marriages Preconception consultation in primary health care settings. J Community Genet 3: 185-192.
4. Jawdat DM, Al Saleh S, Sutton P, Hanan Al Anazi, Abdullah Shubaili, et al. (2009) Chances of finding an HLA matched sibling: the Saudi experience. Biol Blood Marrow Transplant 15: 1342-1344.
5. OLERUP ASSIGN™ SBT v471 Operator Manual, OLERUP SBT HLA Typing Kits instructions for use PCR Amplification and Sequencing of HLA Class I and II Loci. CareDx Pty Ltd. Fremantle, Australia: OLERUP ASSIGN™, October 20, 2017.