Research Article

Open Access

Prediction of Breast Cancer Risk Factors Using Neural Network Analytics: an Empirical Study

Ahed J Alkhatib^{1,2}, Shadi M. Alkhatib³, Mahmoud Atef Ali Ramadan⁴, Ali Atef Ali Ramadan⁵ and Khaled Alshihabat⁶

¹Department of Legal Medicine, Toxicology and Forensic Medicine, Jordan University of Science & Technology, Jordan

²International Mariinskaya Academy, department of medicine and critical care, department of philosophy, Academician secretary of department of Sociology.

³Department of Software Engineering, Hashemite University, Zarqa, Jordan

⁴Computer Information Systems, Al Balqa Applied University

⁵Prince Hussein bin Abdullah College for Information Technology, Al al-Bayt University

⁶Cyprus International University

ABSTRACT

Breast cancer is the most prevalent cancer affecting women. Identifying breast cancer risk factors are crucially to be established. The main objectives of the present study were to identify the predictors of breast cancer risk factors and their relative importance using neural network analysis. The present study depended on neural network analysis of data posted on [1]. The dataset is about predictors of breast cancer. There were 9 covariates included and one dependent variable, output (no disease (1), or disease (2)). The dataset was composed of 116 cases. The category of no disease comprised 79 (68.1%) cases, whereas the disease category included 37 cases (31.9%). Architecture model was built with some characteristics such as training part: gross entropy error was 23.7884, the percent of incorrect predictions was 10.1%. Stopping rule used was 1 consecutive step(s) with no decrease in error. For testing part, gross entropy error was 14,327, the percent of incorrect predictions was 13.5%. The relative importance of breast cancer was in the following order: glucose, resistin, BMI, age, leptin, adiponectin, MCP-1, insulin, and HOMA. Taken together, neural network analysis is an efficient tool to predict breast cancer risk factors.

*Corresponding author

Ahed J Alkhatib, Department of Legal Medicine, Toxicology and Forensic Medicine, Jordan University of Science & Technology, Jordan. Tel no: 00962795905145 E-mail: ajalkhatib@just.edu.jo

Received: January 03, 2021; **Accepted:** January 11, 2021; **Published:** January 14, 2021

Keywords: Breast Cancer, Risk Factor, Neural Network Analytics, Dataset, Architecture Model.

Introduction

Breast cancer is considered the highest prevalent tumor affecting women within the menopause period that usually causes women to lack the ability to cope with routine life functions. Due to the consideration that the incidence of breast cancer has increased over the time, it is crucially to investigate the risk factors that lead to breast cancer [2]. Breast cancer is considered the primary cause of death among women, particularly under the age of 40. Genetic predispositions are important factors affecting young women with breast cancer [3,4]. Other researchers reported that breast cancer impacts 1 woman of 300 before 40 and attributes to approximately 7% of all breast cancer cases [5]. However, the trend of increasing incidence of breast cancer among young women has been reported by various researchers [6,7,8].

In the present study, we investigated some risk factors that can lead to

breast cancer. Leptin is one of the known factors linking obesity with breast cancer by promoting tumor initiation, growth, and metastasis [9]. Leptin is likely to interact with signaling molecules such as ER α , growth factors, Notch, and inflammatory factors that increases the likelihood of cancer occurrence [10,11,12]. Other studies have shown that both leptin and resistin are able to induce signals of inflammatory nature within different immune cells [13,14,15]. On the other hand, adiponectin exhibits anti-inflammatory properties [16].

Inflammation related to obesity mediates the progression of metabolic syndrome ending with insulin resistance [17]. The case of insulin resistance leads to accumulation of insulin as well as growth factors such as insulin-like growth factor 1 (IGF-1), leading to the development of type 2 diabetes [18]. Homeostatic model assessment (HOMA) is usually employed to evaluate the function of β -cell and insulin resistance [19]. The monocyte chemoattractant protein-1 (MCP-1/CCL2) belongs to the C-C chemokine family, and it is considered as an influential attractive factor for monocytes [20,21].

Study Objectives

The main objectives of the present study were to identify the predictors of breast cancer risk factors and their relative importance using neural network analysis.

Materials and Methods

The present study was based on neural network analysis of data posted on [1]. The dataset is about predictors of breast cancer. There were 9 covariates included and one dependent variable, output (no disease (1), or disease (2)). The dataset was composed of 116 cases. The category of no disease comprised 79 (68.1%) cases, whereas the disease category included 37 cases (31.9%).

Case Processing Summary

Using neural network analytics implied two primary training and testing of data. As shown in table 1, a summary of case processing was provided. Training part was performed for 79 (68.1%) cases, whereas testing part was performed for 37 (31.9%) cases. The test was performed for all cases without excluding for any case.

Table 1: Case processing summary

Variable			N	Percent
Sample	Training	Training	79	68.1%
	Testing	Testing	37	31.9%
Valid			116	100.0%
Excluded			0	
Total			116	

The aim of performing neural network analytics is to predict the impact of covariates, or independent variables on the output layer, or dependent variable.

Network Information

As shown in Table 2, network information included three layers, input layer, hidden layer, and output layer. Input layer included the following independent variables, or covariates: age, BMI, glucose, insulin, HOMA, leptin, adiponectin, resistin, and MCP1. Hidden layer (s) included standardized rescaling method for covariates, one hidden layer, five units in hidden layer, and the activation

function was hyperbolic tangent. The output layer included the dependent variable that involved classification of two responses. Softmax was the activation function, whereas error function was represented by cross-entropy.

Table 2: Network information

Input Layer	Covariates		1	Age
			2	BMI
			3	Glucose
			4	Insulin
			5	HOMA
			6	Leptin
			7	Adiponectin
			8	Resistin
			9	MCP1
	Number of Unitsa		9	
	Rescaling Method for Covariates		Standardized	
Hidden Layer(s)	Number of Hidden Layers		1	
	Number of Units in Hidden Layer 1a		5	
	Activation Function		Hyperbolic tangent	
Output Layer	Dependent Variables	1	Classification	
	Number of Units		2	
	Activation Function		Softmax	
	Error Function		Cross-entropy	
a. Excluding the bias unit				

Results

Building the Architecture Model of the Study

As shown in figure 1, the covariates of the study were interacted with hidden layers to give the prediction of disease. Two main colors were drawn, the gray and blue lines. Of each color, there are different intensities and sizes that represent the computed influence of the interacted variable.

Figure 1: A diagram showing the interactions of architecture model

Model Summary

As shown in table 3, the main features of the model were summarized as for training part: gross entropy error was 23.7884, the percent of incorrect predictions was 10.1%. Stopping rule used was 1 consecutive step(s) with no decrease in errora. For testing part, gross entropy error was 14,327, the percent of incorrect predictions was 13.5%.

Table 3: Model Summary

Training	Cross Entropy Error	23.784
	Percent Incorrect Predictions	10.1%
	Stopping Rule Used	1 consecutive step(s) with no decrease in errora
	Training Time	0:00:00.04
Testing	Cross Entropy Error	14.327
	Percent Incorrect Predictions	13.5%
Dependent Variable: Classification		
a. Error computations are based on the testing sample.		

Classification of Output Layer

As shown in Table 4 and Figure 2, in the training part, 37 normal cases were involved, among which 4 cases were predicted as diseased. On the other hand, 42 diseased cases were involved, among which 4 cases were predicted to be normal. The overall percent correct was 89.9%. For training part, 15 normal cases were included, among which 2 cases were predicted as diseased. For testing part, 22 diseased cases were included, among which 3 cases were predicted as normal. The overall percent correct was 86.5%.

Table 4: Classification of output layer

Sample	Observed	Predicted		
		1	2	Percent Correct
Training	1	33	4	89.2%
	2	4	38	90.5%
	Overall Percent	46.8%	53.2%	89.9%
Testing	1	13	2	86.7%
	2	3	19	86.4%
	Overall Percent	43.2%	56.8%	86.5%
Dependent Variable: Classification

Figure 2: Predicting pseudo-probability of output layer classification

Independent Variable Importance

As indicated in table 5 and Figure 3, the independent variable importance was summarized in the following order: glucose (100%), resistin (44.7%), BMI (41.3%), age (32.3%), leptin (29.3%), adiponectin (25.6%), MCP1 (19.2%), insulin (18.1%), and HOMA (10.7%).

Table 5: Independent Variable Importance

Covariates	Importance	Normalized Importance
Age	.101	32.3%
BMI	.129	41.3%
Glucose	.311	100.0%
Insulin	.056	18.1%
HOMA	.033	10.7%
Leptin	.091	29.3%
Adiponectin	.080	25.6%
Resistin	.139	44.7%
MCP1	.060	19.2%

Figure 3: Normalized importance of covariates on breast cancer

Discussion

Breast cancer is a very important medical topic that has impact on the health of women (Kamińska et al., 2015; Bray et al., 2018; Seidler and Huber, 2020) [2,3,4]. We investigated the breast cancer risk factors using neural network analysis. The model we used included 9 risk factors for breast cancer: age, BMI, glucose, insulin, HOMA, leptin, adiponectin, resistin, and MCP-1. According to the predictions made by the model created in this study, glucose was the most important risk factor of breast cancer. Glucose increases the likelihood of cells for proliferation that increased the probability of developing breast cancer [22]. Resistin was ranked as the second most important risk factor for breast cancer. It is not surprising that resistin has a significant impact on diabetes because resistin has inflammatory signals that led to initiation of breast cancer. This agrees with other studies [13,14,15].

BMI ranked the third breast risk factor. However, BMI is associated with developing obesity and increased glucose levels [22]. Age in this study was the fourth breast cancer risk factor. Age has been regarded as a genetic predisposing factor for developing breast cancer risk factor [3,4]. Leptin was the fifth breast cancer risk factor. It has been reported that leptin can link obesity with breast cancer [9]. It can interfere with several signaling molecules such as ERα, growth factors, Notch, and inflammatory factor, a process that increases the probability of developing breast cancer [10,11,12].

Adiponectin is the sixth ranked breast cancer risk factor that is associated with breast cancer. The role of adiponectin in breast cancer looks controversial. Adiponectin exhibits anti-inflammatory properties [16]. In this case, it is difficult to link adiponectin with breast cancer. However, in their study, showed that adiponectin

is a breast cancer risk factor prior to menopausal period [23]. The results showed that MCP-1 as the seventh breast cancer risk factor. MCP-1 has been reported to increase invasiveness and metastasis of breast cancer [24].

Insulin is the eighth breast cancer risk factor. Insulin is associated with obesity related inflammation that increases the probability of breast cancer [17]. HOMA was the last breast cancer risk factor. HOMA has been reported in several studies to increase the probability of developing breast cancer because it is associated with increased status of having insulin resistance [25].

Conclusion

The present study showed the relative importance of breast cancer risk factors from the most important such as glucose to the least important such as HOMA.

References

1. Breast Cancer Coimbra Data Set (2021) <https://www.kaggle.com/yasserhessein/breast-cancer-coimbra-data-set>.
2. Marzena Kamińska, Tomasz Ciszewski, Karolina Łopacka-Szatan, Paweł Miotła, Elżbieta Starosławska (2015) Breast cancer risk factors. *Prz Menopauzalny* 14: 196-202.
3. Bray F, Jacques Ferlay, Isabelle Soerjomataram, Rebecca L Siegel, Lindsey A Torre, et al. (2018) Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians* 68: 394-424.
4. Stephanie J Seidler, DE Huber (2020) Overview of Diagnosis and Treatment of Breast Cancer in Young Women. *EC Gynaecology* SI 02: 18-25.
5. Hussein A Assi, Katia E Khoury, Haifa Dbouk, Lana Khalil,

- Tarek Hassaan Mouhieddine, et al. (2013) Epidemiology and prognosis of breast cancer in young women. *Journal of Thoracic Disease* 5: 2-8.
6. Bouchardy C, G Fioretta, H M Verkooijen, G Vlastos, P Schaefer, et al. (2007) Recent increase of breast cancer incidence among women under the age of forty. *The British Journal of Cancer* 96: 1743-1746.
7. Leclère B, Florence Molinié, Brigitte Trétarre, Fabrizio Stracci, Laetitia Daubisse-Marliac, et al. (2013) Trends in incidence of breast cancer among women under 40 in seven European countries: a GRELL cooperative study. *Cancer Epidemiology* 37: 544-549.
8. Bodmer A, Anita Feller, Andrea Bordoni, Christine Bouchardy, Silvia Dehler, et al. (2015) Breast cancer in younger women in Switzerland 1996-2009: A longitudinal population-based study. *The Breast* 24: 112-117.
9. Barone I, Giordano C, Bonofiglio D, Andò S, Catalano S (2016) Leptin, obesity and breast cancer: progress to understanding the molecular connections. *Curr Opin Pharmacol* 31: 83-89.
10. Saxena NK, Taliaferro-Smith L, Knight BB, Didier Merlin, Frank A Anania, et al. (2008) Bidirectional crosstalk between leptin and insulin-like growth factor-I signaling promotes invasion and migration of breast cancer cells via transactivation of epidermal growth factor receptor. *Cancer Res.*, 68: 9712-9722.
11. Capasso I, Esposito E, Pentimalli F, Montella M, Crispo A, et al. (2013) Homeostasis model assessment to detect insulin resistance and identify patients at high risk of breast cancer development: National Cancer Institute of Naples experience. *Journal of experimental & clinical cancer research: CR*, 32.
12. Lipsey CC, Harbuzariu A, Daley-Brown D, Gonzalez-Perez RR (2016) Oncogenic role of leptin and Notch interleukin-1 leptin crosstalk outcome in cancer. *World J Methodol* 6: 43-55.
13. Silswal N, Singh A K, Aruna B, Mukhopadhyay S, Ghosh S, et al. (2005) Human resistin stimulates the pro-inflammatory cytokines TNF-alpha and IL-12 in macrophages by NF-kappaB-dependent pathway. *Biochem. Biophys. Res. Commun* 334: 1092-1101.
14. Iikuni N, Lam Q L, Lu L, Matarese G, La Cava A (2008) Leptin and Inflammation. *Curr. Immunol. Rev* 4: 70-79.
15. Tripathi D, Kant S, Pandey S, Ehtesham N Z (2020) Resistin in metabolism, inflammation, and disease. *FEBS J* 287: 3141-3149.
- Sun S, Sun Y, Rong X, Bai L (2019) High glucose promotes breast cancer proliferation and metastasis by impairing angiotensinogen expression. *Bioscience reports* 39: 20190436.
16. Fang H, Judd R L (2018) Adiponectin regulation and function. *Compr. Physiol*, 8: 1031-1063.
17. Monteiro R, Azevedo I (2010) Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm* 2010: 289645.
18. S E, Hull RL, Utzschneider K M (2006) Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* 444: 840-846.
19. Tara M Wallace, Jonathan C Levy, David R Matthews (2004) Use and Abuse of HOMA Modeling. *Diabetes Care* Jun 27: 1487-1495.
20. Cochran BH, Reffel AC, Stiles CD (1983) Molecular cloning of gene sequences regulated by platelet-derived growth factor. *Cell* 33: 939-947.
21. Satish L, Deshmane, Sergey Kremlev, Shohreh Amini, Bassel E Sawaya (2009) Monocyte Chemoattractant Protein-1 (MCP-1): An Overview. *J Interferon Cytokine Res* 29: 313-326.
22. Sun S, Sun Y, Rong X, Bai L (2019) High glucose promotes breast cancer proliferation and metastasis by impairing angiotensinogen expression. *Bioscience reports* 39: 20190436.
23. Liu L Y, Wang M, Ma Z B, Yu L X, Zhang Q, et al. (2013) The role of adiponectin in breast cancer: a meta-analysis. *Plos one* 8: 73183.
24. Dutta P, Sarkissyan M, Paico K, Wu Y, Vadgama J V (2018) MCP-1 is overexpressed in triple-negative breast cancers and drives cancer invasiveness and metastasis. *Breast cancer research and treatment* 170: 477-486.
25. Newman G, Gonzalez-Perez RR (2014) Leptin-cytokine crosstalk in breast cancer. *Mol Cell Endocrinol* 382: 570-582.