Case Report

Open Access

Cervical Lipoblastoma in a Child: An Extremely Rare Localisation

Hicham Lyoubi*, Adil Lekhbal, Omar Wydadi, Mouna Lyoubi, Youssef Oukessou, Reda Allah Abada, Sami Rouadi, Mohammed Roubal and Mohammed Mahtar

ENT and Head and Neck Surgery, Hospital August 20, 1953, Ibn Rochd Teaching Hospital, Casablanca–Morocco

ABSTRACT

Lipoblastoma is a benign neoplasm, which occurs usually in infants and children less than 3 years. It is usually seen on the trunk and on the limbs and rarely in the head and neck. In our case, a child of age 9 months presented with a cervical lower left mass. The mass extends from the upper edge of the clavicle to the left side of the hyoid bone.

Magnetic resonance imaging (MRI) showed a well-encapsulated mass, hypointense on the T1 sequence. Histologically, the analysis identified fat lobules with collections of lipoblasts and areas of myxoid degeneration. The mass was resected without notable complications.

*Corresponding author

Hicham Lyoubi, ENT and Head and Neck Surgery, Hospital August 20, 1953, Ibn Rochd Teaching Hospital, Casablanca–Morocco. E-mail: Hicham.lyoubi23@gmail.com

Received: October 23, 2020; **Accepted:** October 27, 2020; **Published:** November 11, 2020

Keywords: Head and Neck Region, Lipoblastoma, Child

Introduction

Lipoblastoma is a benign neoplasm of fetal white fat tissue, which occurs usually in infants and children less than 3 years. Less than 100 cases of lipoblastoma have been reported in literature till now, with involvement of head and neck being reported in very few cases [1]. We are reporting a case of cervical lipoblastoma in a child.

Case Report

A 9 month old female child was brought by his parents with complaints of a big swelling of the neck. They had noticed a small swelling at birth, but during the last 6 months there was a rapid increase in size. It was not associated with pain or fever. The child had a restriction of neck movement. There was no personal history or similar cases in the family.

The mass extends from the upper edge of the clavicle to the left side of the hyoid bone without any inflammatory skin signs. On palpation, the mass was non-fluctuating, firm, non-reducible and mobile. The mass was 45 × 32 × 14 mm. Movements of the neck were limited.

Figure 1: Picture of the cervical mass

Magnetic resonance imaging (MRI) showed a well-encapsulated mass, hypointense on the T1 sequence. On the fat suppression sequence, the corresponding mass demonstrated a weak homogeneous signal intensity.

Histologically, the analysis identified fat lobules with collections of lipoblasts and areas of myxoid degeneration. The mass was resected without notable complications. The follow up is over 1 year, and has not showed any complications or recurrence.

Discussion

Jaffe coined the term “lipoblastoma” in 1926 to describe an atypical lipomatous lesion that consisted of cells resembling embryonic white fat [2]. It usually occurs in children under 3 years of age with a predilection for the extremities and can rarely occur in other places such as the head and neck, as seen in our patient. Men are 3 times more frequently than women [3, 4].

Patients with lipoblastoma may present with various symptoms depending upon its site [5]. The imaging modality of choice to aid in diagnosis is a T1-weighted MRI showing hyperintense signals [6]. Rasmussen et al., mentioned a case of cervical lipoblastoma causing intermittent airway obstruction [7]. The differential diagnosis for lipoblastoma includes cystic hygroma and adipose-derived tumors [8]. The management of lipoblastomas requires complete excision of the tumor while preserving the vital organs in the process. Although there is a high recurrence rate with incomplete resections, it depends on the extent of the resection [9].

Conclusion

Although lipoblastoma is a rare, benign tumor, it should be considered in the diagnosis of a neck mass with stridor in pediatric patients.

Acknowledgements

We like to acknowledge the patient and his relatives for accepting their case to be published. The study was funded by the authors themselves.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to

The research, authorship, and/or publication of this article.

References

1. Premkumar K, Basle MA1, Jassim K, Waseem Ahamed TP (2015) An unusual case of cervical lipoblastoma with review of literature. J Can Res 11: 1025
2. Jaffe RH (1926) Recurrent lipomatous tumors of the groin: Liposarcoma and lipoma pseudomyxomatodes. Arch Pathol 1: 381 387.
3. Ghosh P, Das RN, Ghosh R, Chatterjee U, Datta C, et al. (2015) Lipoblastoma and lipoblastomatosis: a clinicopathological study of six cases. J Can Res Ther 11: 1040.
4. Abdul-Ghafar J, Ahmad Z, Tariq MU, Kayani N, Uddin N (2018) Lipoblastoma: a clinicopathologic review of 23 cases from a major tertiary care center plus detailed review of literature. BMC Res Notes. 11: 42.
5. Dilley AV, Patel DL, Hicks MJ, Brandt ML (2001) Lipoblastoma: Pathophysiology and surgical management. J Pediatr Surg 36: 229 231.
6. Petersen C, Ure B (2017) Thoracic Surgery in Children and Adolescents [e-book]. Berlin, Germany: De Gruyter.
7. Rasmussen IS, Kirkegaard J, Kaasbol M (1997) Intermittent airway obstruction in a child caused by a cervical lipoblastoma. Acta Anaesthesiol Scand 41: 945 946.
8. Rosen A, Jedynak AR, Respler D (2005) Lipoblastoma of the neck mimicking cystic hygroma. Otolaryngol Head Neck Surg 132 : 511-513.
9. Elder DE (2015) Lever's Histopathology of the Skin. 11th ed. Philadelphia, PA: Wolters Kluwer Health.

Copyright: ©2020 Hicham Lyoubi, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.