Research Article

Open Access

Telomere Length and Glutathione Level as Potential Biomarkers for Autism

Ayat M Abdaltawab¹, Nagwa A. Meguid², Eman M. Khaled¹, Adel Hashish², Mohamed H. Bahary³, Mohamed S. Taha^{2*}

¹Department of Pediatrics, Faculty of medicine, Al-Azhar University, Cairo, Egypt

²Department of Research on Children with Special Needs, Medical Research Division, National Research Centre, Giza, Egypt

³Department of Psychiatry, Faculty of medicine, Al-Azhar University, Cairo, Egypt

ABSTRACT

Oxidative stress is a common feature in nearly all neurodegenerative disorders including autism spectrum disorder (ASD). ASD is behaviorally defined disorder associated with abnormalities in social communication, behavior, language and perception. Chronic exposure to different common stress factors like immune inflammation, malnutrition, alcohol, abuse and drugs induce congenital and behavioral impairment. To the best of our knowledge this study is the first exploring the rate of oxidative stress and its effect on telomere length simultaneously in Egyptian autistic children. In this study we investigated the effect of oxidative stress associated with ASD. Patients were selected based on full clinical and neurological examination and the severity of autism evaluated using childhood autism rating scale (CARS) and Autism Diagnostic Interview-Revised (ADI[™]-R). Expectedly we found the majority of autism affected children were males, and consanguinity rate between the ASD families was ~ 27% while it was completely absent in the control subjects. We found a highly significant correlation between ASD and low levels of total glutathione ($P < 0.001$) compared to controls. We reported a remarkable highly significant correlation between ASD children and shorter relative telomere length ($P < 0.001$). Moreover we identified a correlation between telomere length and behavioral characteristics in autistic children represented by social interaction impairment (ADI-R 1st domain) ($r = -0.354$, $P = 0.025$). Here we conclude that oxidative stress represented by low glutathione level is an early sign of autism, and is affecting the telomere length that protecting the genetic material. Behavioral abnormalities associated with autism are dependent on telomere length.

*Corresponding author

Mohamed S. Taha, Department of Research on Children with Special Needs, Medical Research Branch, National Research Centre, El Buhouth St, 12622, Cairo, Egypt; E-mail: ms.taha@nrc.sci.eg.

Received: June 01, 2021; **Accepted:** June 07, 2021; **Published:** June 10, 2021

Keywords: Autism Spectrum Disorder - Neurodevelopmental- Oxidative Stress- Telomere- Glutathione

Introduction

Autism spectrum disorder (ASD) is a complex neurodevelopmental syndrome, which usually occurs at early childhood (~2 years of age). In 1943 Autism first identified by Leo Kanner as an innate inability to create normal, biologically determined, emotional contact with others [1,2]. In general, ASD is characterized by pervasive deficits in social interaction, impairments in verbal and nonverbal communication, and stereotyped patterns of interests and activities [3]. ASD varies in severity and the clinical phenotype reflects multifactorial background [4]. ASD prevalence was 18.5 per 1,000 (one in 54) children aged 8 years (ADDM Network sites) in the United States [5]. ASD prevalent by about 4 times higher in boys than in girls [5-7]. The estimated prevalence of ASD among the neurodevelopmentally disabled children in Egypt was to be 33.6% [8]. Recently both telomere length and oxidative stress are considered a promising biomarker for early a neurodevelopmental evaluation [9].

Telomeres are nucleoprotein structures localized at the end of chromosomes that function to protect chromosomes from degradation. Telomeres are formed from repetitive noncoding

deoxynucleotide sequences composed of conserved sequence of TTAGGG and associated proteins located at chromosomal ends [10]. This telomeric repeats synthesized by telomerase activity [11]. Telomere length (TL) is affected by both genetic and environmental factors, and shortening of telomeres is associated with multiple neuropsychiatric disorders, early life stress, and age-related cognitive dysfunction [11,12]. Considering that telomere shortening has been associated with some psychiatric diseases, there is a possibility that dysfunctional telomeres might be involved in autism pathogenesis and that altered telomere length may be associated with childhood autism [13].

Glutathione (GSH) is a tripeptide, γ -L-glutamyl-L cysteinyl glycine, ubiquitously exist in all mammalian tissue types and crucially in liver tissue with high concentrations [14]. GSH is considered as the most abundant non-protein thiol that defends against oxidative stress. GSH is also a key determinant of redox signaling, vital in detoxification of xenobiotics, modulates cell proliferation, apoptosis, immune function, and fibrogenesis [15]. Glutathione can exist in either thiol-reduced (GSH) or disulfide-oxidized (GSSG) forms. The antioxidant reaction of GSH can be summarized in two oxidation and reduction reactions, where the oxidation of GSH to GSSG is accompanied by the reduction of hydrogen peroxide and lipid peroxide. GSSG in turn is reduced

back to GSH by GSSG reductase at the expense of NADPH, forming a redox cycle [15]. In brain, glutathione play an important cellular protective function by scavenging reactive oxygen species (ROS) and consequently is involved in various cellular survival pathways in response to oxidative stress [16]. To the best of our knowledge this study is the first to investigate the rate and progression of oxidative stress in Egyptian autistic children, by analyzing the tGSH and RTL.

Subject & Methods

Subjects: This study included 80 children, 40 of them are affected by childhood ASD and the other 40 are healthy with no history of any major neurodegenerative disease in their families. ASD children were recruited from autism disorders clinic. Autistics diagnosed by a senior clinician in neurology and human genetics. Patients' consents were read, fully comprehended and signed by tutoring parents of the recruited pediatric subjects. The typically-developing (TD) controls sampled from one common kindergarten with no abnormal histories of motor, language, or social developmental disorders, as determined according to reports of parents. Children in both groups were matched in age (3 – 6 years) and sex. Detailed history of patients has been recorded, including three generation pedigree analysis, pregnancy history, perinatal history, developmental history, similarly affected family members, and age of onset of presenting manifestation. Clinical diagnosis performed based on the criteria for autistic disorder as defined in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-IV) [17,18]. And Autism Diagnostic Interview-Revised (ADI-R) [19]. The Childhood Autism Rating Scale (CARS) was conducted for assessing autism severity [20]. The Autism Diagnostic Interview – Revised (ADI-R) was employed on three domains: (Qualitative abnormalities in reciprocal social interaction), (Qualitative abnormalities communications), and (Restricted repetitive & stereotyped pattern of behavior). Patient with other neurodegenerative diseases e.g. fragile-X syndrome, Asperger's syndrome, Rett syndrome, and patients with childhood schizophrenia have been excluded from this study.

Telomere Length Measurements: Blood samples were collected into EDTA tube (BD Vacutainer®) by following standard laboratory procedure. Samples were centrifuged at 1200 rpm for 10 minutes. Cells pellet were used for DNA isolation from leukocyte cell using Qiagen DNeasy Blood & Tissue isolation kit (Qiagen, Germany). Isolated DNA samples were stored at -20C for later relative telomere length analysis (RTL). RTL was measured by the quantitative real-time polymerase chain reaction (PCR) method originally described in details elsewhere [21]. Real time PCR was performed using DTlite Real-Time PCR System (DNA technology, Russia). Using the same primers and protocol previously described in [22]. A comparative quantitation approach was used to measure the telomere length regarding a relative ratio (T/S ratio) of telomere (T) repeat copy number to a single copy of reference gene (S) copy number.

Glutathione Concentration Measurements: Blood samples were collected using heparin as anticoagulant. Total glutathione level was detected in blood erythrocytes of patient and control samples simultaneously by following the manufacturer protocol using OxiSelect™ Total Glutathione (GSSG/GSH) Assay Kit (STA-312) (Cell Biolabs, USA).

Statistical Analysis: Recorded data were analyzed using the statistical package for social sciences, version 20.0 (SPSS Inc., Chicago, Illinois, USA). Quantitative data were expressed as mean $\pm$ standard deviation (SD). Qualitative data were expressed

as frequency and percentage. Comparing proportion was done using Pearson's correlation coefficient (r) and regression analysis. Receiver operating characteristic (ROC curve) analysis was used to find out the overall sensitivity and specificity. All P-values were two-sided and considered statistically significant at $P < 0.05$, and highly significant at $P < 0.001$.

Results

Subject Group Analysis

The present study included 80 children with age range 2-6 years old. They divided equally into two groups autistics and controls, each group contain 40 children. The autistic group (ASD children) was evaluated according to the criteria defined in the Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-IV), Childhood Autism Rating Scale (CARS), and Autism Diagnostic Interview-Revised (ADI™-R). The control group is healthy children with no history of any associated diseases. Both groups are matched in age and sex, the autistics were (70% males & 30% females) with age mean of 4.28 ± 1.09 . While the control group includes (62.5% males & 37.5% females) with age mean of 4.54 ± 0.99 . The parents' consanguinity marriage crucially reported in autistic group.

Clinical Manifestation

Autistics were categorized into three categories according to the severity of autism associated symptoms based on the CARS score. About two third of autistics (62.5%) had mild to moderate autism with CARS score of (30 to 37), 20% with sever autism (CARS score > 37) and 17.5% with mild autism (CARS score = 30) (Fig 1A). in addition to CARS scores, the autistics were reevaluated clinically by Autism Diagnostic Interview-Revised (AIDR). Both methods showed a similar results for ASD symptoms classification, and we found a significant direct correlation ($R^2 = 0.376$ & $P < 0.001$) as shown in the linear regression curve (Figure 1B). Demographic and clinical characteristics including height, weight and head circumference values were recorded for autistic group. We didn't find a significant difference in the severity of symptoms between patients based on the demographic and clinical characteristics e.g. sex, height, weight or head circumference.

Figure 1: Clinical analysis of autistic group: A) three categories of autism severity based on CARS score, where 7 patients (17.5%) were mild with CARS score ($= 30$), 25 patients (62.5%) were moderate to severe with CARS score of (30-37) and 8 patients (20%) were severe with CARS score of (38-60). B) Scatter plot showing a linear regression and a significant correlation between CARS scores and the mean of the three domains of ADI-R (Qualitative abnormalities in reciprocal social interaction & Qualitative abnormalities communications, and Restricted repetitive with stereotyped pattern of behavior) scores for patient group ($R^2 = 0.376$ & $P < 0.001$).

Relative Telomere Length And Glutathione Levels In Autism

The mean of relative telomere length (T/S) in the ASD group (mean $\pm$ SD) (0.66 ± 0.11) was significantly shorter than the mean RTL in controls (0.8 ± 0.18), ($P < 0.001$) (Figure 2A & Table 1).

Similarly the ASD group showed a lower tGSH level (6.21 ± 1.12) in comparison to controls (7.76 ± 1.32), ($P < 0.001$) (Figure 2B & Table 2). The mean of RTL in the autistic females (0.65 ± 0.09) was significantly shorter than those females in control group (0.78 ± 0.16), $P = 0.025$ and the tGSH level for the females in ASD group was (6.11 ± 1.13) less than those females in control group (7.17 ± 1.23), ($P = 0.03$). In the same way the males in ASD group showed shorter RTL (0.66 ± 0.11) than the males in control group (0.81 ± 0.19), $p < 0.001$, and the tGSH level was much lower (6.26 ± 1.14) among males in ASD versus those in controls (8.12 ± 1.27), $p < 0.001$ (Table 1&2). There is no correlation between gender identity and RTL in both autistic and control groups ($r = 0.045$,

$P = 0.79$ & $r = 0.08$, $P = 0.62$) respectively. Although the tGSH had a significant correlation with gender in control group only ($r = 0.35$, $P = 0.026$) (Table 3). RTL and tGSH showed a direct positive correlation in the ASD group using linear regression analysis and Pearson correlation ($r = 0.499$, $P^{**} = 0.001$) (Figure 2C). In autistic group the RTL and tGSH showed no correlation with age ($r = 0.198$, $P = 0.221$) & ($r = 0.275$, $P^{*} = 0.086$) respectively. The severity of autism symptoms based on CARS score hadn't any correlation with RTL or tGSH (Table 1&2). Similarly RTL and tGSH levels were both not correlated to patient height, weight or head circumference (Tables 4&5). RTL showed an inverse correlation with qualitative abnormalities in reciprocal social

Figure 2:Distribution of relative telomere length (RTL) and glutathione (tGSH) in patient and control groups: A) Relative telomere length in autistic group and control group (mean $\pm$ SD), showing a significant telomere shortening in autistic group $P < 0.001$. B) Total glutathione level in autism and control group (mean $\pm$ SD), showing lower level of glutathione in autistic group. C) Linear regression showing a significant correlation between relative telomere length and glutathione in autistic group ($R^2 = 0.29$ & $P < 0.001$). ** = highly significant

Figure 3: Glutathione and relative telomere length as biomarker for autism: A) Scatter plot showing a non-significant correlation between Qualitative abnormalities in reciprocal social interaction (ADI-R 1st domain) and tGSH in autistic group, $P = 0.221$. B) Scatter plot showing a significant correlation between Qualitative abnormalities in reciprocal social interaction (ADI-R 1st domain) and RTL, $P = 0.025$. C) Receiver operating characteristic curves analysis of glutathione (red, continuous) area under the curve (AUC) = 0.828 and relative telomere length (blue, dotted) (AUC=0.732). The ROC analysis showing both tGSH and RTL are good biomarkers for autism with higher specificity and sensitivity.

Table 1: Relative telomere length showed a significant difference between autistic and control groups.

	RTL (relative telomere length) (T/S)				
	Autistics		Control		P-value
	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	
females	0.65 ± 0.09	0.48 - 0.82	0.78 ± 0.16	0.59 - 1.04	0.025*
males	0.66 ± 0.11	0.49 - 0.84	0.81 ± 0.19	0.49 - 1.21	<0.001**
Mixed	0.66 ± 0.11	0.48 - 0.84	0.8 ± 0.18	0.49 - 0.21	<0.001**

SD: standard deviation, * : statistically significant at $P < 0.05$, ** : highly significant at $P < 0.001$

Table 2: Total glutathione level showed a significant difference between autistic and control groups

	tGSH (total glutathione level) (μmol/g Hb)				
	Autistics		Control		P-value
	Mean $\pm$ SD	Range	Mean $\pm$ SD	Range	
females	6.11 ± 1.13	4.3 - 7.8	7.17 ± 1.23	4.3 - 9	0.03*
males	6.26 ± 1.14	3.8 - 8.2	8.12 ± 1.27	4.2 - 9.9	<0.001**
Mixed	6.21 ± 1.12	3.8 - 8.2	7.76 ± 1.32	4.2 - 9.9	<0.001**

*: statistically significant at $P < 0.05$, **: highly significant at $P < 0.001$, SD: standard deviation.

Table 3: Correlation between (tGSH, RTL) and gender type in both autistics and control groups

	RTL (relative telomere length) (T/S)		tGSH (total glutathione level) (μmol/g Hb)	
	r	P-value	r	P-value
Autistics (Gender)	0.045	0.79	0.062	0.706
Control (Gender)	0.081	0.62	0.353	0.026*

*: statistically significant at $P < 0.05$, r: Pearson's correlation coefficient

Table 4: Correlation between telomere length relative (T/S) and clinical parameters, using Pearson Correlation Coefficient (r) in the active group.

Parameters	Telomere length Relative (T/S)	
	r	p-value
Age (years)	0.198	0.221
Weight %	-0.068	0.679
Height %	0.03	0.853
Head circumference %	-0.099	0.544
CARS	0.012	0.939
Qualitative abnormalities in reciprocal social interaction	-0.035	0.831
Qualitative abnormalities communications	0.02	0.902
Restricted repetitive & stereotyped pattern of behavior	0.038	0.815

r: Pearson's correlation coefficient

Table 5: Correlation between glutathione level (μmol/g Hb) and clinical parameters, using Pearson Correlation Coefficient in the active group.

Parameters	Telomere length Relative (T/S)	
	r	p-value
Age (years)	0.275	0.086
Weight %	-0.117	0.471
Height %	-0.037	0.822
Head circumference %	-0.249	0.122
CARS	0.163	0.314
Qualitative abnormalities in reciprocal social interaction	0.354	0.025*
Qualitative abnormalities communications	0.213	0.186
Restricted repetitive & stereotyped pattern of behavior	0.096	0.555

*:Statistically significant at $P < 0.05$, r: Pearson's correlation coefficient

RTL and Glutathione Level as Biological Biomarker in Autism
Both the RTL and tGSH were evaluated as biological biomarkers using statistical analysis and receiver operating characteristics (ROC) curves to analyze the sensitivity and specificity of both

biomarkers for autism. RTL (AUC = 0.732, (95% CI 0.623 - 0.841), $P < 0.001$, tGSH (AUC = 0.828 (95% CI 0.735 - 0.92), $P < 0.001$.

Discussion

It was reported that both oxidative stress and mitochondrial dysfunction are common features of neurodegeneration [23-25]. In this study, we analyzed the possible association of oxidative stress with one of the most common neurodevelopmental disorders (Autism spectrum disorder, ASD) among Egyptian children. Autistics in this study were evaluated using the Childhood Autism Rating Scale (CARS) for reconfirming the diagnosis of ASD and to assess the severity of ASD symptoms (Figure 1A). In addition to, using the Autism Diagnostic Interview – Revised (ADI-R). ADI-R is a highly recognized evaluative measure for diagnosing Autism Spectrum Disorder (ASD). The ADI-R provides a helpful standardized approach to gathering diagnostic information. In this study, abnormal restricted repetitive & stereotyped patterns of behavior have the highest percentage of cases. In addition, we found that severe CARS is associated with the highest percentage of restricted repetitive & stereotyped patterns of behavior [26]. Consequently, we noted a significant direct correlation between CARS and ADI-R scores in the patient group (Figure 1B).

Here we evaluated the peripheral blood leukocytes relative telomere length as a biomarker for oxidative stress in autism. Telomeres are nucleoprotein structures at the end of chromosomes and are required for integrity and stability of chromosome ends and prevent them from being recognized as DNA double-strand breaks, through their binding with an adequate amount of a six-unit protein complex (shelterin) [27]. Shelterin protein has a specific and crucial role in telomere maintenance and function. Shortening or absence of telomere due to age or oxidative stress leading to a subsequent decrease in the amount of shelterin protein at chromosome ends and further increasing the vulnerability of DNA to degradation and instability [28,29]. Our results are confirming that relative telomere length (RTL) (telomere length in base pair compared to a reference gene (T/S)) is shorter in autistics compared to the control group (Figure 1B). Autism patients have elevated inflammatory activity and abnormal oxidative stress which make them susceptible to short telomeres [30]. Telomere length shortening is a remarkable biomarker for other neurodegenerative diseases like Alzheimer disease confirming the role of environmental factors and chronic stress caused in inducing age related as well as neurodegenerative diseases [31-33]. We reported no significant difference in RTL between males and females in the control group (Table 3). Considering the hypothesis that females possess a more efficient antioxidant defense mechanism more than males and consequently a longer RTL than males. A reasonable explanation for our result that detecting gender effect on telomere length requires a larger number of subjects of significantly older age on other hand we noted that previous studies performed on a small group of subjects and relatively young individual (or children) like our study have been failed to detect gender effect on RTL [13,30,34-38].

Moreover, we reported a significantly low level of total glutathione (tGSH, GSH/GSSG) concentrations in autistics (Figure 2A) [39,40]. Glutathione is the major antioxidant in nearly all mammalian tissue types, with an exceptionally abundant concentration in liver tissues due to detoxification processes [15]. And it is necessary for the neutralizing of

reactive oxygen species (or free radicals), as well as the regeneration of other antioxidants such as vitamins C and E [40]. In the brain, GSH is a key determinant of cell signaling, cell proliferation, cell differentiation, protein function, apoptosis, immune function, fibrogenesis, and gene expression [40,41]. Reported that glutathione plays an important role in the brain as an antioxidant, however, it has been found that the GSH levels are the same in the brain of adult men with and without ASD, using magnetic resonance spectroscopy (MRS) [42]. Moreover glutathione reductase level in blood was the same in control and autistics [39]. In the control group, we found that males have slightly more glutathione levels than females ($p = 0.026$). It was previously reported that females have lower GSH levels and higher GSSG levels than males [43]. Females have more level of GPx (glutathione peroxidase), GPx is important for oxidizing GSH into GSSG as well as for catabolizing organic hydroperoxides and H_2O_2 [44]. Accordingly, females' mitochondria produce significantly less hydrogen peroxide than males [45,46]. Interestingly, taking the hypothesis that oxidative stress precedes and causing autism by inducing telomere shortening and DNA damage, this could be an additional explanation for why the majority of autistics are males (70% males and 30 % females) [30,47]. In the autistics group, we didn't report any correlation between gender identity and tGSH level (Table 3).

tGSH had no significant correlation with the severity of autism or with the patient clinical features including CARS, ADI-R scores, weight, height, and head circumference. This may agree with the hypothesis that oxidative stress is preceding and causing autism. While we reported an inverse correlation between RTL and ADI-R 1st domain (qualitative abnormalities in reciprocal social interaction. ($r = -0.354$, $P = 0.037$). A correlation between telomere length and sensory symptoms of autism reported in previous studies [12]. Taking the hypothesis that chronic oxidative stress can induce telomere shortening and DNA damage the leading to autism progression [48].

Regardless of the previously reported inverse correlation between age and either tGSH or RTL in normal individuals, we found no correlation between age and neither tGSH nor RTL in the autistic group [49,50]. The corresponding p-values for the correlation between either RTL or (tGSH) and age were ($P = 0.22$ & $P = 0.447$) respectively (Table 4&5) (Figure 3 A&B). Interestingly, several studies including us reported no correlation between age and oxidative stress markers (RTL & tGSH) among affected groups (e.g. neurodegeneration, cardiovascular and diabetes) [30,51,52]. Here we hypothesis that oxidative stress in autistic is masking and dominate the age-related oxidative stress, so we were no longer able to detect age effect inactive groups (autistic). We can also consider the technical difficulty in recording the variations among critically short RTL or low level of tGSH that human body maintains to avoid apoptotic e.g. (ferroptotic) [53,54]. On other hand, chronic exposure to some toxins and pollution is producing a significant effect on glutathione level and was detectable among autistics [40].

We evaluated tGSH and RTL as a possible biomarker for autism using receiver operating characteristics (ROC) curves. A cut off for RTL at ≤ 0.72 is giving sensitivity of 72.5%, specificity of 65%. While in tGSH level a cut off value at ≤ 7.1 producing sensitivity of 80%, and specificity was 75%. We found that tGSH and RTL are a potential biomarker for autism, and maybe recommended for future studies on evaluating the effect of dietary adequacy on oxidative stress as well as improving autism associated symptoms [55,56].

Conclusions

In conclusion it was highly mandated to analyze the extent of oxidative stress in autistic children who lives in a developing country like Egypt. Where, the high rate of consanguinity and daily life burden that inducing additional stress. We found that oxidative stress associated to autism has a significant effect on telomere length and glutathione level among autistic Egyptian children. We found no correlation between tGSH levels and autism symptoms. Nevertheless the RTL showed an inverse correlation with some autism symptoms. This finding supports the hypothesis that oxidative stress is preceding and causing autism by affecting the integrity of human genome including telomeres. ROC analysis confirmed the higher sensitivity and specificity between tGSH and autism. Finally we are suggesting both RTL & tGSH as potential biomarkers for autism in early childhood.

Highlights

- Versatile types of stressors affecting mental as well as physical health of adults and adolescents.
- Oxidative stress represented by low glutathione level is suggested to be a leading cause of autism.
- Shorter relative telomere length is a common feature in autism.
- Both tGSH and RTL are recommended to be a biological biomarkers in Egyptian autistic children.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval: The whole study is approved by the Ethics Committee of National Research Centre, Giza, Egypt.

Acknowledgements: we are grateful to the Autism Spectrum Disorders Clinic, Medical Research Center of Excellence, at the National Research Centre for their support.

Conflicts of Interest: The authors declare that they have no competing interests.

References

1. Waight PA, Andrews NJ, Ladhani SN, Sheppard CL, Slack MP, Miller E (2015) Effect of the 13-valent pneumococcal conjugate vaccine on invasive pneumococcal disease in England and Wales 4 years after its introduction: an observational cohort study. *Lancet Infect Dis*, 15: 535-543.
2. Hategan A, Bourgeois JA, Goldberg J (2017) Aging with autism spectrum disorder: an emerging public health problem. *Int Psychogeriatr*, 29: 695-697.
3. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders 5th edn American Psychiatric Association.
4. Hallmayer J, Cleveland S, Torres A, Phillips J, Cohen B, et al. (2011) Genetic heritability and shared environmental factors among twin pairs with autism. *Arch Gen Psychiatry*, 68: 1095-1102.
5. Maenner MJ, Shaw KA, Baio J, Washington A, Patrick M, et al. (2020) Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years - Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2016. *MMWR Surveill Summ* 69: 1-12.
6. Loomes R, Hull L, Mandy WPL (2017) What Is the Male-to-Female Ratio in Autism Spectrum Disorder? A Systematic Review and Meta-Analysis. *J Am Acad Child Adolesc Psychiatry* 56: 466-474.
7. Baig U, Mehdi SM, Imtiaz Afzal HM (2021) Sleep disturbances in children with autism spectrum disorder in

- Lahore, Pakistan; a cross-sectional study. *J Pak Med Assoc* 71: 783-785.
8. Seif Eldin A, Habib D, Noufal A, Farrag S, Bazaid K, et al. (2008) Use of M-CHAT for a multinational screening of young children with autism in the Arab countries. *Int Rev Psychiatry* 20: 281-289.
 9. Provenzi L, Scotto di Minico G, Giorda R, Montirosso R (2017) Telomere Length in Preterm Infants: A Promising Biomarker of Early Adversity and Care in the Neonatal Intensive Care Unit? *Front Endocrinol (Lausanne)*, 8: 295.
 10. Blackburn EH, Greider CW, Szostak JW (2006) Telomeres and telomerase: the path from maize, Tetrahymena and yeast to human cancer and aging. *Nat Med* 12: 1133-1138.
 11. Turner KJ, Vasu V, Griffin DK (2019) Telomere Biology and Human Phenotype. *Cells* 8: 1-19.
 12. Lewis CR, Taguinod F, Jepsen W M, Cohen J, Agrawal K, et al. (2020) Telomere Length and Autism Spectrum Disorder within the Family: Relationships with Cognition and Sensory Symptoms. *Autism Res* 13: 1094-1101.
 13. Li Z, Tang J, Li H, Chen S, He Y, et al. (2014). Shorter telomere length in peripheral blood leukocytes is associated with childhood autism. *Sci Rep* 4, 7073.
 14. Pizzorno J (2014) Glutathione! *Integr Med (Encinitas)*, 13: 8-12.
 15. Lu SC (2013) Glutathione synthesis. *Biochim Biophys Acta* 1830: 3143-3153.
 16. Zhao S, Zhao X, Dong S, Yu J, Pan G, et al. (2015). Correction: A hierarchical, stretchable and stiff fibrous biotemplate engineered using stagger- electrospinning for augmentation of rotator cuff tendon-healing. *J Mater Chem B* 3.
 17. Roehr B (2013) American Psychiatric Association explains DSM-5. *BMJ*, 346, f3591.
 18. Kim SK (2015) Recent update of autism spectrum disorders. *Korean J Pediatr*, 58: 8-14.
 19. Pilowsky T, Yirmiya N, Shulman C, Dover R (1998) The Autism Diagnostic Interview-Revised and the Childhood Autism Rating Scale: differences between diagnostic systems and comparison between genders. *J Autism Dev Disord*, 28: 143-151.
 20. Schopler E, Reichler RJ (1988) The Childhood Autism Rating Scale (CARS). Western Psychological Services, Los Angeles, CA.
 21. Cawthon RM (2002) Telomere measurement by quantitative PCR. *Nucleic Acids Res*, 30 e47.
 22. O'Callaghan NJ, Fenech M (2011) A quantitative PCR method for measuring absolute telomere length. *Biol Proced Online* 13.
 23. Carpita B, Muti D, Dell'Osso L (2018) Oxidative Stress, Maternal Diabetes, and Autism Spectrum Disorders. *Oxid Med Cell Longev*, 3717215.
 24. Singh A, Kukreti R, Saso L, Kukreti S (2019). Oxidative Stress: A Key Modulator in Neurodegenerative Diseases. *Molecules*, 24.
 25. Taha MS, Haghighi F, Stefanski A, Nakhaei-Rad S, Kazemeini Jasemi NS, et al. (2020) Novel FMRP interaction networks linked to cellular stress. *FEBS J*.
 26. McKinnon CJ, Eggebrecht AT, Todorov A, Wolff JJ, Elison JT, et al. (2019). Restricted and Repetitive Behavior and Brain Functional Connectivity in Infants at Risk for Developing Autism Spectrum Disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging* 4: 50-61.
 27. Montpetit AJ, Alhareeri AA, Montpetit M, Starkweather AR, Elmore LW, et al. (2014) Telomere length: a review of methods for measurement. *Nurs Res* 63: 289-299.
 28. De Lange T (2018) Shelterin-Mediated Telomere Protection. *Annu Rev Genet* 52: 223-247.
 29. Jacobs JJ (2013) Loss of telomere protection: consequences and opportunities. *Front Oncol*, 88.
 30. Gardner M, Bann D, Wiley L, Cooper R, Hardy R, et al. (2014). Gender and telomere length: systematic review and meta-analysis. *Exp Gerontol*, 51: 15-27.
 31. Forero DA (2016) Meta-analysis of Telomere Length in Alzheimer's Disease. *J Gerontol A Biol Sci Med Sci* 71: 1069-1073.
 32. Bernardi RB (2021) The impact of chronic exposure to air pollution over oxidative stress parameters and brain histology. *Environ Sci Pollut Res Int*.
 33. Shueb M (2020) A possible relationship between telomere length and markers of neurodegeneration in rat brain after welding fume inhalation exposure. *Environ Res* 180: 108900.
 34. Aviv A (2004) Telomeres and human aging: facts and fids. *Sci Aging Knowledge Environ* 43.
 35. Benetos A, Okuda K, Lajemi M, Kimura M, Thomas F, et al. (2001). Telomere length as an indicator of biological aging: the gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension* 37: 381-385.
 36. Hewakapuge S, van Oorschot R A, Lewandowski P, Baidur-Hudson S (2008) Investigation of telomere lengths measurement by quantitative real-time PCR to predict age. *Leg Med (Tokyo)* 10: 236-242.
 37. McGrath M, Wong JY, Michaud D, Hunter DJ, De Vivo I (2007) Telomere length, cigarette smoking, and bladder cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev* 16: 815-819.
 38. Das B, Saini D, Seshadri M (2009). Telomere length in human adults and high level natural background radiation. *PLoS One*, 4.
 39. Al-Yafee YA, Al-Ayadhi LY, Haq SH, El-Ansary AK (2011) Novel metabolic biomarkers related to sulfur-dependent detoxification pathways in autistic patients of Saudi Arabia. *BMC Neurol*, 11.
 40. Faber S, Fahrenholz T, Wolle MM, Kern JC, Pamuku M, et al. (2019) Chronic exposure to xenobiotic pollution leads to significantly higher total glutathione and lower reduced to oxidized glutathione ratio in red blood cells of children with autism. *Free Radic Biol Med* 134: 666-677.
 41. Aoyama K, Nakaki T (2013) Impaired glutathione synthesis in neurodegeneration. *Int J Mol Sci* 14: 21021-21044.
 42. Durieux AMS, Horder J, Mendez MA, Egerton A, Williams SCR, et al. (2016). Cortical and subcortical glutathione levels in adults with autism spectrum disorder. *Autism Res* 9: 429-435.
 43. Kowalska K, Milnerowicz H (2016) The Influence of Age and Gender on the Pro/Antioxidant Status in Young Healthy People. *Ann Clin Lab Sci* 46: 480-488.
 44. Hill KE, Taylor MA, Burk RF (1987) Influence of selenium deficiency on glutathione disulfide metabolism in isolated perfused rat heart. *Biochim Biophys Acta* 923: 431-435.
 45. Borrás C, Sastre J, García-Sala D, Lloret A, Pallardó FV (2003) Mitochondria from females exhibit higher antioxidant gene expression and lower oxidative damage than males. *Free Radic Biol Med* 34: 546-552.
 46. Viña J, Sastre J, Pallardó F, Borrás C (2003) Mitochondrial theory of aging: importance to explain why females live longer than males. *Antioxid Redox Signal* 5: 549-556.
 47. Wang X, Lu J, Xie W, Lu X, Liang Y, et al. (2019) Maternal

- diabetes induces autism-like behavior by hyperglycemia-mediated persistent oxidative stress and suppression of superoxide dismutase 2. *Proc Natl Acad Sci U S A* 116: 23743-23752.
48. Sarkar T, Patro N, Patro IK (2019) Cumulative multiple early life hits- a potent threat leading to neurological disorders. *Brain Res Bull* 147: 58-68.
 49. Liu J, Mori A (1993) Age-associated changes in superoxide dismutase activity, thiobarbituric acid reactivity and reduced glutathione level in the brain and liver in senescence accelerated mice (SAM): a comparison with ddY mice. *Mech Ageing Dev* 71: 23-30.
 50. Shay JW (2018) Telomeres and aging. *Curr Opin Cell Biol* 52: 1-7.
 51. Kota LN, Bharath S, Purushottam M, Moily NS, Sivakumar PT, et al. (2015). Reduced telomere length in neurodegenerative disorders may suggest shared biology. *J Neuropsychiatry Clin Neurosci* 27: e92-96.
 52. Rodrigues B, Nóbrega AC, Sampaio M, Argolo N, Melo A (2011) Silent saliva aspiration in Parkinson's disease. *Mov Disord* 26: 138-141.
 53. Guiney SJ, Adlard PA, Bush AI, Finkelstein DI, Ayton S (2017) Ferroptosis and cell death mechanisms in Parkinson's disease. *Neurochem Int* 104: 34-48.
 54. Zhang Q, Wu H, Zou M, Li L, Li Q, et al. (2019) Folic acid improves abnormal behavior via mitigation of oxidative stress, inflammation, and ferroptosis in the BTBR T+ tf/J mouse model of autism. *J Nutr Biochem* 71: 98-109.
 55. Thanseem I, Viswambharan V, Poovathinal SA, Anitha A (2017) Is telomere length a biomarker of neurological disorders? *Biomark Med* 11: 799-810.
 56. Meguid NA, Anwar M, Bjørklund G, Hashish A, Chirumbolo S, et al. (2017). Dietary adequacy of Egyptian children with autism spectrum disorder compared to healthy developing children. *Metab Brain Dis* 32: 607-615.