PREVALENCE OF ANTIENDOMYSIAL ANTIBODIES IN SJÖGREN'S SYNDROME PATIENTS

Laís Zanolorenzi^{1,*}, Naiana Diaz, Matheus Burkot Alves de Araújo¹, Matheus Vinicius Conte Laba¹, Aiessa Fedrigo¹, Thelma Larocca Skare¹

1.Hospital Universitário Evangélico Mackenzie, Curitiba (SP), Brazil.

*Corresponding author: laiszn@gmail.com

BACKGROUND

Autoimmune disease may occur in the same patient probably due to a shared genetic background or to a common environmental exposure. Literature shows that celiac disease (CD) and/or its autoantibodies are present in Sjögren's syndrome (SS) patients in a prevalence ranging from 1% to 14.7%. The studies on SS prevalence in patients with celiac disease diagnosis ranges from 1.2% to 5%. The prevalence differs according to studied population as both diseases have influence of genetic background. The aim of this study was to study the prevalence of antiendomysial antibodies (anti-EmA-IgA) in a sample of Brazilian patients with SS.

METHODS

The search of EmA-IgA in SS patients was done by indirect immunofluorescence technique using indirect anti-IgA fluorescent conjugate (Euroimmun, Brazil) and compared with the prevalence of local general population (blood donor's) from literature. Positive patients answered the gastrointestinal symptom rating scale (GSRS) questionnaire for gastrointestinal symptoms.

RESULTS

Thirty-nine SS patients were included: 28 (71.7%) with primary SS (pSS) and 11 (28.3%) with secondary SS (sSS) (9 associated with systemic lupus erythematosus and 2 associated with rheumatoid arthritis). In this sample, there was one patient with known CD diagnosis; other 2 had a positive EmA-IgA. All positive patients had pSS and had some minor gastrointestinal complaint not properly valued previously: 66% complained of loose stools, 66% of passing gas or flatus; 33% of urgency to evacuate and 33% of the sensation of not completely emptying bowel. Comparison of the prevalence of EmA-IgA (3/39 or 7.6%) with the prevalence in general population (1/286), SS patients had an OR = 23.7; 95%CI = 3.4–234.0; $p = 0.005$ for positive EmA-IgA. Upper gastrointestinal endoscopy was offered to them but the patients postponed it because of COVID-19 pandemic.

CONCLUSION

The present work suggests a high prevalence of coexistence of these two diseases.

KEYWORDS

Antiendomysial antibodies, Sjögren's syndrome, Celiac disease.