Caliphate Medical Journal

A Publication of Nigerian Medical Association Sokoto State Chapter.

Vol 9, No. 4, October - December, 2021

ISSN: 2346-7193

In this Issue

CMJ

Also available @ www.cmj.nmasokoto.org

PATTERN OF COMPLEMENTARY AND ALTERNATIVE MEDICINE (CAM) USE AMONG HYPERTENSIVE PATIENTS IN NORTHERN NIGERIA

Jimoh AO¹, Faragai AY², Sani Z³, Tukur UM¹, Hudu SA⁴

¹Department of Pharmacology and Therapeutics, Faculty of Basic Clinical Sciences, College of Health Sciences, Usmanu Danfodiyo University Sokoto, Nigeria.

²Faculty of Pharmaceutical Sciences, Usmanu Danfodiyo University Sokoto, Nigeria.

³Department of Family Medicine, Usmanu Danfodiyo University Teaching Hospital Sokoto Nigeria.

⁴Department of Medical Microbiology and Parasitology, Faculty of Basic Clinical Sciences, College of Health Sciences, Usmanu Danfodiyo University Sokoto, Nigeria.

ABSTRACT

Introduction: Hypertension is the most common cardiovascular diseases worldwide. The frequency of utilization of CAM is increasing worldwide, and is well documented in both African and other global populations.

Objective: The objective of this study was to investigate the prevalence and correlates of CAM use among hypertensive patients.

Methods: The study was conducted at the department of medicine of Murtala Muhammad General Hospital of Kano State, Nigeria. A cross-sectional study was conducted, on 208 patients, using a semi structured questionnaire on the use of CAM among hypertensive patients. Data analysis was done using Microsoft Office Excel 2007 and SPSS V16.

Result and Conclusion: The frequency of CAM use in this study was (67.8%) and the use of CAM was significantly associated with marital status, duration of diagnosis, knowledge of antihypertensives and ability to pay for drugs, $p < 0.05$.

Keywords: Hypertension, Cardiovascular diseases, CAM, Anti hypertensive

Corresponding Author's:

Abdulgafar O. Jimoh,

Department of Pharmacology and Therapeutics,
Faculty of Basic Clinical Sciences,
College of Health Sciences,
Usmanu Danfodiyo University Sokoto, Nigeria.
Email: abdulgafarjimoh@gmail.com,
jimoh.abdulgafar@udusok.edu.ng,
Phone: +2348035950558

Introduction

Hypertension is a clinical and family health problem¹ and is the principal cardiovascular disease worldwide.² It is defined currently as systolic and/or diastolic blood pressures $\geq 140/90$ mmHg in tandem with The Seventh Report of the Joint National Committee on prevention, detection, evaluation and treatment of high blood pressure in adults aged 18 years and older.⁴ In recent years there has been increasing concern about the burden of non-communicable diseases especially cardiovascular diseases and particularly hypertension in sub-Saharan African countries.⁴ In Nigeria, hypertension is the most common non-communicable and cardiovascular disease.⁵ The consequences of this preventable epidemic of hypertension will be substantial on the individual and family dynamics resulting in individual and family suffering, stress, distress and dysfunction.⁶ The prevalence of hypertension in populations has been reported to differ geographically⁷ since it may be influenced by family socio-biological and other diverse factors.⁸

In Nigeria, the estimated prevalence by World Health Organization was 42.8%.² The prevalence range of 8%-46% was reported in a systematic review of prevalence of hypertension in adult Nigerians⁵, 22.5% -31.0% was reported in Abia state non-communicable disease survey⁵, 26.4% was reported among patients with family history of cardiovascular diseases in Umuahia, Abia state⁵ and 42.7%, were reported among obese adult in primary care clinic in South-east Nigeria⁶, and 18.4%, was reported in obese adult patients in semi-urban hospital in Eastern Nigeria.⁶ Complementary and Alternative Medicine is any practice that is perceived by its users to have the healing effects of medicine, but does not originate from evidence gathered using the scientific method^{9,11}, is not part of biomedicine^{9,11,13} or is contradicted by scientific evidence or established science.^{9,11}

The frequency of utilization of CAM is increasing worldwide, and is well documented in both African and other global populations to be between 20 - 80%.¹⁴ In an ethnobotanical survey conducted by (Nurudeen *et al.*, 2013)

at Sokoto, Nigeria, it was shown that a large number of medicinal plants are used in Sokoto state to treat hypertension.

The objective of this study was to investigate the prevalence and correlates of CAM use among hypertensive patients attending hypertension clinic of the Murtala Muhammad General Hospital of Kano State, Nigeria.

Methodology

Study design

A cross-sectional descriptive study design was used to address the research goals. The study population were interviewed using a semi structured questionnaire.

Study Area

Kano is a city in Nigeria and the capital of Kano State in Northern Nigeria, in the Sahelian geographic region south of the Sahara. Kano is the commercial nerve center of Northern Nigeria and is the second largest city in Nigeria after Lagos. According to the 2006 census, Kano is the most populous state in Nigeria, with about 9,383,682 million people. The Kano urban area covers 137 km².

Study Site

The study was conducted in hypertension outpatients' clinic at Murtala Muhammad General Hospital of Kano State, Nigeria. It is a government owned General Hospital in Kano Municipal local government area of Kano State, Nigeria. It is located at Muhammad Abdullai Wase Way, Kano, Kano State. It offers general care to Kano populace as well as other people from outside the states.

Sample Size Calculation

Calculation of sample size for this study is based on the findings of Trevisol DJ *et al.*, 2000²⁰, that the percentage of patients with controlled hypertension has been reported to vary between 5.4% and 58% worldwide, with minimum number of 78 samples.

Data Collection Procedure

A pilot study was carried out to ensure a comprehensive analysis for a range of perspectives, to detect any possible problems associated with the format, wording, and measurement, and to make sure that the respondents comprehend the instructions and questions. Data was collected using a set of questionnaires. Three main variables were used; 1. Socio-demographic data, 2. Clinical hypertension related variables, 4. CAM profile of hypertensive patients.

Data Analysis

The data collected was entered and analysed using Microsoft Office Excel 2007 and SPSS V16. The descriptive analysis was done using the Microsoft Office Excel 2007. While, the statistical tests of association were done using the SPSS V16. Ethical approval for the study was sought from the hospital research and ethics committee.

Results

The mean age of the study population (mean $\pm$ SD) was 50.1 $\pm$ 10.7 years. Age of the patients ranged from 21-80, with the majority 67 (32.2%) falling within 41-50 years of age. One Hundred and twenty-five (60.1%) were female patients. Majority of the patients 62 (29.8%) had a monthly income of ₦11,000-₦50,000 only. Majority 174 (83.6%) of the patients were married and 73 (35.1%) had tertiary level of education (Table 1).

Table 1: Socio-Demographic and Clinical-Related Variables of the Study Population

Variables	N=208	Percentage
Age		
21-30	8	3.8
31-40	42	20.2
41-50	67	32.2
51-60	59	28.4
61-70	2	0.9
71-80	6	2.9
Sex		
Female	125	60.1
Male	83	39.9
Monthly Income		
≥₦100,000	13	6.3
₦51,000-₦100,000	40	19.2
₦11,000-₦50,000	62	29.8
≤₦10,000	42	20.2
Blanks	51	24.5
Marital Status		
Married	174	83.6
Divorced	6	2.9
Widow	28	13.5
Single	-	-
Family Size		
Nuclear	132	63.5
Extended	61	29.3
Blanks	15	7.2
Educational Level		
Non-formal	53	25.5
Primary	14	6.7
Secondary	67	32.2
Tertiary	73	35.1
Blanks	1	0.5

The duration of the illness of the majority of the patients 84 (40.4%) fall within 2-5 years. Majority 182 (87.6%) of the patients were on once daily drug dosage and most 187 (87.5%) of the patients took their antihypertensive agents once daily. The medication bills of 126 (60.6%) patients were paid by their families. Also, 185 (88.9%) of the patients had not experience any kind of adverse drug reactions with

their medications (Table 2). More than half of the patients were reported to be CAM users (Figure 1).

Table 2: Clinical-Related Variables of the Study Population

Variables	N=208	Percentage
Duration of Diagnosis		
2-5 Years	84	40.4
6-10 Years	72	34.6
≥11 Years	52	25.0
Dosage Frequency of the Anti Hypertensive Agent		
Once Daily	182	87.5
Twice Daily	26	12.5
Mode of Payment for the Anti Hypertensive Agent		
Self	82	39.4
Family	126	60.6
Improvement with Anti Hypertensive Agent use		
Yes	207	99.5
No	-	-
Blanks	1	0.5
Unto Wards Effects with Anti Hypertensive Agent use		
Yes	25	12.1
No	183	87.9

Figure 1: The prevalence of CAM Use among the Study Population.

Majority of those using CAM were of the age range of 41-50 years (31.9%), female (64.5%), married (78.7%) and of secondary level of education (33.3%). However, only being married was found to be statistically associated with CAM use (Table 3).

The result also shows that majority of those using CAM were diagnosed with hypertension more than 10 years away from the time of the study (79.8%), were also reported to not have been improving with their anti hypertensive medications (77.7%), and relied on their families for their medications (73.8%). All these were found to be statistically associated with CAM use (Table 4).

Discussion

The majority of the patients were of the age range of 41-50. This is expected as the onset of systemic hypertension is usually within this age bracket. This is in contrast to a study conducted by.¹⁷ Surprisingly, even though essential hypertension is higher in males than females, majority of the population were female. This is similar (65.9%) to a study conducted by.¹⁸

The ₦11,000-₦50,000 monthly income of the study

Table 3: Relationship between Socio-Demographic data and CAM Use

Variables	N (208)	CAM Use		χ^2	P-value
		Yes (141)	No (67)		
Age				1.061	0.588
21-30	8	5(3.5%)	39(4.5%)		
31-40	42	24(17.0%)	18(26.9%)		
41-50	67	45(31.9%)	22(32.8%)		
51-60	59	37(26.2%)	23(34.3%)		
61-70	25	25(17.7%)	-		
71-80	6	5(3.5%)	1(1.5%)		
Sex				3.603	0.069
Male	125	50(35.5%)	33(49.3%)		
Female	83	91(64.5%)	34(50.7%)		
Monthly Income				3.358	0.187
≥100,000	13	6(4.3%)	7(10.4%)		
51,000-100,000	40	24(17.0%)	16(23.9%)		
11,000-50,000	62	38(26.9%)	24(35.8%)		
≤10,000	42	31(21.9%)	11(16.4%)		
Blanks	51	42(29.8%)	9(13.4%)		
Marital Status				7.635	0.022*
Married	174	111(78.7%)	63(94.0%)		
Divorced	6	4(2.8%)	2(3.0%)		
Widow	28	26(18.4%)	2(3.0%)		
Single	0	-	-		
Educational Level				5.977	0.113
Non-formal	53	41(29.1%)	12(17.9%)		
Primary	14	10(7.1%)	4(5.9%)		
Secondary	67	47(33.3%)	20(29.9%)		
Tertiary	73	42(29.8%)	31(46.3%)		
Blanks	1	1(0.7%)	-		

*There is statistically significant relationship between the variable and CAM use $p < 0.05$

population is in line with the average monthly earning of many Nigerians. The fact that majority of the patients were married is also in line with the tradition of the Hausa community. Usually, people within the age group of the patients should be married unless there is strong reason not to. The finding that majority of the patients had tertiary level of education is an indication of how the contemporary people of Hausa land embrace Western education.

The frequency of CAM use in this study (67.8%) is less than what was reported in Sokoto Nigeria (84.0%)¹⁹, and is higher to what was reported by a study conducted at Lagos University Teaching Hospital (39.1%)²⁰ and higher than the one obtained in the study conducted in Ibadan Nigeria (29%).²¹ The reported frequencies of CAM use in the general population (irrespective of disease entity) from different parts of the world include 40% in the US²², 38.5% among the Indian community of Chatsworth in South Africa²³ and 48.5% in Australia.²⁴ However, this finding is similar to what Shafiq *et al* reported in a cohort study that as many as 63.9% of their hypertensive subjects in a clinic in India took herbal medicines²⁵, while in Morocco 80% of patients with hypertension and diabetes used medicinal plants to treat their ailments.²⁶ The high prevalence of CAM use in our study is likely because before the advent of orthodox medicine in Nigeria, herbal medicines had been the

Table 4: Relationship between Clinical Variables and CAM Use

Variables	N (208)	CAM Use		χ^2	P-value
		Yes (141)	No (67)		
Duration Of Diagnosis				7.635	0.023*
<5 Years	55	35(63.6%)	20(36.4%)		
5-10 Years	101	63(62.4%)	38(37.6%)		
>10 Years	84	67(79.8%)	17(20.2%)		
Anti Hypertensive Knowledge				10.870	0.001*
Yes	96	54(56.3%)	42(43.7%)		
No	112	87(77.7%)	25(22.3%)		
Mode Of Payment				5.306	0.021*
Self	82	48(58.5%)	34(41.5%)		
Family	126	93(73.8%)	33(26.2%)		
Improvement With Anti Hypertensive				0.477	0.490
Yes	207	140(67.6%)	67(32.4%)		
No	1	1(100%)	0(0%)		
Unto Wards Effects with Anti Hypertensive Use				1.808	0.179
Yes	25	14(9.9%)	11(16.4%)		
No	183	127(90.1%)	56(83.6%)		

*There is statistically significant relationship between the variable and CAM use $p < 0.05$;
CAM = Complementary and Alternative Medicine

mainstay of treatment for various ailments and were dispensed by traditional herbalists involved in their cultivation and preparation. The observed high use of herbal remedies in this study may be linked to this cultural background and history.

This study found no statistically significant relationship between CAM use and socio-demographic variables with exception of marital status. This is similar to a study carried out in hypertensive patients attending an urban tertiary care centre in Nigeria²⁰ where CAM use was found to be independent of demographic variables. However, it differs from the finding of a study conducted at a tertiary hospital in Imo State, South-East, Nigeria²⁷ where age, marital status, educational level and monthly income were the determinants of CAM use in the study. Our finding also differs from differ from the American and Australian studies, which both found several associations between demographic factors and CAM usage.^{22,24} Notably college education and wealth were predictors common to both the US and Australian studies.

This study found statistically significant relationship between CAM use and duration of the disease, $p < 0.05$. Majority of CAM users as displayed in table 4 were those diagnosed with hypertension more than 10 years ago from the time of the study. However, this is differs from the study carried out in hypertensive patients attending an urban tertiary care centre in Nigeria²⁰ where CAM use was found to be independent of duration of the illness, duration of clinic attendance or initial B.P at presentation. The statistically significant relationship between CAM use and late onset of the disease is expected. This probably happened because the patients were disappointed with conventional treatment and wanted to try everything that may help them.

CAM use is associated with low knowledge of antihypertensives and family funding of medications, $p < 0.05$. Low knowledge level and poverty are two factors that can predispose patients to increase CAM use.

Conclusion

This study has reinforced a high prevalence of CAM use among hypertensive patients in a northern Nigerian community. marital status, duration of hypertension, knowledge of antihypertensives and ability to pay for antihypertensives associated with increase CAM use.

Limitations

A relatively small sample size. The study only targeted the outpatients diagnosed with essential hypertension and treated for hypertension in the General Hospital of Kano metropolis.

References

1. Chuku A, Gabriel U, Pascal I, Nnadozie PO, John NO, Augustin OI: Frequency of cardiovascular risk factors in adult Nigerians with family history of non-communicable cardiovascular disease in a primary care clinic of a tertiary hospital in a resource-constrained environment of eastern Nigeria. *American Journal of Health Research* 2013; 1(1): 17-25.
2. Mendis S, Johnston SC, Fan W, et al. Cardiovascular risk management and its impact on hypertension control in primary care in low- resource settings: a cluster-randomized trial. *Bull World Health Organ* 2010;88:412-19.
3. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones 8DW, Materson BJ, Oparil S, Wright JT Jr, Roccella E J . The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* 2003; 289: 2560-72.
4. Twagirumukiza M, De Bacquer D, Kips JG, de Backer G, Stichele RV & Van Bortel LM. Current and projected prevalence of arterial hypertension in sub-Saharan Africa by sex, age and habitat: an estimate from population studies. *J Hypertens*. 2011; 29(7): 1243-1252.
5. Ogah OS, Okpechi I, Chukwuonye II, Akinyemi JO, Onwubere BJ, Falase AO, et al. Blood pressure, prevalence of hypertension and hypertension related complications in Nigerian Africans: A review. *World J Cardiol*. 2012;4:327-40.
6. Iloh GUP, Amadi AN, Nwankwo BO, Ugwu VC. Medication adherence and blood pressure control amongst adults with primary hypertension attending a tertiary hospital primary care clinic in Eastern Nigeria. *Niger J Clin Pract*. 2011;14:212-218.
7. Kearney PM, Whelton M, Reynolds K, Whelton PK, He J. Worldwide prevalence of hypertension: a systematic review. *J Hypertens* 2004; 22:11-19.
8. Olatunbosun ST, Kaufman JS, Cooper RS, Bella AF (2000). Hypertension in black population: prevalence and biosocial determinants of high blood pressure in a group of urban Nigerians. *J. Human Hypertension* 14(4)249-257.
9. Akpa MR, Alasia DD, Emem-chioma PC. An appraisal of hospital based blood pressure control in Port Harcourt, Nigeria. *Nigerian Health J*. 2008;8:27-30.
10. Sampson W (1995). "Antiscience Trends in the Rise of the "Alternative Medicine 'Movement". *Annals of the New York Academy of Sciences* 775 (1): 188-197.
11. Beyerstein BL (2001). "Alternative medicine and common errors of reasoning". *Academic medicine : journal of the Association of American Medical Colleges* 76 (3): 230-237.
12. Terence, Hines. *Pseudoscience and the Paranormal* (2nd ed.) (2003). Amerst, New York: Prometheus Books.
13. Nissen, N.; et al. (2013). "Researching alternative and complementary therapies: Mapping the field". *Medical Anthropology* 32(1): 1-7.
14. Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR, Delbanco TL: Unconventional Medicine in the United States – Prevalence, Costs, and Patterns of Use. *N Engl J Med* 1993, 328:246-252.
15. Nurudeen O R, Iyabo M A, Shaibu O B. Ethnobotanical Survey Of Antihypertensive Agents In Sokoto, Northwest Nigeria. *International Journal of Innovative Research & Development*. 2013, 2:1820-1835.
16. Trevisol DJ, Moreira LB, Kerkhoff A, Fuchs SC, Fuchs FD: Health-related quality of life and hypertension: a systematic review and meta-analysis of observational studies. *J Hypertens* 2011, 29:179-188.
17. Amal KT, Syed AS. Elevated Blood Pressure Among Patients With Hypertension In General Hospital Of Penang, Malaysia: Does Poor Adherence Matter? *International Journal of Pharmacy and Pharmaceutical Sciences* 2010, 2(1):24-32.
18. Jelena L, Radmila VR, Branka M, Valentina N, Ana S, Goran K. Medication Adherence in Outpatients with Arterial Hypertension. *Scientific Journal of the Faculty of Medicine in Niš* 2013, 30(4):209-218.
19. Abdulgafar O. Jimoh, Zuwaira Sani, Kabiru Abubakar, Halilu E. Mshelia. Safety Concerns and Determinants of Complementary and Alternative Medicine Use in a Sub-urban area of Sokoto, North Western Nigeria. *Journal of Medical Sciences*, 2013, 13:737-742.
20. Oluwatoyin C Amira and Njideka U Okubadejo. Frequency of complementary and alternative medicine utilization in hypertensive patients attending an urban tertiary care centre in Nigeria. *BMC Complementary and Alternative Medicine* 2007, 7:30.
21. Pauline EO, Bernard EO: Complementary and alternative medicine in the management of hypertension in an urban Nigerian community. *BMC Complementary and Alternative Medicine* 2010, 10:36.
22. Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M, Kessler RC: Trends in alternative medicine use in the United States, 1990 – 1997: results of a follow-up national survey. *JAMA* 1998, 280:1569-1575.
23. Singh V, Raidoo DM, Harries CS: The prevalence, patterns of usage and people's attitude towards complementary and alternative medicine (CAM) among the Indian community in Chatsworth, South Africa. *BMC Complement* 2004, 4(3):1-7.
24. MacLennan AH, Wilson DH, Taylor AW: Prevalence and cost of alternative medicine in Australia. *Lancet* 1996, 347:569-573.
25. Shafiq N, Gupta M, Kumari S, Pandhi P: Prevalence and pattern of use of complementary and alternative medicine (CAM) in hypertensive patients of a tertiary care center in India. *Int J Clin Pharmacol Ther* 2003, 41(7):294-298.
26. Eddouks M, Maghrani M, Lemhadri A, Ouahidi ML, Jouad H: Ethnopharmacological survey of medicinal plants used for the treatment of diabetes mellitus, hypertension and cardiac diseases in the south-east region of Morocco (Tafilalet). *J Ethnopharmacol* 2002, 82(2-3):97-103.
27. Chukwuma B. Duru, Kenechi A. Uwakwe, Nnebue C. Chinomnso, Ikechukwu I. Mbachi, Kevin C. Diwe, Chuka C. Agunwa, Anthony C. Iwu, Irene A. Merenu. Socio-demographic Determinants of Herbal Medicine Use in Pregnancy Among Nigerian Women Attending Clinics in a Tertiary Hospital in Imo State, South-East, Nigeria. *American Journal of Medicine Studies* 2016, 4:1-10.