COPY RIGHT

ELSEVIER

SSRN

2021 IJIEMR. Personal use of this material is permitted. Permission from IJIEMR must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collective works, for resale or redistribution to servers or lists, or reuse of any copyrighted component of this work in other works. No Reprint should be done to this paper, all copy right is authenticated to Paper Authors

IJIEMR Transactions, online available on 13th Apr 2021. Link

[:http://www.ijiemr.org/downloads.php?vol=Volume-10&issue=ISSUE-04](http://www.ijiemr.org/downloads.php?vol=Volume-10&issue=ISSUE-04)

DOI: 10.48047/IJIEMR/V10/I04/44

Title: **Tadpole graphs and their laceability.**

Volume 10, Issue 04, Pages: 172'-176

Paper Authors

Annapoorna M S, Murali R, Manjunath G

USE THIS BARCODE TO ACCESS YOUR ONLINE PAPER

To Secure Your Paper As Per **UGC Guidelines** We Are Providing A Electronic Bar Code

Tadpole graphs and their laceability

Annapoorna M S¹, Murali R² and Manjunath G³

^{1*} Department of Mathematics, BMS Institute of Technology & Management, Bangalore.

Email: anuraga@bmsit.in

² Department of Mathematics, Dr. Ambedkar Institute of Technology, Bangalore.

Email: muralir2968@dr-ait.org

³ Department of Mathematics, Rajiv Gandhi Institute of Technology, Bangalore.

Email: ghanshan.1981@gmail.com

ABSTRACT

A connected graph G is termed Hamiltonian- t -laceable (t^* -laceable) if there exists in G a Hamiltonian path between every pair (at least one pair) of its vertices u and v with the property $d(u, v) = t$. The Tadpole graph is the graph obtained by joining a cycle graph C_m to a path graph P_n with a bridge. In this paper, we discuss the laceability properties associated with the Tadpole graph.

Keywords: Hamiltonian- t -laceability, Hamiltonian- t^* -laceability, Tadpole graph

INTRODUCTION

All graphs considered here are finite, simple, connected and undirected. A graph G is hamiltonian- t -laceable (hamiltonian- t^* -laceable) if there exists in G a hamiltonian path between every pair (at least one pair) of vertices u and v such that $d(u, v) = t$, $1 \leq t \leq \text{diam}G$, where t is a positive integer. G is termed t^* -connected if it is hamiltonian- t^* -laceable for all t such that $1 \leq t \leq \text{diam}G$. Various results on the laceability properties of graph structures like the fan graph, the gear graph, the wheel graph, brick product of even and odd cycles and the cyclic product can be found in [5]. In this paper we study the hamiltonian- t^* -laceability property of the Tadpole graphs.

Definition: The Tadpole graph is the graph obtained by joining a cycle graph C_m to a path graph P_n with a bridge.

Fig 1: The Tadpole graphs $T_{(4,1)}$,

$T_{(5,1)}$, $T_{(n,1)}$, $T_{(4,2)}$ and $T_{(5,3)}$

Definition

Let P be a path from the vertices a_i to a_j in a graph G and let P' be a path from a_j to a_k . Then the path $P \cup P'$ is the path obtained by extending the path P from a_i to a_j to a_k , through the common vertex a_j .

Definition

In a graph $G = (V, E)$, let X be a minimal set of edges not in G and $G' = (V, E \cup X)$ is t^* -connected graph then $|X|$ is called the t^* -laceability number $\lambda^*(t)$ and the edges in X are called the t^* -laceability edges.

RESULTS

We need to introduce the following terminologies.

For a vertex x_i of G , we write:

$$x_i^1[n] = x_i, x_{(i+1)}, x_{(i+2)}, \dots, x_n.$$

$$x_i^{-1}[m] = x_i, x_{(i-1)}, x_{(i-2)}, \dots, x_m.$$

$$\{x_i y_j z_k\}^r[l, m, n] = x_{(i+r)}, y_{(j+r)}, z_{(k+r)}, x_{(i+2r)}, y_{(j+2r)}, z_{(k+2r)}, \dots, x_l, y_m, z_n.$$

$$x_i^{(l, m, n)} = x_{i+l}, x_{i+m}, x_{i+n}.$$

Theorem 1: The graph $G = T_{(n,1)}$ is hamiltonian- t^* -laceable with $\lambda_{(t)} = 1$ for all

t such that $2 \leq t \leq \text{diam}(G)$, where $3 < n \leq 6$

Proof: Let $G = T_{(n,1)}$ with vertex set

$V = \{a_0, a_1, a_2, a_3, \dots, a_n\}$ and edge set

$E = \{a_1 a_2, a_2 a_3, a_3 a_4, \dots, a_{(n-2)} a_{(n-1)}, a_{(n-1)} a_n\} \cup \{a_n a_0\}$, where $3 < n \leq 6$.

Clearly $\text{diam}(G) = 3$.

To establish the results, we consider the following cases,

Case (i): $t = 2$

In $G = T_{(n,1)}$, $d(a_1, a_0) = 2$ and the path, $P: a_1 I^1[n] P^0 a_0$ is the hamiltonian path between the vertices a_1 and a_0 containing the laceability edge (a_1, a_3) .

Case (ii): $t = 3$

In $G = T_{(n,1)}$, $d(a_2, a_0) = 3$ and the path, $P: a_2 I^{-1} I^2 P^0 I^1[n] P^0 a_0$ is the hamiltonian path between the vertices a_2 and a_0 containing the laceability edge (a_1, a_3) .

Hamiltonian path from vertex a_2 to a_0 in the graph $T_{(6,1)}$
Hence the proof.

Theorem 2: The graph $G = T_{(n,1)}$ is hamiltonian- t^* -laceable with $\lambda_{(t)} = 1$ for $t=2,3$ and with $\lambda_{(t)} = 2$ for $t=4$, where $7 \leq n \leq 10$.

Proof: Let $G = T_{(n,1)}$

For $t=2,3$, we follow the proof provided in theorem 1. For $t=4$, we observe that since $\text{diam}(G)=4$,

$d(a_4, a_0) = 4$ in G and the path,

$P: a_4 I^{-1}[1] P^0 I^5 P^0 I^1[n] P^0 a_0$ is

a hamiltonian path between the vertices a_4 and a_0 with the laceability edges (a_1, a_3) and (a_1, a_5) .

Hamiltonian path from vertex a_4 to a_0 in the graph $T_{(9,1)}$

Hence the proof.

Theorem 3: The graph $G = T_{(n,1)}$ is hamiltonian- t^* -laceable with $\lambda_{(t)} = 1$ for $t=2,3$ with $\lambda_{(t)} = 2$ for $t=4$ and with $\lambda_{(t)} = 3$ for $t=5$, where $11 \leq n \leq 13$.

Proof: Let $G = T_{(n,1)}$

For $t=2,3$, we follow the proof provided in theorem 1. For $t=4$, we follow the proof provided in theorem 2 and for $t=5$, we observe that since $\text{diam}(G)=5$, $d(a_7, a_0) = 5$ in G and the path,

$P: a_7 I^{-1}[1] P^0 I^1 P^0 I^6 P^0 I^1[n] P^0 a_0$ is the

hamiltonian path between the vertices a_7 and a_0 with the laceability edges (a_1, a_3) , (a_1, a_5) and (a_2, a_8) .

Hamiltonian path from vertex a_7 to a_0 in the graph $T_{(13,1)}$

Hence the proof.

The following lemma is an immediate consequence of theorem 3.

Lemma 4: The graph $G = T_{(n,1)}$ is hamiltonian- t^* -laceable with $\lambda_{(t)} = 4$ for $n=14$, where $2 \leq t \leq \text{diam}(G)$.

Remark: The graph $T_{(n,1)}$ is t^* -connected with $\lambda^*(t) = 4$ for $n = 14$

Theorem 5: The graph $G = T_{(n,1)}$ is hamiltonian- t^* -laceable with $\lambda_{(t)} = 1$ for $t=2,3$ with $\lambda_{(t)} = 2$ for $t=4$, with $\lambda_{(t)} = 3$

for $t=5$ and with $\lambda_{(t)} = 4$ for $t=6$, where $15 \leq n \leq 18$.

Proof: Let $G = T_{(n,1)}$

For $t=2,3$, we follow the proof provided in theorem 1. For $t=4$, we follow the proof provided in theorem 2, for $t=5$, we follow the proof provided in theorem 3 and we observe that since $diam(G)=6$, $d(a_{10}, a_0) = 6$ in G and the path, $P: a_{10}I^{-1}[5]P^0a_1I^1[4]P^0a_{11}I^1[n]P^0a_0$ is the hamiltonian path between the vertices a_{10} and a_0 with the laceability edges (a_1, a_3) , (a_1, a_5) , (a_2, a_8) and (a_4, a_{11}) .

Hamiltonian path from vertex a_{10} to a_0 in the graph $T_{(18,1)}$

Hence the proof.

Theorem 6: The graph $G = T_{(n,1)}$ is hamiltonian- t^* -laceable with $\lambda_{(t)} = 1$ for $t=2,3$ with $\lambda_{(t)} = 2$ for $t=4$, with $\lambda_{(t)} = 3$ for $t=5$, $\lambda_{(t)} = 4$ for $t=6$ and with $\lambda_{(t)} = 5$ for $t=7$ where $19 \leq n \leq 22$.

Proof: Let $G = T_{(n,1)}$

For $t=2,3$, we follow the proof provided in theorem 1. For $t=4$, we follow the proof provided in theorem 2, for $t=5$, we follow the proof provided in theorem 3, for $t=6$, we follow the proof provided in theorem 5 and for $t=7$ we observe that since $diam(G)=7$, $d(a_{13}, a_0) = 7$ in G and the path, $P: a_{13}I^{-1}[8]P^0a_2I^{-1}P^0a_3I^1[7]P^0a_{14}I^1[n]P^0a_0$ is the hamiltonian path between the vertices a_{13} and a_0 with the laceability edges (a_1, a_3) , (a_1, a_5) , (a_2, a_8) , (a_4, a_{11}) and (a_7, a_{14}) .

Hamiltonian path from vertex a_{13} to a_0 in the graph $T_{(22,1)}$

Hence the proof.

CONCLUSION

In this paper we have explored the hamiltonian- t^* -laceability properties of tadpole graphs $T_{(n,1)}$ for $3 < n \leq 22$ and $2 \leq t \leq diam(G)$.

ACKNOWLEDGEMENT

The first author thankfully acknowledges the support and encouragement provided by the management and staff of the department of Mathematics, BMS Institute of Technology, Bangalore. The authors are also thankful to the management and R & D centre, Department of Mathematics, Dr. Ambedkar Institute of Technology, Bangalore and Rajiv Gandhi Institute of Technology, Bangalore.

REFERENCES

- [1] "**Distance in Graphs**" by Fred Buckley and Frank Harary, Addison-Wesley Pub. Co., 1990 – Mathematics - 335 pages.
- [2] Jain Akiyama and Takashi Hamada and I.Yoshimura, "**On Charecterization of middle graphs**", True Mathematics 11 (1975),pp 35-39.
- [3] VernoldVivin J and K. Thilagavathi, "**On Harmonious Colouring of Line Graph of Central Graph of Paths**", Applied MathematicalmSciences, Vol.3,2009, No.5, 205-214.
- [4] Manjunath. G, R. Murali and Girisha .A, "**Hamiltonian Laceability in Line Graphs**", International Journal of Computer Applications (0975-8887), volume 98-N0.12,July 2014.
- [5]Manjunath. G, Murali.R and B. Shanmukha, "**Hamiltonian Laceability in Middle**

Graphs", International Journal Of Engineering Sciences and Research Technology (203-

209), volume 4 issue 1, January 2015.