Overview of Sleep

S. Chokroverty

Professor and Co-Chair of Neurology; Chairman of Clinical Neurophysiology and Sleep Medicine; Program Director of Clinical Neurophysiology and Sleep Medicine; NJ Neuroscience Institute at JFK; Seton Hall University, Edison, NJ, USA

Indian J Sleep Med 2007; 2.2, 40-49

Introduction

The history of sleep medicine and sleep research is a history of remarkable progress and remarkable ignorance. In the 1940's and 1950's sleep had been in the forefront of neuroscience, and then again in the late 1990's there has been a resurgence of our understanding of the neurobiology of sleep. Sleeping and waking brain cycles can now be studied by sophisticated neuroimaging techniques which have shown remarkable progress by mapping different areas of the brain during different sleep states and stages. Electrophysiologic research has shown that even a single neuron sleeps, as evidenced by the electrophysiologic correlates of sleep-waking at the cellular (single cell) level. Despite recent progress, we are still groping for answers to two fundamental questions: What is Sleep? and Why do we sleep? Sleep is not simply an absence of wakefulness, nor is it just a suspension of sensorial processes but is the result of a combination of a passive withdrawal of afferent stimuli to the brain and functional activation of certain neurons in selective brain areas. The discovery of human electroencephalography (EEG) in 1929 and REM sleep in 1953, and studies to elucidate consciousness and wakefulness in the 1930's and 1940's, are the driving forces in our recent understanding of sleep. This chapter outlines a general overview of sleep in order to understand the nature of sleep and its

Address for correspondence:

Dr. S. Chokroverty, MD, FRCP, FACP
Professor and Co-Chair of Neurology;
Chairman of Clinical Neurophysiology and Sleep Medicine; Program Director of Clinical Neurophysiology and Sleep Medicine; NJ Neuroscience Institute at JFK; Seton Hall University, Edison, NJ, USA

function, as well as physiological and anatomical substrates.

Definition of Sleep, Sleep Architecture and Sleep Stages

The modern sleep scientists define sleep on the basis of behavioral and physiological criteria. Behavioral criteria (Table 1) include lack of mobility or slight mobility, closed eyes, characteristic sleep posture, a markedly reduced response to external stimulation and a reversible unconscious state. Physiological criteria (Table 1) are based on the findings of EEG, electro-oculography (EOG) and electromyography (EMG). Based on physiological criteria, sleep is divided into two states with independent controls and functions: non-rapid eye movement

Table 1: Behavioral and physiological criteria of wakefulness and sleep

Criteria	Awake	Nonrapid eye movement sleep	Rapid eye movement sleep
Posture	Erect, sitting, or recumbent	Recumbent	Recumbent
Mobility	Normal	Slightly reduced or immobile; postural shifts	Moderately reduced or immobile; myoclonic jerks
Response to stimulation	Normal	Mildly to moderately reduced	Moderately reduced to no response
Level of alertness	Alert	Unconscious but reversible	Unconscious but reversible
Eyelids	Open	Closed	Closed
Eye movements	Waking eye movements	Slow rolling eye movements	Rapid eye movements
Electroencephalography	Alpha waves; desynchronized	Synchronized	Theta or saw tooth waves; desynchronized
Electromyography (muscle tone)	Normal	Mildly reduced	Moderately to severely reduced or absent
Electro-oculography	Waking eye movements	Slow rolling eye movements	Rapid eye movements

(NREM) and REM sleep (Table 2) alternating in a cyclical manner throughout the night, each cycle lasting approximately 90 to 110 minutes during the normal sleep in adults (four to six cycles are noted). The first third of normal sleep is dominated by slow wave sleep (Stages 3 and 4 of NREM sleep or SWS), and the last third is dominated by REM sleep. The first REM cycle is short, and the last REM cycle (toward the end of the night or early hours of the morning) is generally the longest and can last up to an hour.

Table 2: Sleep States and Stages

- NREM Sleep: 75 - 80 % of sleep time
 - Stage 1: 3 - 8 % of sleep time
 - Stage 2: 45 - 55% of sleep time
 - Stage 3: 3 - 8 % of sleep time
 - Stage 4: 20 - 25% of sleep time
- REM Sleep: 20 - 25% of sleep time

NREM sleep accounts for 75-80% of sleep time in adult humans and is subdivided into four stages based mainly on EEG criteria. Stage 1 NREM comprises 3-8% of sleep time; Stage 2 NREM occupies 45-55%, and Stage 3 and 4 make up 15-20% of total sleep time. In Stage 1 NREM sleep, the alpha rhythm (8-13 Hz), characteristic of wakefulness, diminishes to less than 50% in an epoch (typically a 30-second polysomnographic tracing) and a mixture of slower theta rhythms (4-7 Hz) and beta (greater than 13 Hz waves) appear (Fig. 1). Slow, rolling eye movements may be noticed at this time.

Fig 1: Polysomnographic recording shows stage I non-rapid eye movement (NREM) sleep. Electroencephalograms (top 10 channels) show a decrease of alpha activity to less than 50% and low-amplitude theta and beta activities. Electrooculographic channels show slow rolling eye movements. Rest of the montage is the same as in Figure 6.

Vertex sharp waves begin to appear towards the end of Stage 1 NREM sleep. Stage 2 NREM sleep begins normally about 10-12 minutes after Stage 1 NREM. Stage 2 NREM is characterized by sleep spindles (12-18 Hz, mostly 14 Hz) and K complexes intermixed with vertex sharp waves (Fig. 2). The background rhythm consists of a mixture of theta and delta waves (less than 4 Hz) comprising less than 20% of the epoch. During Stage 3 NREM, delta waves occupy 20-50% of the epoch (Fig. 3) and in Stage 4 of NREM, delta waves occupy more than 50% of the epoch (Fig. 4). Stage 3 and Stage 4 NREM sleep are grouped together as slow wave sleep.

Fig 2: Polysomnographic recording shows stage II non-rapid eye movement (NREM) sleep. Note 14-Hz sleep spindles in the electroencephalographic channels seen prominently in C3-A2 and C4-A1 intermixed with delta waves up to 2 Hz and 75 μ V in amplitude occupying less than 20% of the epoch. See Figure 6 for description of the rest of the montage.

Fig 3: Stage III non-rapid eye movement (NREM) sleep in this segment of polysomnogram. Electroencephalograms (top 10 channels) show 20% to 50% of the epoch occupied by delta waves as defined in Figure 2. See Figure 6 for description of the rest of the montage.

REM sleep begins 60-90 minutes after the onset of NREM sleep. EEG during REM sleep is characterized by fast rhythms and theta waves, some of which may have a characteristic sawtooth appearance (Fig. 5). Fundamental characteristics of REM sleep are the presence of rapid eye movements and markedly diminished or absent muscle activities. These traditional scoring criteria will soon be modified (a committee appointed by the American Academy of Sleep Medicine

Fig 4: Polysomnographic recording shows stage IV non-rapid eye movement (NREM) sleep. More than 50% of the epoch in the electroencephalograms (top 10 channels) consists of delta waves as defined in Figure 2. See Figure 6 for description of the rest of the montage.

Fig5: Polysomnographic recording shows rapid eye movement (REM) sleep. Electroencephalograms (top 10 channels) show mixed frequency theta, some alpha, and low-amplitude beta activities intermixed with sawtooth waves (middle of the recording) seen prominently in C3-A2 and C4-A1 derivations. Rapid eye movements are seen in electrooculographic channels. Chin electromyogram shows marked hypotonia, whereas Tib electromyogram shows phasic myoclonic bursts

is working to revise the original staging criteria), and NREM sleep will be divided into three stages. REM sleep will not be subdivided into tonic and phasic stages. In a normal adult, there is an orderly progression from wakefulness to sleep onset to NREM sleep and finally REM sleep. Relaxed wakefulness is characterized by a behavioral state of quietness and a physiologic state of alpha and beta frequencies on EEG (Fig. 6), waking eye movements and increased muscle tone. NREM sleep is characterized by progressively decreased responsiveness to external stimulation accompanied by slow eye movements, followed by EEG slow activity associated with sleep spindles and K complexes as well as decreased muscle tone. REM sleep state is characterized by further reduction of responses to stimulation, absent muscle tone and low amplitude fast EEG activity mixed with distinct sawtooth waves (sharp-contoured waves at 2-7 Hz).

Sleep macrostructure consists of sleep states and stages, sleep cycles, sleep latency, sleep efficiency (duration of total sleep time to total time in bed expressed in percent) and wake after sleep onset, whereas sleep microstructure includes momentary dynamic phenomena such as arousals, K complexes and sleep spindles and cyclic alternating pattern (CAP). The CAP indicates sleep instability, whereas frequent sleep arousals signify

Fig 6: Polysomnographic recording showing wakefulness. Top 10 channels show posterior dominant 10-Hz alpha activity intermixed with some low-amplitude beta rhythms in the electroencephalogram (International nomenclature). A2, right ear; A1, left ear. Waking eye movements are seen in the electrooculogram of the left (Lf) and right (Rt) eyes referred to linked ear. Chin electromyography (EMG) shows much tonic activity. Ononasal, oronasal airflow; chest, respiratory effort—chest; abdomen, respiratory effort—abdomen; snoring, this channel shows no snoring artifact; ECG, electrocardiogram; SaO₂, oxygen saturation by finger oximetry.

Table 3: Physiological changes during wakefulness, NREM sleep, and REM sleep

<i>Physiology</i>	<i>Wakefulness</i>	<i>NREM sleep</i>	<i>REM sleep</i>
Parasympathetic activity	++	+++	++++
Sympathetic activity	++	+	Decreases or variable (++)
Heart rate	Normal sinus rhythm	Bradycardia	Brady-tach yarrhythmia
Blood pressure	Normal	Decreases	Variable
Cardiac output	Normal	Decreases	Decreases further
Peripheral vascular resistance	Normal	Normal or decreases slightly	Decreases further
Respiratory rate	Normal	Decreases	Variable; apneas may occur
Alveolar ventilation	Normal	Decreases	Decreases further
Upper airway muscle tone	++	+	Decreases or
Upper airway resistance	++	+++	++++
Hypoxic and hypercapnic ventilatory responses	Normal	Decreases	Decreases further
Cerebral blood flow	++	++ or +++	++++
Thermoregulation	++	+	
Gastric acid secretion	Normal	Variable	Variable
Gastric motility	Normal	Decreases	Decreases
Swallowing	Normal	Decreases	Decreases
Salivary flow	Normal	Decreases	Decreases
Migrating motor complex (a special type of intestinal motor activity)	Normal	Slow velocity	Slow velocity
Penile or clitoral tumescence	Normal	Normal	Markedly increased

NREM = non-rapid eye movement; REM = rapid eye movement; + = mild; ++ = moderate; +++ = marked; ++++ = very marked; – = absent.

sleep fragmentation. The CAP is characterized by a repetitive EEG pattern in a cyclical manner noted mainly during NREM sleep. A CAP cycle consists of an unstable phase (Phase A) and a relatively stable phase (Phase B) each lasting between 2 and 60 seconds. Phase A of CAP is marked by an increase of EEG activity with contributions from both synchronous high amplitude slow and desynchronized fast waves in the EEG standing out from a relatively low amplitude slow background. Phase A is associated with an increase in heart rate, respiration, blood pressure and muscle tone. CAP rate and arousals both increase in older individuals and in a variety of sleep disorders. A non-CAP (i.e., a sleep period without CAP) is thought to indicate a state of sustained stability.

Ontogeny of Sleep

There are dramatic changes in the requirements of sleep from infancy to old age. Newborns have a polyphasic sleep pattern with 16 hours of sleep per day in the first few days of life. By 3-5 years of age, sleep requirements fall to about 10 hours per day. The polyphasic sleep patterns of newborns change to the biphasic pattern in preschool children. Adults exhibit a monophasic pattern with an average duration of 7 ½ to 8 hours per night; however, this reverts to a biphasic pattern in elderly persons. In the newborn, REM sleep occupies about 50%, but by 6 years of age this is decreased to the normal adult pattern of 25%. By three months of age, NREM/

REM cyclic pattern of sleep is noted and sleep spindles begin to appear generally by six to twelve weeks. K complexes are noted at about six months of age. Sleep in the elderly is characterized by an attenuation of the amplitude of delta waves, reduction of slow wave sleep stage and repeated awakenings, including early morning awakenings. The percentage of REM sleep in normal elders remains relatively constant, and the total duration of sleep time within 24 hours also is generally not different from that of young adults because older individuals often nap during the daytime, compensating for lost sleep during the night.

Phylogeny of Sleep

Sleep is present in both mammals and non-mammals, but sleep architecture and sleep stages differ. Both vertebrates and invertebrates display sleep/wakefulness and basic rest/activity cycles (BRAC). Mammalian sleep shows REM/NREM sleep stages and cycles, excepting dolphins, porpoises, and Australian Spiny Anteaters which do not have REM sleep; however, there is recent evidence indicating that the echidna (spiny anteaters) may have REM sleep. Another characteristic is that the dolphin's brain may show sleep in one half while wakeful activities are seen in the other half of the brain. This rhythm then reverses after 30 to 60 minutes in the two halves of the brain. Birds also have NREM/REM sleep cycles, but they are very short in duration. Sleep is also noted in reptiles and fish as well as invertebrates, such as insects, scorpions and worms, based on behavioral criteria.

Circadian Rhythm and Chronobiology of Sleep

The term circadian rhythm is derived from the Latin word *circa* which means about and *dies* which means day. The cycle lasts approximately 25 hours instead of 24 hours of day/night cycle. The existence of circadian rhythms was first observed in 1731 when the French Astronomer de Mairan noted that a heliotrope plant's leaves closed at sunset and opened at sunrise even when the plant was kept in darkness. This observation suggests that an internal clock controls the 24-hour rhythm. The site of this biological clock has been located in the paired suprachiasmatic nuclei of the hypothalamus above the optic chiasm. Time isolation experiments have shown that daily rhythms exist in many physiological processes

such as sleep, basic rest activity, body temperature and neuroendocrine secretions. The most important of these other rhythms is body temperature rhythm which, although independent of the circadian rhythm of sleep/wakefulness, follows it closely. Dysfunction of circadian rhythm results in some important human sleep disorders. The molecular basis of the mammalian circadian clock is beginning to be identified. A total of 8-9 genes (clock, par, bmal, etc.) and their protein products have been identified within the circadian clock system and they are still evolving. Remarkable progress has been made in the past few years in identifying key components of the circadian clock in both fruit flies (*Drosophila*) and mammals. The molecular mechanism of two human circadian rhythm disorders – advanced sleep phase syndrome (ASPS) and delayed sleep phase syndrome (DSPS) – has been uncovered by applying gene sequencing techniques. ASPS seem to be due to a mutation of the hper 2 gene (a human homologue of the period 2 in *Drosophila*) causing advancing of the clock (i.e., alteration of the circadian timing of sleep propensity). DSPS is due to polymorphism of hper 3 gene.

Regulation of sleep/wakefulness is modulated by homeostatic and circadian patterns. The homeostatic factor refers to the increased propensity for sleepiness with longer periods of prior wakefulness, whereas the circadian factor refers to variations in physiologic alertness and sleepiness (timing, duration, and other characteristics) that vary cyclically with time of day. There are two very highly vulnerable periods of sleepiness: 2-6 a.m., especially 3-5 a.m. and 2-6 p.m.; the former is stronger than the latter. The timing of the physiologic sleepiness (mid-afternoon and early morning hours) depends on circadian factors. The highest number of sleep-related accidents has been observed during this period. It has been recently suggested that the suprachiasmatic clock and the genes may also affect sleep regulation and sleep homeostasis independent of circadian rhythm generation. This is based on the findings of anatomical projections from the suprachiasmatic nucleus to lateral hypothalamic neurons containing hypocretin (wake-promoting) peptide and anterior hypothalamic ventrolateral preoptic (VLPO) containing sleep-promoting neurons, as well as on experimental evidence. Experimental studies in mice using knockout mice lacking core clock genes and those with mutant clock genes suggested that circadian clock genes may also affect sleep regulation independent of

circadian rhythm generation.

The role of various sleep factors in maintaining homeostasis has not been clearly established. Several cytokines such as interleukin-1, interferon alpha and tumor necrosis factor promote sleep. There are other sleep factors which increase in concentration during prolonged wakefulness and infection. It has been shown that adenosine can fulfill the major criteria for the neural sleep factor that mediates this somnogenic effect of prolonged wakefulness. There are several other endogenous compounds which may serve as sleep factors and include delta sleep-inducing peptides, muramyl peptides, cholecystokinins, arginine vasotocin, vasoactive intestinal peptide, growth hormone-releasing factors and somatostatin.

Physiological Changes in Sleep

A variety of physical changes occurs during NREM and REM sleep that are different from those noted during wakefulness. Profound changes occur in the autonomic nervous system, causing important physiologic alterations in the respiratory and cardiovascular functions (Table 3). Profound hemodynamic changes (unstable blood pressure and heart rate, and progressive decrease in cardiac output causing maximum oxygen desaturation and periodic breathing, and intermittent increase of sympathetic activity during REM sleep) may explain increasing mortality during the early morning hours, especially in patients with cardiopulmonary disease. These hemodynamic and sympathetic alterations may initiate increased platelet aggregations, plaque rupture and coronary arterial spasm which may trigger thrombotic events that cause myocardial infarction, ventricular arrhythmia, and even sudden cardiac death and stroke. Sleep-related alveolar hypoventilation and increased upper airway resistance may predispose susceptible individuals to upper airway occlusion and obstructive apnea. Patients with neuromuscular disorders, chronic obstructive pulmonary disease, and bronchial asthma may be affected adversely from such hypoventilation. Asthmatic attacks may be exacerbated at night as a result of bronchoconstriction during sleep. There are also important physiologic changes during sleep in the gastrointestinal system, endocrine, renal and sexual functions as well as in thermal regulation. Body temperature begins to fall at the onset of sleep and reaches its lowest point during the third sleep cycle. Thermal regulation is maintained during NREM sleep but is non-

existent in REM sleep.

Functional Neuroanatomy of Sleep

The neuroanatomic substrates for REM sleep, NREM sleep and wakefulness are located in separate parts of the central nervous system. There are no discrete sleep-promoting centers, but the sleep/wake stages are produced by changes of interconnected systems rather than by discrete centers. The methods to characterize sleep/wake generator sites include lesion, stimulation, ablation, intracellular (neuronal unit recording), C-fos immunoreactivity (C-fos or immediate early gene is a nuclear protein released during activation of neurons) and neuroimaging mapping of neuronal networks.

Neuroanatomical substrates of REM sleep

Transection experiments through different regions of the midbrain, pons and medulla clearly show the existence of REM sleep-generating neurons in the pontine cat. A transection at the junction of the pons and midbrain produced all the physiologic findings compatible with REM sleep in the section caudal to that transaction, whereas in the forebrain region, rostral to the section, the recording showed no signs of REM sleep. After a transection between the pons and medulla, structures rostral to the section showed signs of REM sleep, but, in structures caudal to the section, there were no signs of REM sleep. After a section at the junction of the spinal cord and medulla, REM sleep signs are noted in the rostral brain areas. Finally, transection at the pontomesencephalic and the pontomedullary junctions produced an isolated pons that showed all the signs of REM sleep. The pons is, therefore, sufficient and necessary to generate all the signs of REM sleep. Neurons in the pedunculopontine tegmental (PPT) and the laterodorsal tegmental (LDT) nuclei in the pontomesencephalic region are cholinergic and are the REM-on cells which are responsible for REM sleep and show high firing rates at this stage. The REM-off cells are located in the locus coeruleus and raphe nucleus; these cells are aminergic neurons and are inactive during REM sleep. Histaminergic neurons in the tuberomammillary region of the posterior hypothalamus can also be considered REM-off cells. Thus, the cholinergic REM-on and the aminergic REM-off cells are all located within transections in the pons. A reciprocal interaction between REM-on and REM-off

neurons in the brainstem is thought to be responsible for REM generation and maintenance. Cholinergic neurons of the PPT and LDT projecting to the thalamus and basal forebrain regions as well as to the medial pontine reticular formation are responsible for activation and generation of REM sleep. Aminergic cells seem to play a permissive role in the appearance of REM sleep state. Muscle hypotonia or atonia during REM sleep is thought to depend on inhibitory postsynaptic potentials generated by dorsal pontine interneurons sending descending axons. The pathway from the perilocus coeruleus alpha region ventral to the locus coeruleus to the lateral tegmental reticular tract and then to the medial medullary region (e.g., nucleus magnocellularis and paramedianus) and the reticulospinal tract, projecting to the anterior horn cells of the spinal cord, controls REM sleep-induced muscle atonia. An experimental lesion in the perilocus coeruleus alpha region as well as in the medial medullary region produced REM sleep without muscle atonia. In human REM behavior disorder, causing dream-enacting behavior associated with REM sleep without muscle atonia, a structural or functional alteration of the pathway maintaining muscle atonia during REM sleep is most likely responsible for such disorder.

Neuroanatomical substrates of wakefulness

The neuroanatomic substrates of wakefulness consist of the ascending reticular activating system (ARAS), cholinergic neurons, aminergic neurons and hypocretin systems. Projections from the ARAS terminating in the thalamus and thalamocortical projections to widespread areas of the cerebral cortex produce cerebral cortical activation during wakefulness. Extrathalamic projections from the brainstem reticular neurons terminate in the posterior hypothalamus and the basal forebrain region; the latter, in turn, project to the cerebral cortex to maintain wakefulness. The major neurotransmitter pathways that regulate the wakefulness systems (ARAS projecting to the thalamus, the hypothalamus and basal forebrain) utilize the cholinergic, noradrenergic dopaminergic and histaminergic neurons. Cholinergic neurons in the LDT and PPT nuclei in the pontomesencephalic region project to the thalamus, posterior hypothalamus and basal forebrain region. These cholinergic neurons fire at the highest rate during wakefulness and during REM sleep but decrease their

rates of firing at the onset of NREM sleep. Acetylcholine, through both nicotinic and muscarinic receptors, depolarizes and excites the thalamocortical projecting neurons, promoting tonic firing. Cholinergic receptor antagonists (e.g., scopoline) inhibit cortical activation. Wake-promoting aminergic neurons include noradrenergic neurons in the locus coeruleus, serotonergic neurons in the dorsal raphe of the brainstem, histaminergic neurons in the tuberomammillary nucleus of the hypothalamus and possibly also dopaminergic neurons in the ventral tegmental area, substantia nigra and the ventral periaqueductal area. The role of dopamine is uncertain, however, pharmacologic, biochemical, and physiologic studies suggest that dopamine probably through D_1 , and possibly through D_2 receptors along with noradrenergic system, promotes wakefulness. Norepinephrine-containing locus coeruleus neurons show their highest firing rates during wakefulness, their lowest during REM sleep and intermediate rate during NREM. Pharmacologic studies suggest that posterior hypothalamic histaminergic neurons also help maintain wakefulness. The excitatory amino acids glutamate and aspartate are intermingled within the ARAS and are present in many neurons projecting to the cerebral cortex, forebrain and brainstem. These excitatory amino acids are released maximally during wakefulness. Recent discovery of hypothalamic hypocretin neurons and their widespread CNS projections have directed attention to the role of hypocretin system mediating sleep/wake regulation. In 1998, deLecea and co-workers described two neuropeptides in the lateral hypothalamic and perifornical region which they named hypocretin 1 and hypocretin 2. Independently in the same year, Sakurai et al. described two neuropeptides in the same region which they named Orexin A (corresponding to hypocretin 1) and Orexin B (corresponding to hypocretin 2). Shortly after recognition, it was shown that these hypocretin systems have widespread ascending and descending projections to locus coeruleus, dorsal raphe, ventral tegmental area, tuberomammillary nuclei of the posterior hypothalamus, LDT/PPT, ventrolateral preoptic (VLPO) nucleus of the hypothalamus, basal forebrain, limbic system (hippocampus and amygdala), cerebral cortex, thalamus (intralaminar and midline nuclei) and autonomic neurons (nucleus tractus solitarius, dorsal vagal nuclei and intermediolateral neurons of the spinal cord). Hypocretin systems promote wakefulness mainly through excitation of histaminergic, noradrenergic and

serotonergic neurons. These systems also suppress REM sleep through activation of the aminergic neurons (REM-off) which in turn inhibit REM-generating neurons of the LDT/PPT (REM-on).

Neuroanatomic substrates of NREM sleep

During the great epidemic of encephalitis lethargica in the first quarter of the 20th century, von Economo noted that patients with encephalitis lethargica, who had excessive sleepiness, had pathological alterations in the posterior hypothalamus, whereas those with severe insomnia had predominant lesions in the anterior hypothalamus. These findings immediately suggested the existence of sleep/wake centers in the hypothalamus. Sleep-promoting neurons are thought to reside in the ventrolateral preoptic (VLPO) area of the anterior hypothalamus as well as in the region of the nucleus tractus solitarius in the medulla. VLPO neurons consist of two subgroups: "tightly clustered" and "diffuse" depending on their distribution pattern. The tightly clustered neurons project to the tuberomammillary nuclei and promote NREM sleep, whereas diffusely distributed neurons project to the aminergic nuclei in the locus coeruleus and dorsal raphe region of the brainstem, participating in REM sleep. The VLPO neurons fire actively during NREM sleep and their lesion induces insomnia. The GABA- and Galanin-containing VLPO neurons project to inhibit locus coeruleus, dorsal raphe and tuberomammillary aminergic nuclei which in turn inhibit VLPO neurons. The contemporary theory for the mechanism of NREM sleep suggests a reciprocal interaction between two antagonistic neurons in the VLPO (activation) of the anterior hypothalamus and wake-promoting (inhibition) neurons in the tuberomammillary nuclei of the posterior hypothalamus, basal forebrain and mesopontine tegmentum. Reciprocal interaction between sleep-promoting (activation) neurons in the region of the nucleus tractus solitarius and wake-promoting (inhibition) neurons within the ascending reticular activating system of the brainstem, independently of the reciprocal interaction of the neurons in the forebrain, also plays a role in the generation of NREM sleep. The latter is the old reticular passive hypothesis or the disfacilitation hypothesis of sleep. In this model of passive theory sleep results from a cascade of disfacilitation within the brainstem. The passive theory originated in two classic preparations in cats by Bremer: *cerveau isole* and *encéphale isole*. Bremer found that in

cerveau isole preparation (e.g., midcollicular transection) all specific sensory stimuli were withdrawn and the animals were somnolent, whereas in *encéphale isole* preparation (transection at C1 vertebral level disconnecting the entire brain from the spinal cord) these specific stimuli maintained activation of the brain and the animals were awake. Moruzzi and Magoun postulated that withdrawal of generalized activation from ARAS is responsible for somnolence in *cerveau isole* preparation, whereas activation of the midbrain reticular neurons causing direct excitation of the thalamocortical projections results in EEG desynchronization and behavioral arousal. These observations support the later suggestion of Steriade et al. that at the onset of NREM there is a deafferentation of the brain because of blockage of afferent information, first at the thalamic level, causing the waking "open" brain to be converted into a "closed" brain resulting from thalamocortical inhibition. This passive reticular theory was challenged by the experiments of Batini et al., producing the midpontine pretrigeminal section which is only a few millimeters below the section that produced *cerveau isole* preparation. In this midpontine preparation there were persistent EEG and behavioral signs of alertness, suggesting that structures located in the brainstem between *cerveau isole* and midpontine pretrigeminal preparations are responsible for wakefulness. The active inhibitory role of lower brainstem hypnogenic neurons in the region of the nucleus tractus solitarius on the upper brainstem ARAS was also clearly demonstrated by the pretrigeminal midpontine preparation of Batini et al. Thus both active and passive theories play a role in the generation of NREM sleep; however, the contemporary theory favors activation of VLPO neurons and inhibition of posterior hypothalamic neurons containing histamine and hypocretin. Reduced activity of the hypocretin projections to the locus coeruleus noradrenergic, midline raphe serotonergic, mesopontine dopaminergic and tuberomammillary hypothalamic histaminergic cells may also reduce the level of arousal causing sleepiness.

The role of the suprachiasmatic nuclei in the regulation of circadian as well as homeostatic factors determining initiation and maintenance of sleep has been discussed above. Finally, recent advances in neuroimaging studies, including positron emission tomographic (PET) and single photon emission computed tomographic (SPECT) scans, have been able to visualize dramatic changes in function in cortical and subcortical neuronal networks in different sleep states and stages,

advancing our understanding of the functional neuroanatomy of sleep/wakefulness. PET scans have shown marked activation of the amygdala and the anterior cingulate region (part of the limbic system) during REM sleep which is, of course, generated by the brainstem neurons. In contrast, in NREM sleep, neuroimaging techniques have shown declining function in the thalamocortical circuits including association cortex of the frontal, parietal and temporal lobes. Thus, the brainstem and hypothalamic and forebrain sleep/wake-promoting neurons modulate functions of widespread forebrain cortical areas, keeping in balance cortical and subcortical tone to control sleep/wake regulation.

Function of Sleep

Why do we sleep? The function of sleep remains the greatest biologic mystery of all times. There are several theories about the function of sleep (Table 4), but none are satisfactory. There have been some recent advances made in the molecular mechanism of memory consolidation during sleep. Studies have revealed that new memories can be strengthened by sleep. In other words, sleep can rescue memories that are lost during wakefulness during the daytime, and consolidated memories can be re-consolidated when they are re-activated. Memory is stored in two stages: short-term memory, which is labile, and long-term memory, which is stable and consolidated requiring synthesis of new RNA and proteins by the neurons. Recent studies have shown clearly a significant contribution of sleep and wakefulness in the development of memory consolidation. There is suggestion that memory consolidation occurs during both slow wave and REM sleep. There is a further suggestion that REM/NREM sleep cycling is also important for memory consolidation.

The theory of maintenance of synaptic and neuronal network integrity is an emerging concept that is concerned with the primary function of sleep. Intermittent stimulation of neural network synapses is

Table 4: Biological functions of sleep

Body and brain tissue restoration
Energy conservation
Adaptation
Memory reinforcement and consolidation
Synaptic neuronal network integrity
Thermoregulation

necessary to preserve CNS function. Despite all these theories as listed in Table 4, we are still searching for clues about the functions of sleep.

International Classification of Sleep Disorders

In the new revised International Classification of Sleep Disorders (ICSD-2), sleep disorders are classified into eight broad categories (Table 5). Each of these eight categories is subdivided into several disorders, and pertinent diagnostic points are listed. An approach to a patient with sleep complaints must begin with a comprehensive knowledge of the ICSD-2 so that the patient can be evaluated in a proper manner, paying particular attention to the history and physical findings before ordering laboratory tests.

Table 5: International Classification of Sleep Disorders (ICSD-2)

1. Insomnias
2. Sleep related breathing disorders
3. Hypersomnias of central origin not due to circadian rhythm sleep disorder, sleep related breathing disorder, or other cause of disturbed nocturnal sleep
4. Circadian rhythm sleep disorders
5. Parasomnias
6. Sleep related movement disorders
7. Isolated symptoms, apparently normal variants, and unresolved issues
8. Other sleep disorders

Appendix A. Sleep disorders associated with conditions classifiable elsewhere

Appendix B. Other psychiatric and behavioral disorders frequently encountered in the differential diagnosis of sleep disorders

References

1. **McCarley RW.** Sleep neurophysiology; basic mechanisms underlying control of wakefulness and sleep. In: Chokroverty S. (ed), *Sleep Disorders Medicine*, ed. 2. Boston: Butterworth-Heinemann, 1999: 21-50.
2. **Siegel JM.** REM sleep. In Kryger MH, Roth T, Dement WC (eds). *Principles and Practice of Sleep Medicine*, ed. 4. Philadelphia: Elsevier/Saunders, 2005: 120-135.

3. **Steriade M**, McCarly RW. Brainstem control of wakefulness and sleep. New York: Plenum Publishers, 2005.
4. **Sakai K**, Crochet S. A neural mechanism of sleep and wakefulness. *Sleep and Biological Rhythms* 2003; 1: 29-42.
5. **Nishino S**, Yoshida Y. History and perspectives of hypocretin/orexin research in sleep medicine. *Sleep and Biological Rhythms* 2003; 43-54.
6. **Scammell M**. The neurobiology, diagnosis, and treatment of narcolepsy. *Ann Neurol* 2003; 53: 154-166.
7. **Steriade M**. The Neurophysiology mechanisms of non-rapid eye movement (resting) sleep. In: Chokroverty S (ed), *Sleep Disorders Medicine*, ed. 2. Boston: Butterworth-Heinemann, 1999: 51-62.
8. **Nofzinger EA**. What can neuroimaging findings tell us about sleep disorders? *Sleep Medicine* (Vol. 5, Supplement 1) 2004: S16-S22.
9. **Chokroverty S**. An overview of normal sleep. In: Chokroverty S, Hening W, Walters A (eds), *Sleep and Movement Disorders*. Boston: Butterworth-Heinemann, 2003: 23-43.
10. **De Lecea L**, Kilduff TS, Peyron C, et al. The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc National Academy Sci USA*, 199; 98: 365
11. **Sakurai T**, Amemiya A, Ishii M, et al. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell* 1998; 92: 573
12. The International Classification of Sleep Disorders, Second Edition. *Diagnostic and Coding Manual*. American Academy of Sleep Medicine, Westchester, IL, USA, 2005.
13. **Chokroverty S**. Physiological Changes in Sleep. In: Chokroverty S. (ed), *Sleep Disorders Medicine*, ed. 2. Boston: Butterworth-Heinemann, 1999: 95-126.
14. **Terzano M**, Parrino L, Smerieri A, et al. Atlas, rules and recording techniques for scoring of cyclic alternation pattern (CAP) on human sleep. *Sleep Medicine* 2002; 3: 187