Erratum to

Kidney Histology of Mice After Seven Days Oral Intake of Cyanobacterial Extract

[*J. Braz. Soc. Ecotoxicol.* 2 (2007) 39-43]

E. G. CARVALHO,^{1*} R. B. SOTERO-SANTOS,¹ C. B. R. MARTINEZ,³ E. C. FREITAS,¹
N. FENERICH-VERANI,² M. J. DELLAMANO-OLIVEIRA¹ & O. ROCHA¹

¹Departamento de Ecologia e Biologia Evolutiva, Universidade Federal de São Carlos, São Carlos, SP, Brasil

²Departamento de Hidrobiologia, Universidade Federal de São Carlos, São Carlos, SP, Brasil

³Centro de Ciências Biológicas, Departamento de Ciências Fisiológicas, Campus
Universitário Perobal, Universidade Estadual de Londrina, Londrina, PR, Brasil

The publisher regrets that Figure 3 of this paper was missing. The missing Figure is reproduced below. The publisher apologizes for any inconvenience caused.

Figure 3 – Kidney histology after treatment with 0.5, 1.0 and 2.0 mg mL⁻¹ concentrations of crude cyanobacterial extract. Kidney section (7 μ m) showing glomeruli (G) and renal tubules (T) from control (a, b), 0.5 (c, d), 1.0 (e, f) and 2.0 mg mL⁻¹ (g, h) groups. 640X.

*Corresponding author: Elisa Garcia Carvalho, e-mail: elisa.garciacarvalho@gmail.com.