Original Research

DOI: 10.5152/cjms.2021.1470

THE SMOKING STATUS AND ATTITUDES OF MEDICAL SCHOOL STUDENTS IN A UNIVERSITY OF NICOSIA

Authors: Ozen Asut¹, Songul Vaizoglu¹, Gulpiya Abduxcur¹, Cali Sanda¹

Institutions:

¹Near East University Faculty of Medicine, Public Health, Nicosia (Lefkosa), Turkey

Corresponding Author: Ozen Asut

E-mail: ozen.asut@neu.edu.tr

Received Date: 04.11.2019

Accepted Date: 30.06.2020

How to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470

Abstract

Background and aims

The smoking prevalences of physicians and medical students continue to preserve their high levels around the world. This study is the 2018 survey of a broader intervention started in 2017, aiming to monitor the smoking status of medical students in a multi-national university of Northern Cyprus.

Methods

The universe of this cross-sectional study was all of the 1461 medical students registered at the medical school for the academic year 2018-2019. The data were collected by a self-administered

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

questionnaire applied under direct observation. Of the students, 1304 responded to the questionnaire with a response rate of 89.3%. The data were analyzed by IBM SPSS 18.0, with $p < 0.01$ evaluated as significant.

Results

The participants were from 31 counties, the majority being citizens of the European-Central Asian country group. The overall smoking frequency was 33.7%, with a frequency of 42.1% for males and 26.1% for females. While 26.8% consumed tobacco products other than cigarettes, 50.8% of smokers had starting smoking after registering to medical school. Sub-Saharan African country citizens smoked least by 6.0% and the European-Central Asian group smoked most by 35.7%. Non-smokers displayed significantly more positive attitudes for tobacco regulations and doctors being role models, compared to smokers.

Conclusion

The results of this survey revealed a high tobacco use frequency in spite of a medical curriculum including a specific tobacco program. Stronger educational and other behavioural interventions are needed to alter the smoking behaviours of medical students and thus, medical education should be modified accordingly.

Keywords: Smoking, Tobacco use frequency, Attitudes, Medical students, Northern Cyprus

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Abstract in Turkish

Giriş ve amaç

Tüm dünyada hekimlerde ve tıp öğrencilerinde sigara içme prevalansı yüksek düzeylerde seyretmektedir. Bu araştırma 2017'de başlatılan bir müdahalenin 2018 araştırmasını oluşturmaktadır. Amaç Kuzey Kıbrıs'taki çok uluslu bir üniversitede sigara içme durumunu izlemektir.

Yöntem

Bu kesitsel araştırmanın evreni, 2018-2019 akademik döneminde Tıp Fakültesi'nde kayıtlı 1461 tıp öğrencisidir. Veriler, gözlem altında uygulanan bir anketle toplanmıştır. Öğrencilerin 1304'ü ankete yanıt vermiş olup yanıt hızı %89,3'tür. Veriler IBM SPSS 18.0 programı ile değerlendirilmiş, $p < 0.01$ önemli kabul edilmiştir.

Bulgular

Katılımcılar 31 ülke yurttaşı olup çoğunluğu Avrupa-Orta Asya ülkeler grubundandır. Sigara içme sıklığı %33,7, erkeklerde %42.1, kadınlarda % 26.1'dir. Katılımcıların %26,8'i sigara dışı tütün ürünü kullanmaktadır. Sigara içenlerin %50,8'i sigaraya tıp fakültesinde başladığını belirtmiştir. Sahraaltı Afrika ülkeleri üyeleri %6 ile en az sigara tüketirken, Avrupa-Orta Asya grubu %35,7 ile en çok sigara içen ülke grubu üyeleridir. Sigara içmeyenler, içenlere göre, tütün kontrolü düzenlemeleri ve doktorların rol modeli olmaları konularında önemli ölçüde daha olumlu tutum içindedirler.

Sonuç

Bu araştırmanın sonuçları, tıp müfredatında uygulanan özgün tütün eğitime karşın, öğrencilerde yüksek tütün kullanım sıklığı olduğunu göstermektedir. Sigara içme davranışlarını değiştirmek için daha güçlü eğitim ve diğer davranışsal müdahaleler gerekmektedir ve tıp eğitimi bu doğrultuda değiştirilmelidir.

Keywords in Turkish

Sigara içme, Tütün kullanım sıklığı, Tutum, Tıp öğrencisi, Kuzey Kıbrıs

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Introduction

Tobacco use is a major public health problem globally and specifically in developing countries, resulting in more than seven million deaths per year according to the World Health Organisation (WHO). More than one billion people of our planet are smokers (1).

Health professionals and particularly doctors are in a position to curb health problems related to tobacco use by community approaches. Attitudes and behaviours of doctors and their role model functions are influential on increasing the willingness of their patients to quit smoking, as well as in affecting other lifestyle behaviours (2-5). A number of studies conducted during the last decades of the previous century and up to the present time have revealed the efficacy of physician's role in assisting individuals for smoking cessation, and doctors with good lifestyle behaviours have a better chance for this end (3-5). Yet, smoking among doctors and medical students continues to stay high in many countries, in spite of the WHO and World Medical Association efforts to assist medical doctors globally and to guide national medical associations for effective measures (2).

The tobacco use attitudes and behaviours of doctors and medical students are well documented by ample research worldwide (5-10). For example, a study of 2008-2012 from Brazil, found the level of agreement of the students with the WHO recommendations as high. The prevalence of cigarette smoking was 5.2%, but 43.8% of the students reported experimenting water-pipe (11), whereas a study among medical students of Saudi Arabia reported a 13% smoking frequency among male students (12). According to an international review of 2007 on tobacco smoking among medical students, the prevalence ranged from 3% in the United States to 58% in Japan (13). In general, the attitudes and behaviours of medical students in this respect closely reflect the community they are a part of, just as the medical doctors and other health professionals.

Turkey is one of the countries with a considerable number of studies on the tobacco issue conducted during recent decades. On the average, the frequencies of smoking among medical students were found in the range of 30-40% in Turkey during late 1990's and early 2000's (14-16).

Smoking frequency is consistently higher for male medical students than females and the smoking rates increase with the advancing years of medical education (14-17) in Turkey. In one study, smoking frequency was 34.5% in the first year rising up to 44.2% in the 6th year of medical school (17).

Comprehensive tobacco control education in the medical curriculum is crucial for preventing the initiation of the smoking habit or lowering the prevalence of smoking among medical students and assisting the students to identify themselves as non-smoking role models for the community (5,13).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abdurcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

This study is part of an intervention study designed to follow up the students in a medical school of the Turkish Republic of Northern Cyprus (TRNC). The aim is to monitor the students as to the changes in trends of smoking attitudes and behaviours in relation to the impact of educational interventions.

Objectives: The objective of this study is to determine the tobacco use prevalence and attitudes regarding tobacco use among medical students in a university of Northern Cyprus for the year 2018.

The broader and long term objectives of this study include monitoring tobacco use prevalence among medical students of the Near East University between 2017-2021 and to follow up the changes in smoking behaviours of the students after integrating comprehensive tobacco education into the NEU medical curriculum.

Material and methods

The university where the study was conducted is located in Nicosia, Northern Cyprus and accepts citizens of countries from various regions of the world. The current study was designed as a cross-sectional survey with the aim of accessing all the 1461 medical students registered at the Turkish and English programs of the Medical School in December 2018.

The data were collected by a questionnaire of 12 questions developed by the researchers, and applied under direct observation. The questionnaire included 4 questions on socio-demographic features and 8 questions on smoking behaviours and attitudes, which were developed in compliance with WHO adult questionnaires for Global Adult Tobacco Survey (GATS) (18, 19, 20). The questions were revised so that the best input from the participants would be attained. Besides, some adaptations were performed relevant to the local conditions, ie, the grouping of number of cigarettes per day. The questionnaire has been attached as Annex 1.

Statistical Analysis:

The data collected were analysed by IBM SPSS 18.0 program. Chi square and Fisher's exact tests were used for determining the differences of the groups and $p < 0.01$ was evaluated as significant

Ethics approval and consent to participate:

The informed consent of the participants was requested and anonymity of the responses was guaranteed. Permission from the dean of the Faculty of Medicine and the approval of the Ethics Committee university were provided (Report No.XXX/2018/63-673, dated 22 November 2018).

Terms and definitions: (20, 21, 22)

The following terms and definitions based on WHO guidelines were used in accordance with the context of the study:

Current smoker: Percentage of adults who currently smoke tobacco.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Daily smoker: Percentage of adults who currently smoke tobacco daily.

Occasional smoker: Percentage of adults who currently smoke tobacco less than daily.

Former smoker: Percentage of adults who are ever daily and occasionally tobacco smokers and currently do not smoke tobacco.

Cigarette smoker: Percentage of adults who currently smoke cigarettes.

Current tobacco user: Percentage of adults who currently use tobacco (current daily and less than daily tobacco smokers and/or smokeless tobacco users).

Results

The total number of students studying at the Turkish and English programs of the Medical School was 1461 in December 2018. The number of students who responded to the questionnaire was 1304, comprising a response rate of 89.3%. The majority of the Turkish medical program students (97.5%) are citizens of Turkey and TRNC. Most of the English medical program students (75.6%) are from five countries: Turkey, Nigeria, Syria, Jordan and TRNC, in descending order.

Table 1 shows some socio-demographic characteristics and the cigarette and non-cigarette tobacco product use features of the participants.

Table 1.

Of the participants, 57.9% are lifetime non-smokers and 33.7% are current smokers, while 76.4% of the smokers are daily smokers and 12.4% smoke more than 20 per day. Of the smokers and former smokers together, 54.8% reported a smoking duration of 2-5 years. Notably, 50.8% of the smokers stated they started smoking after admittance to medical school.

Consumers of non-cigarette tobacco products comprise 26.8% of the total participants. Of the non-smokers 17.7% and of the smokers 44.2% consume non-cigarette tobacco products. Cigarette smokers consume non-cigarette tobacco products at significantly higher levels than non-smokers

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

($\chi^2=86.0$, $p<0.001$). Of the male students 38.6% and of the females, 16.2% consume non-cigarette tobacco products; the difference between the genders being significant as well.

On the other hand, 80.7% of the students indicated being exposed to some form of second-hand smoke. Notably, 39.5% of the passive smokers were exposed at cafes and restaurants. Other locations of exposure included homes, polluted either by smoker household members (22.9%) or by visitors (19.3%) and indoor places other than home (34.7%) (Not shown in tables).

Table 2 shows the smoking status of the participants in relation to some socio-demographic features and their countries of origin classified as World Bank country groups by region and by income.

Table 2.

Male students smoke more than females at significantly higher rates ($\chi^2=36.9$, $p<0.001$); 42.1% of the male students and 26.1% of the females smoke. Third grade students smoke most with 37.4% followed by 5th and 6th grade students with 35.5% each.

Sub-Saharan country citizens smoke least by 6.0% while European and Central Asian region members smoke most with 35.7%. Smoking is highest among citizens of higher-middle income countries with 37.4%. Smoking frequency among lower-middle income country citizens (12.4%) is significantly lower than citizens of high income, higher-middle income and low income countries ($p<0.001$) (Table 2).

The comparison of the attitudes of smoker and non-smoker medical students regarding their attitudes on support of indoor smoking bans and their opinions on doctors being non-smoking role models is presented in Table 3.

Table 3.

Of the non-smokers, 91.9%, as compared to 73.9% of smokers are in support of smoking bans indoors. Non-smokers support smoking bans at significantly higher levels than smokers ($\chi^2=76.9$, $p<0.001$). Besides, non-smokers adopt the role model function of doctors at significantly higher levels than smokers at 87.7% for non-smokers and 54.3% for smokers.

Smoking status of all students according to their nationalities is indicated in Table 4, country by country.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Table 4.

Overall, citizens of Turkey, comprising 63% of the participants smoke at a higher frequency by 37.5% than the total average of 33.7%.

The analyses of medical students of various country income groups in regard to use of non-cigarette tobacco products revealed that low income countries of sub-Saharan Africa consume these products at the lowest level with 7.2% (not shown Tables). The consumption of these products in all other regions is above 25%. Surprisingly, consumption is highest among students of low economy countries with 31.3%, while 23.1% for high income countries and 28.9% for upper middle income countries. The students from lower middle income countries have the lowest frequency in this respect with 12.5%, which is significantly lower than other country groups.

Similar to the cigarette smoking data, 74.8% of non-cigarette tobacco users compared to 89.2 of non-consumers support the ban in indoor public places, the difference being significant ($\chi^2= 34.0$, $p< 0.001$). Similar results are valid for the opinions of the participants about the role model function of the physician.

Discussion

The status of tobacco use among medical students and their attitudes on the tobacco issue were studied in this cross-sectional study. The smoking frequency was 33.7% in general, 37.5% among Turkish citizens and 27.7% among Northern Cyprus citizens. The smoking prevalences are higher in our study than most medical student studies in Turkey (16, 17, 23). However, the smoking frequency was much higher (39%) in a study among students of a private medical school in Turkey (24).

Recent studies among university students and particularly medical students in the Middle East and African countries have found lower prevalences compared to the results of our study in general. However, two studies in Nigeria found higher frequencies compared to our Nigerian group; lifetime tobacco use was 9.6% (25) and 10.5% (26) in these surveys. The Nigerian medical students in our study showed a lower frequency as 3.1% .

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

A 2007 study in Syria on medical students found the cigarette smoking prevalence to be 10.9% (27), much lower than our general result of 33.7% and the Syrian students in particular, who had a smoker rate of 31.3% in our study.

A study in Jordan among medical students found the smoking prevalence as 26% for male and 7% for female students (28). On the contrary, the findings of the Jordanian students in our study showed a much higher frequency as 43.6%. Syrian and Jordanian results in the current study may be a reflection of the social status of our students, as well as the pro-smoking atmosphere of Cyprus grounds.

According to an international review, the prevalence of smoking among medical students ranged from 3% in the United States to 58% in Japan, with 29.5% for Italy, 29% for Saudi Arabia and 10-10.5% for Nigeria (13). Medical students are subject to similar impacts of their social environment and present behaviours as other members of their society, in spite of the fact that they are studying medicine. Thus, the higher smoking rates displayed by the present study may be attributed to the high smoking atmosphere of Northern Cyprus.

Smoking frequency was significantly higher among male students than female students in our study as expected. Furthermore, smokers were higher for both sexes among Turkish medical program students with 43.9% for men and 29.1% for women. These results are similar to but higher than the results of some medical schools of Turkey, which show a range of 8.2-17.3% for female students and 25.1-35.0% for males (23, 24).

The finding that male students smoke more than female students is consistent with other studies' findings(29). The study among medical students in Jordan revealed the prevalences as 26% for male and 7% among female students (28).

Contrary to other research, a study from China found that medical students have higher rates of smoking than other students but most of the smokers were occasional smokers(30). On the other hand, a study in Greece found smoking rates of medical students as 35.3%, while 50.2% for other students (31).

Smoking frequency is 28.9% among first year students and 35.5% for sixth grade students in the current study, while it is highest in the 3rd year with 37.4%. Smoking prevalences range from 8 to 18% among first year medical students and 21 to 39% among 6th year students according to various studies conducted in Turkey (16,32).

Our results show higher frequencies for almost all parameters. In a study comparing the students of seven medical schools and the same grades of other schools of science in Turkey, a significant

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

difference between the medical and other students was found in the first grades, with lower frequencies for medical students. However, the difference was not statistically significant for last grade students, although smoking rates were lower for medical students (32).

Of the participants in our study, 26.8% have stated using non-cigarette tobacco products, of which water pipe (narghile) is the leading product with 23.5%. According to a 2011 study among medical students in Turkey, 17% of the first year students and 14% of the 6th year students were using water pipe (33).

Exposure to various forms of passive tobacco smoke was stated by %80.7 of the participants in our study. Although the law regulating tobacco use has been enforced since 2008 in Northern Cyprus, these results display the non-compliance regarding indoor bans. On the other hand, the students are highly in favor of the indoor bans regardless of their smoking status. Still the rate of supporters is significantly higher among non-smokers.

The majority of the students are in the opinion that doctors should be non-smoking role models for the community. Non-smokers share this view at significantly higher levels than smokers (87.7% of non-smokers versus 54.3% of smokers). The 1994 study from Turkey had displayed a much lower frequency for this parameter; 44% of the smokers and 70% of the never smokers were in support of the role model function of doctors at that time (32). The difference between the two studies may be evaluated as an indicator of progress of the community and particularly of the medical students on the tobacco control issue during the years in between. Interestingly, 26.1% of smoker students and 8.1% of non-smokers do not support legal measures for tobacco control in the current study, the difference of the groups being highly significant. In comparison, these rates were 24% and 6% respectively in the seven university study of 1994 (32). These results reveal the role of smoking on attitudes of doctors and the importance of non-smoking behaviours.

More students have started smoking after registering to the medical school (50.8%) than those who initiated smoking before their medical education period (%49.2%). These findings indicate the importance of early tobacco control education in the medical school for prevention of picking up the habit. The findings of the present study are similar to the results of the first baseline study in our university (34) and indicative of inadequate tobacco control interventions, including current content of the medical education.

The role of medical education on the smoking trends of doctors has been a subject of research in recent years. According to global surveys, the tobacco education in medical curricula varies widely among countries and medical schools (13, 35).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

An international study among medical students of 42 countries in Europe, Asia, and Africa found significant deficiencies in medical education on tobacco and recommended urgent changes in the medical curricula (35). The content, program and methods of the tobacco education curriculum continue to be topics of discussion in the context of medical education.

The restrictions of the study

The study was conducted in one medical school of Northern Cyprus. Thus, the results of our study cannot be representative of all the medical schools of Cyprus, nor can it be associated with students studying in other schools of universities.

The questions of the study questionnaire were limited as for a descriptive study to obtain baseline data with the intention of accessing more students and more responses. Therefore, in depth and detailed analyses of the data could not be performed in this study, nor could we derive a wider variety of results from the data obtained.

Conclusion

This study about the smoking status of medical students revealed high smoking frequencies for both male and female students, compared to medical students of other countries, probably related to the high smoking atmosphere of Northern Cyprus. More than half of the smokers started smoking after entering the medical school and smoking status increased with advancing years of medical education, all showing the necessity of more tobacco control measures in the university and more tobacco education throughout the medical curriculum.

In addition to increasing education on tobacco, it is crucial to fully implement all tobacco control measures in medical schools. University campuses should be smoke-free and the rules and regulations regarding tobacco use should be strictly supported and enforced. Related educational and other activities about tobacco control should be available for all medical students throughout all the years and content of medical education.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Medical education should be re-structured to include the skills for tobacco cessation and other relevant subjects. Tobacco control and cessation techniques education in the curriculum of medical education is one of the intervention methods, which is in the agenda of our medical faculty in the near future.

List of abbreviations

TRNC: Turkish Republic of Northern Cyprus

WB: World Bank

WHO: World Health Organisation

Declarations

Consent for publication

Not applicable.

Availability of data and material

The datasets analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

No funding information to be declared.

Authors' contributions

All authors contributed to the development of the questionnaire, designing of the study, data collection and analyses. OA wrote the final manuscript. All authors read and approved the final manuscript.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

References

1. Tobacco Breaks Hearts. World Health Organisation; 2018. (<https://apps.who.int/iris/bitstream/handle/10665/272675/WHO-NMH-PND-18.4-eng.pdf?ua=1&ua=1>, accessed 09 March 2019).
2. Tobacco. Health Effects of Tobacco Secondhand Smoke. World Medical Association; 2019 (<https://www.wma.net/what-we-do/education/tobacco/> , accessed 10 March 2019).
3. Canadian Medical Association. The role of physicians in prevention and health promotion. CMAJ. 1995; 153: 208 A-D. (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1338069/pdf/cmaj00074-0083>, accessed 15 December 2018).
4. Frank E, Segura C, Shen H, Oberg E. Predictors of Canadian physicians' prevention counseling practices. Can J Publ Health. 2010; 101:390-5. PMID:21214054
5. Pipe A, Sorensen M, Reid R. Physician smoking status, attitudes toward smoking, and cessation advice to patients: an international survey. Patient Educ Couns. 2009;74:118-23. PMID:18774670 DOI:10.1016/j.pec.2008.07.042
6. Melani A, Verponziani W, Boccoli E, Trianni GL, Federici A, Amerini R et al. Tobacco smoking habits, attitudes and beliefs among nurse and medical students in Tuscany. Eur J Epidemiol. 2000; 16(7): 607-611. (<https://link.springer.com/article/10.1023%2FA%3A1007697126844?LI=true>, accessed 14 October 2018).
7. Omair A, Kazmi T, Alam SE. Smoking prevalence and awareness about tobacco related diseases among medical students of Ziauddin Medical University. J Pak Med Assoc. 2002 Oct; 52: 389. (https://www.researchgate.net/publication/10945862_Smoking_prevalence_and_a_wareness_about_tobacco_related_diseases_among_medical_students_of_Ziauddin_Medical_University, accessed 15 October 2018).
8. Maira JP and Akshava KM. Association between the duration of stay in medical colleges and the smoking behaviour among medical students. Indian J Community

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Med. 2017 Oct-Dec; 42 (4): 193–6.

(<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5682714/>, accessed 11 January 2019).

9. Vakeflliu Y, Argjiri D, Peposhi I, Agron S, Melani AS. Tobacco smoking habits, beliefs, and attitudes among medical students in Tirana, Albania. *Prev Med.* 2002Mar; 34(3): 370-3. (<http://www.sciencedirect.com/science/article/pii/S0091743501909946>, accessed 15 October 2018).
10. La Torre G, Kirch W, Bes-Rastrollo M, Ramos RM, Czaplicki M, Gualano MR et al. Tobacco use among medical students in Europe: Results of a multicentre study using the Global Health Professions Student Survey. *Public Health.* Feb 2012; 126(2): 159-64. (<https://iris.unipa.it/retrieve/handle/10447/99320/133097/tobacco%20%281%29.pdf>, accessed 15 October 2018).
11. Martins SR, Paceli RB, Bussacos MA, Fernandes FLA, Prado GF, Lombardi EMS et al. *J Bras Pneumol.* 2017;43(3): 202-7. Effective tobacco control measures: agreement among medical students. (<http://dx.doi.org/10.1590/s1806-37562015000000316>)
12. Al-Turki YA. Smoking habits among medical students in Central Saudi Arabia. *Saudi Med J.* 2006; 27(5):700-3. PMID: 16680263
13. Smith DR, Leggat PA. An international review of tobacco smoking among medical students. *J Postgrad Med.* 2007 Jan1; 53 (1): 55. (<http://www.jpgmonline.com/text.asp?2007/53/1/55/3033>, accessed 11 Apr 2019).
14. Akvardar Y, Demiral Y, Ergor G, Ergor A. Substance use among medical students and physicians in a medical school in Turkey. *Soc Psychiatry Psychiatr Epidemiol.* 2004; 39(6): 502-6. PMID: 15205736
15. Oguş C, Özdemir T, Kara A, Senol Y, Cilli A. Smoking habits of grade I and VI medical students at Akdeniz University Medical Faculty. *Türkiye Klinikleri Arch Lung* 2004; 5 (3): 139-42. (<http://www.turkiyeklinikleri.com/article/en-akdeniz-universitesi-tip-fakultesi-donem-i-ve-vi-ogrencilerinin-sigara-icme-aliskanliklari-54904.html>, accessed 11 April 2019). (In Turkish)
16. Vatan I, Ocakoglu H, Irgil E. Determining smoking prevalence in Uludağ University Faculty of Medicine. *TAF Prev Med Bull.* 2009; 8 (1): 43-8.
17. Kocabas A. Smoking prevalence and features of tobacco use in Turkey. *Solunum Hastalıkları.* 1994; 5: 133-47. (In Turkish)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. *Cyprus J Med Sci* 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

18. WHO Tobacco Questionnaire. World Health Organisation.
(https://www.who.int/tobacco/surveillance/6_GATS_CoreQuestionnairewithOptionalQuestions_v2.1_FINAL_13June2014.pdf?ua=1, accessed May 24, 2019).
19. Global Adult Tobacco Survey Collaborative Groups. Tobacco questions for survey: A subset of key questions from the Global Adult Tobacco Survey (GATS). 2nd Edition. Atlanta GA: Centers for Disease Control and Prevention; 2011.
20. WHO Report On The Global Tobacco Epidemic, 2009. Implementing smoke-free environments. World Health Organisation.
(https://afro.who.int/sites/default/files/2017-06/9789241563918_eng_full.pdf, accessed December 2018).
21. Global Adult Tobacco Survey (GATS) Fact Sheet Template Version 2.1 June 2012 .
(<https://www.paho.org/hq/dmdocuments/2017/18-GATS-AnalysisPackage-12June2014.pdf>, accessed May 27 2019).
22. Global Adult Tobacco Survey (GATS) Indicator Guidelines: Definition and Syntax Final Edition July 2009.
(https://www.who.int/tobacco/surveillance/en_tfi_gats_indicator_guidelines.pdf, accessed May 27 2019).
23. Oktay G, Hodoglugil NS, Agca E, Akdogdu E, Ardagil A, Celik E. Smoking behaviour and attitudes of 6th year medical students of Hacettepe University. In: International Public Health Congress Health 21 in Action October 2000; Istanbul, Turkey. p. 149.
24. Gunes G, Pehlivan E, Egri M, Genc M. The frequency of smoking among physicians, nurses and medical students of Turgut Ozal Health Center [Turgut Ozal Tip Merkezi hekim, hemşire ve tıp öğrencilerinde sigara içme sıklığı]. J Turgut Ozal Med Cent. 1997; 4(4):407-11. (In Turkish)
25. Makanjuola AB, Daramola TO, Obembe AO. Psychoactive substance use among medical students in a Nigerian university. World Psychiatry. 2007 Jun; 6(2):112. PMID: 18235869
26. Dania MG, Ozoh OB, Bandele EO. Smoking habits, awareness of risks, and attitude towards tobacco control policies among medical students in Lagos, Nigeria. Ann Afr Med. 2015 Jan 1; 14(1):1. doi: 10.4103/1596-3519.148701.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

27. Almerie MQ, Matar HE, Salam M, Morad A, Abdulaal M, Koudsi A et al. Cigarettes and water pipe smoking among medical students in Syria: a cross-sectional study. *IJTL*. 2008 Sep 1; 12(9):1085-91. PMID: 18713509
28. Merrill RM, Madanat HN, Cox E, Merrill JM. Perceived effectiveness of counseling patients about smoking among medical students in Amman, Jordan. *East Mediterr Health J*. 2009 Sep-Oct; 15(5):1180-91. PMID: 20214132
29. Hitchman SC, Fong GT. Gender empowerment and female-to-male smoking prevalence ratios. *Bull World Health Organ* 2011 Mar;89(3): 195-202. DOI:10.2471/BLT.10.079905
30. Zhu T, Feng B, Wong S, Choi W, Zhu SH. A comparison of smoking behaviours among medical and other college students in China. *Health Promot Int*. 2004 Jun 1; 19(2):189-96. PMID:15128710
31. Alexopoulos EC, Jelastopulu E, Aronis K, Dougenis D. Cigarette smoking among university students in Greece: a comparison between medical and other students. *Environ Health Prev Med*. 2010 Mar; 15(2):115-20. PMID: 19802656
32. Kocabas A, Burgut R, Bozdemir N, Akkoclu A, Cildag O, Dagli E et al. Smoking patterns at different medical schools in Turkey. *Tob Control* 1994; 3: 228-35. (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1759350/> accessed 10 March 2019).
33. Emlek G, Keklikoglu I, Demir N, Pekcan H, Erbaydar PN. The opinions on smoking and tobacco use experiences of the first and sixth grade students of Hacettepe University Medical School (Hacettepe Universitesi Tip Fakültesi birinci ve altinci sınıf öğrencilerinin tutun ürünlerini kullanma deneyimleri ve sigara içmeye yönelik bazı düşünceleri.) In: *Besinci Sigara ve Sağlık Ulusal Kongre Kitabı*, 8-11 Mayıs 2011, Kocaeli. (Proceedings of the 5th Tobacco and Health Congress, 8-11 May, Kocaeli.) 2011. p. 114. (In Turkish)
34. Asut O, Cali S, Ozcan A, Dundar F, Palaz C, Duran A et al. Tobacco use status among the students of the medical faculty Turkish medical program at a university in Nicosia. *STED* 2019; 28(1):17-27. DOI: 10.17942/sted.415640. (In Turkish)
35. Robyn L, Richmond RL, Debono DS, Larcos D, Kehoe L. Worldwide survey of education on tobacco in medical schools. *Tob Control* 1998; 7(3): 247-52. DOI: 10.1136/tc.7.3.247. (<https://tobaccocontrol.bmj.com/content/7/3/247>).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. *Cyprus J Med Sci* 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

TABLES

Table 1. Some socio-demographic characteristics and smoking features of the NEU medical students (Nicosia, December 2018) (N=1304)

Socio-demographic characteristics	n	%
Age group (years) (n=1292)		
≤ 17	24	1.9
18 – 24	1180	91.3
≥ 25	88	6.8
	<i>Mean± SD =21.3± 2.3</i>	<i>Minimum=16 Maximum=39</i>
Sex (n=1304)		
Male	623	47.8
Female	681	52.2
Lifetime smoking (n=1304)		
Never	541	41.5
Only tried	214	16.4
Former smoker	109	8.4
At least 1cigarette daily	337	25.8
Less than 1 cigarette daily	103	7.9
Use of non-cigarette tobacco products (n=1060)		
Never	776	73.2
Water pipe (Narghile)	249	23.5
Electronic cigarettes	15	1.4
Cigar	11	1.0
Pipe	9	0.9
Male students (n=503)	194	38.6
Total	284	26.8
Current cigarette smoking status (n=1304)		
Non-smoker	865	66.3

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Smoker	439	33.7
Number of daily cigarettes consumed by smoker students (n=435)		
Occasional (less than 1 daily)	70	16.1
1-5	88	20.2
6-10	95	21.9
11-20	128	29.4
Over 20	54	12.4
Duration of smoking (n=578) *		
One year or less	136	23.5
2-5 years	317	54.8
6-10 years	109	18.9
More than 10 years	16	2.8
Initiation of smoking in relation to starting medical school (n =478)		
Before	235	49.2
After	243	50.8
*Includes ex-smokers as well		

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Table 2. Smoking status of NEU medical students in relation to some socio-demographic features and to their countries of origin classified as WB country groups by region and by income groups ¹ (Nicosia, December 2018) (N=1304)

Smoking status								
Socio-demographic feature	Non-smoker		Smoker		Total		χ^2	P
	n	%	n	%	n	%		
Sex (n=1304)							36.9	<0.001
Male	361	57.9	262	42.1	623	100		
Female	503	73.9	178	26.1	681	100		
Age group (n=1292)							1.1	0.292
<25	805	66.9	399	33.1	1204	100		
≥25	54	61.4	34	38.6	88	100		
Grade of medical school (n=1304)								
1. grade	182	71.1	74	28.9	256	100		
2. grade	176	68.0	83	32.0	259	100		
3. grade	149	62.6	89	37.4	238	100		
4. grade	206	65.0	111	35.0	317	100		
5. grade	91	64.5	50	35.5	141	100		
6. grade	60	64.5	33	35.5	93	100		
Country group (region) (n=1290)								

¹ World Bank Country and Lending Groups (2019) (online).

<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519> Accessed 13 March 2019.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Middle East&North Africa	124	66.0	64	34.0	188	100
Europe&Central Asia	642	64.3	357	35.7	999	100
Latin America&Caribbeans	-	-	1	100	1	100
Sub-Saharan Africa	78	94.0	5	6.0	83	100
South Asia	13	76.5	4	23.5	17	100
North America	1	50.0	1	50.0	2	100
WB income group (n= 1282)						31.7 <0.001
High income	135	70.7	56	29.3	191	100
Higher middle income	564	62.6	337	37.4	901	100
Lower middle income	99	87.6	14	12.4	113	100
Low income	56	72.7	21	27.3	77	100

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Table 3. The comparison of the attitudes of smoking and non-smoking NEU medical students regarding support of indoor smoking bans and doctors being non-smoking role models (Nicosia, December 2018) (N=1304).

Attitudes regarding indoor smoking bans (n=1290)								
	Supporting		Not supporting		Total		χ^2	<i>P</i>
	n	%	n	%	n	%		
Smoking status							76.9	<0.001
Non-smoker	784	91.9	69	8.1	853	100		
Smoker	323	73.9	114	26.1	437	100		
Total	1107	85.8	183	14.2	129	100		
Attitude on doctors being non-smoking role models (n=1291)								
							179.5	<0,001
Non-smoker	751	87.7	105	12.3	856	100		
Smoker	236	54.3	199	45.7	435	100		
Total	987	76.5	304	23.5	129	100		

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

Table 4. Smoking status according to country of origin of NEU medical students ((Nicosia, December 2018) (N=1304)

Smoking status (n=1293)		Non-smoker		Smoker	Total
% *	n	% *	n	%	
62.5	305	37.5	814	63.0	
72.5	48	27.7	173	13.4	
96.9	2	3.1	64	4.9	
68.8	20	31.3	64	4.9	
56.4	24	43.6	55	4.3	
75.0	5	25	20	1.5	
75.0	3	25	12	0.9	
54.5	5	45.5	11	0.9	
77.8	2	22.2	9	0.7	
50	4	50	8	0.6	
85.7	1	14.3	7	0.5	
71.4	2	28.6	7	0.5	
66.7	2	33.3	6	0.5	
80	1	20	5	0.4	
100	-	-	4	0.3	
75	1	25	4	0.3	
50	4	50	4	0.3	
100	-	-	3	0.2	
100	-	-	3	0.2	
33.3	2	66.7	3	0.2	

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

50	1	50	2	0.2
50	1	50	2	0.2
50	1	50	2	0.2
50	1	50	2	0.2
100	-	-	2	0.2
100	-	-	2	0.2
80	1	20	5	0.4

*Row percentages

**Algeria, Armenia, Bulgaria, Romania, Venezuela

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

TOBACCO USE ATTITUDES AND BEHAVIOURS OF MEDICAL STUDENTS

QUESTIONNAIRE FOR DATA COLLECTION

Dear student,

This study has been planned to determine your attitudes and behaviours on tobacco use. The data of the study shall not be used for any purpose other than the one stated above. Your participation and appropriate and sincere responses to the questions will contribute to the access of right and reliable conclusions. The results of the study will provide the data for the needed arrangements in our educational program and for the development of knowledge, attitudes and behaviours of the students. Thank you for your participation.

Near East University Faculty of Medicine Public Health Department

Assoc. Prof. Dr. Ozen Asut, Prof. Dr. Songul Vaizoglu, Prof. Dr. Sanda Cali, Dr. Gulifeiya Abuduxike
Tel. 675 10 00/3026 ☐ I am willing to participate

Answer the following questions by encircling the appropriate choice and /or filling in the blanks.

Multiple choice questions have only one answer unless indicated otherwise.

1. Which grade are you in? 1. 2. 3. 4. 5. 6.
2. What is your gender? 1. Male 2. Female
3. State you nationality:
4. State your age:
5. Did you ever smoke or do you currently smoke cigarettes?
 1. No, never smoked
 2. Only tried
 3. Smoked once but currently do not smoke
 4. Yes, I currently smoke every day (at least 1 cigarette/day)
 5. Occasionally smoke (smoke less than one cigarette a day)
6. Do you use any tobacco products other than cigarettes? (More than one choice may be indicated)
 1. I don't use other tobacco products

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

2. Hookah (Narghile, Argilah)
3. Other (Please state)
7. How long have you been smoking or consuming a tobacco product?
1. Never smoked / never consumed any tobacco product
 2. Equal to or less than one year
 3. 2 – 5 years
 4. 6-10 years
 5. More than 10 years
8. How many cigarettes a day do you smoke?
1. Don't smoke
 2. Less than one cigarette a day
 3. 1-5 cigarettes a day
 4. 6-10 cigarettes
 5. 11-20 cigarettes
 6. More than 20
9. Are you exposed to the tobacco smoke of other people? (More than one choice may be selected)
1. No exposure
 2. Continuous exposure at home
 3. Visitors smoking at home
 4. People smoking when I visit someone or somewhere
 5. In cafés and restaurants I attend
 6. Other (Please state).....
10. Do you believe in and support the ban on smoking indoors? 1. Yes 2. No
11. Do you think that **doctors** should be non-smoking role models for patients and the community as a whole?
1. Yes 2. No

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>

12. Only smokers will answer: When did you start consuming cigarettes/other tobacco products?

1. Before starting the medical school
2. After starting the medical school

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Asut O, Vaizoglu S, Abduxcu G, Sanda C. The Smoking Status And Attitudes Of Medical School Students In A University Of Nicosia. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.1470- Available online at <https://cyprusjmedsci.com/EN>