Original Research

DOI: 10.5152/cjms.2021.224-328

Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC

Authors: Filiz Yarıcı¹, Betül Mammadov¹, Dilay Necipoğlu²

Institutions:

¹Near East University, Faculty of Health. Near East Boulevard, ZIP: 99138 Nicosia

² Near East University, Faculty of Nursing. Near East Boulevard, ZIP: 99138 Nicosia

Corresponding Author: Filiz Yarıcı

E-mail: filiz_atis01@hotmail.com

Received Date: 26.08.2021

Accepted Date: 26.10.2021

How to cite: Yarıcı F, Mammadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328

Abstract

Aim: This research has been conducted with the aim of evaluating vaginal discharge and genital hygiene in women between the ages of 18-49 who live in the TRNC.

Materials and Method: The research, which was planned as a descriptive study, was carried out on an online basis. The sample scope was determined as 95% confidence interval and 5% margin of error and at least 382 women using a well-known sampling formula and 408 women volunteered for this study. The data were collected using a personal information form, vaginal discharge question form and Genital Hygiene Behaviours Inventory (GHBI). Number, percentage and average tests were used for

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mammadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

the descriptive analysis. The conformity of the dependent and independent variables to the normal distribution was tested with the Kolmogorov-Smirnov Test. The Independent t Test and One-way ANOVA test were used to determine the difference between the dependent and independent variables.

Findings: This study 36.5% of the participants were between the ages of 25-34, 59.8% were married and in terms of level of education, 54.9% had university or higher level qualifications. 8.3% of the women had a previous medical history of a sexually transmitted disease and this was continuing in 3.9% of them. It was determined that 46.8% of the women participating in the study had abnormal vaginal discharge. It was determined that the abnormal discharge of 14.2% of the women lasted 1 month and for 13% of the women, it lasted for 1 year or more. Also, it was seen that 30.2% of the women resorted to traditional methods for dealing with their abnormal vaginal discharge, including 11.5% of them who cleaned their vagina with vinegar, soda or lemon juice. The average score that the women obtained from the “Genital Hygiene Behaviour Inventory” was 78.24 ± 12.07 (min=49.0, max=108.0). A statistically significant difference was found between the GHBI mean score of the women participating in the study according to their past experience of abnormal vaginal discharge, the presence of a previous sexually transmitted disease and vaginal douching ($p < 0.05$).

Conclusion: As a result, it was found that nearly half of the women had abnormal vaginal discharge (curd, green-yellow or grey in colour and foul-smelling). In addition, it was determined that 30.2% of women resorted to traditional methods for treating abnormal vaginal discharge, including 11.5% of the participants who cleaned their vagina with vinegar, soda and lemon juice. Within the scope of primary health care services, it is recommended that midwives and nurses question the genital hygiene behaviours of women by both planning home visits and during the visits to health institutions, and by providing accurate information on reproductive health, genital hygiene, sexually transmitted diseases and family planning through brochures or training booklets.

Key words: vaginal discharge, genital hygiene, women’s health

INTRODUCTION

Genital infections are recognised as the most common health problems seen in women in their reproductive stage between the ages of 18-49. Every year, one million women contract urogenital infections around the world and it has been reported that at least 75% of them have a history of vaginal infection (1–3). Research conducted in Turkey shows that the frequency of genital tract infections in women is between 52% and 92% (4–9).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

There are many reasons why genital infections are so common in women. The anatomic structure in which the urethra, vagina and anus are close together increases the risk of genital tract infections. Additionally, having multiple sexual partners, poor genital and menstrual hygiene, tight and synthetic underwear, bad nutrition, systemic diseases such as diabetes, poor obstetric history, stress, long term antibiotic and steroid use, incorrect genital hygiene applications, vaginal douche (VD), tampon use, poor environmental conditions, lack of education and many other factors can cause genital infections (9–13). Ensuring genital hygiene is one of the most important steps in preventing serious problems that can be caused by genital infections, as well as protecting and maintaining women's health and reproductive health. This is because when genital hygiene is neglected, the tendency for infection increases, and if the infection is not treated correctly and in a timely manner, this can have a negative impact on the woman's reproductive ability (1,8,9). Additionally, genital infections can result in problems such as babies with low birth weight, abortion, foetal death, congenital infections in newborns, ectopic pregnancy, sepsis, cervical cancer, chronic pelvic pain etc. (1,9,13). In order to protect and enhance women's health, it is important to provide early diagnosis and treatment services along with protective measures against genital infections, and for health professionals to provide education and counselling for women. (9).

The way each woman describes and perceives vaginal discharge is different because many women do not know their body or how to identify abnormal vaginal discharge. Some women perceive their vaginal discharge as abnormal and consult health institutions, whereas even though others may have excessive discharge, they think that this is normal and take no action. Women's perception of genital infection risks differ according to cultural structure, socioeconomic situation, environmental conditions, stories they hear from their loved ones, traditional methods they use for treatment, negative social exclusion and embarrassment with regard to gynaecological examinations (1,9). Due to these reasons, women often resort to traditional methods to solve their problems rather than using medical treatments. However genital infections that are not treated can cause serious reproductive health problems (1,6). Many studies have examined the genital hygiene behaviours of women in Turkey (9,14,15). However, no studies have evaluated the genital hygiene behaviours of women living in the TRNC. It is clear that midwives and nurses have an important role in protecting, maintaining and treating women's health conditions. Midwives in particular play an important role in determining the groups at risk of genital infections, evaluating the presence of genital infections in women and maintaining preventive health services in order to prevent their occurrence. For this reason, this study aims to evaluate the abnormal vaginal discharge and genital hygiene behaviours of women between the ages of 18-49 living in the TRNC.

Research Questions

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

1. Is there a statistically significant difference between the mean scores of the genital hygiene behaviors inventory of women according to some socio-demographic characteristics and vaginal discharge status?
2. Is there a significant difference between women's socio-demographic characteristics and differentiating between experiencing abnormal vaginal discharge?
3. Is there a significant difference between the socio-demographic characteristics of women and vaginal douching?

MATERIALS AND METHODS

Study design and sampling

The cross-sectional and descriptive study was carried out between 01.06.2021 and 30.06.2021 on an online basis. The universe of the study comprised n=77,920 women between the ages of 18-49 in accordance with the TRNC 2011 census. The sample, on the other hand, was calculated to be at least 382 women, with a 95% confidence interval and a 5% margin of error, according to the known sampling formula. The sample of the study consisted of n=408 women.

Data Collection

The data collection phase was conducted online. The data were collected by researchers at a specified time on a web basis with an online questionnaire form. Invitations to participate in this study were shared via social media accounts and the WhatsApp application. Individuals who were directed to the link address were presented with an informative and consent section. Individuals who provided their consent were then directed to the research form and participated in the study. Completing the online form took an average of 5 minutes.

Data Collection Tools

The data collection form consists of two sections. The first section includes all information regarding the study as well as a consent form. The second section consists of the data collection forms. In the first part of the introductory information form, there are 20 questions including information about age, marital status, socio-economic characteristics and obstetric history, and introductory features for vaginal discharge. The second part consists of 15 questions about vaginal discharge. These questions were prepared in line with the purpose of the study and their validity and reliability were not determined. While the scope validity was being evaluated regarding the items on the question

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

form, the expert opinions of five specialist lecturers of whom one specialised in the basics of nursing, one in public health nursing and three in the field of midwifery, were taken into account. Necessary changes were made to the data collection form according to the feedback from these experts. A pilot study of the data collection form was conducted with 20 women in the TRNC that were not included in the sample group. The necessary changes were made to the data collection form according to the results of the pilot study.

Genital Hygiene Behaviours Inventory (GHBI): The Genital Hygiene Behaviour Inventory (GHBI) is a one-dimensional scale consisting of 27 questions developed by Ege and Eryilmaz in 2005 (16). Every statement in the inventory is graded between 1 and 4. For the positive statements, one (1) point is given for the answer “Never”, (2) for “Sometimes”, (3) for “Often” and (4) for “Always. For the negative statements (1) is given for “Always”, (2) for “Often”, (3) for “Sometimes” and (4) for “Never”. Due to the fact that items 17, 26 and 27 in the inventory include negative statements, they are reverse scored. The minimum score that can be obtained from the GHBI is 27 and the maximum is 108. The total scores obtained from the inventory indicate the genital hygiene behaviour scores. As the scores obtained from the inventory increase, it is evaluated that the genital hygiene behaviours are satisfactory. The Cronbach’s alpha reliability coefficient is 0.86 (16). In the current study, the Cronbach’s alpha value of the scale was determined to be 0.77.

Data analysis

The data collected online were evaluated by transferring to the Statistical Package for Social Sciences software version (SPSS 21.0- IBM SPSS Corp.; Armonk, NY USA) package program. The Cronbach’s α value was calculated to test the internal reliability of the scale. The number, percentage and average tests were used for descriptive analysis. The conformity of the dependent and independent variables to normal distribution was tested with the Kolmogorov-Smirnov Test. The Independent t Test and One-way ANOVA test were used to determine the difference between the dependent and independent variables. The data were evaluated with a 95% confidence interval and $p=0.05$ margin of error.

Ethical considerations

Approval from the Scientific Research Ethics Board was obtained before starting the research (ethical board no: 2021/1367). It was stated to the participants in the informed consent section that the study would be voluntary, could be completed at any time and data would remain anonymous. No

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

payment was made or charged to the participants in the study. However, it was stated that they could receive summary information about the results of the study upon request.

RESULTS

The women's socio-demographic information is given in Table 1. 36.5% of the participants were between the ages of 25-34, 59.8% were married and the level of education of 54.9% was university or higher. The obstetric and gynaecological details of the women are given in Table 2. It was determined that 46.8% of the women used family planning, 15.9% of the women used condoms, and 10.5% used intrauterine devices for family planning. 8.3% of the participants had a previous sexually transmitted disease and 3.9% of them still had the disease. It was determined that 46.8% of the women in the study had abnormal vaginal discharge. It was determined that the discharge of 14.2% of the women lasted for 1 month or less, and 13.0% of them had discharge that lasted for 1 year or more. In addition, it was determined that 30.2% of women resorted to traditional methods for treating abnormal vaginal discharge, including 11.5% of the participants who cleaned their vagina with vinegar, soda and lemon juice.

The women who participated in this research had a mean score of 78.24 ± 12.07 (min=49.0, max=108.0) from the 'Genital Hygiene Behaviour Inventory'. A comparison of the GHBI mean scores according to the participants age and level of education is given in Table 3. According to this, statistically difference was found between the GHBI mean scores of the women according to their age and level of education ($p < 0.05$).

Table 4 shows a comparison of the GHBI mean scores of the participants according to certain obstetric and gynaecological specifications and their discharge situation. It was found that there was a statistically significant difference between the mean scores of the 'Genital Hygiene Behaviour Inventory' according to the women's status of distinguishing the normality of vaginal discharge, using perfume, deodorant, or soap to prevent discharge, and keeping the vagina dry to prevent discharge ($p < 0.05$). Additionally, a statistically significant difference was found between the GHBI mean scores of those who live with abnormal vaginal discharge, who have had a previous history of sexually transmitted disease and who apply vaginal douche ($p < 0.05$).

DISCUSSION

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

Genital tract infections, which are commonly seen in women, can be prevented and easily treated by changing certain hygiene habits and consulting health facilities in the presence of abnormal discharge. Therefore, the reproductive health, sexual health, emotional health and quality of life of women can be positively affected. According to the data of our study, in which the vaginal discharge and genital hygiene behaviours of 408 women in the TRNC was evaluated, it was determined that nearly half of the women had a history of abnormal vaginal discharge. When other studies are examined, it is seen that one out of every two women experience abnormal vaginal discharge due to poor genital hygiene habits (3,17–20). Our study showed that this was not only related to hygiene habits, but it was also determined that about a third of the participants found it difficult to distinguish abnormal discharge. Additionally, it was determined that more than half of the women who experienced abnormal vaginal discharge did not consult health services and that about a third resorted to traditional methods (vinegar, soda, lemon juice etc.) in the case of abnormal vaginal discharge. The study conducted by Okumuş and Demirci showed that although 22.8% of the women had abnormal vaginal discharge, they did not consult a health facility (21). The study conducted by Özcan et.al showed that 26,6% of the women waited for their abnormal vaginal discharge to clear up on its own and that 13.9% of them resorted to traditional methods to treat themselves (22). Many other studies have shown that women use traditional applications for their abnormal vaginal discharge and urogenital tract infections, which are commonly seen in women (23–25). In our study, the rates of applying to health services and using traditional methods in the case of abnormal vaginal discharge were found to be slightly higher compared to the literature. There are many reasons for this, such as delaying applying to health services for economic reasons, exchanging information about health among women, considering and abnormal vaginal discharge to be an infection that heals easily, being embarrassed about gynaecological examinations, and regarding the situation as normal.

Vaginal Douche (VD) is a common method applied by women and it is where the vagina is cleaned with water or antiseptic liquids (26). In our study, about a third of women stated that they applied vaginal douche. The study conducted by Yağmur in 2007 showed that 57.2% (8) of women applied vaginal douche, and in similar studies, it was reported that 88% (27), 59% (10), and 72.7% of women apply vaginal douche and that this practice increases the risk of infection by 2.5 times (28). Although it is not as high as in the literature, the number of women who apply vaginal douche in the TRNC is also high. Approximately one out of three women stated that they apply vaginal douche. The most common reasons that women apply vaginal douche includes “being hygienic” (2) as well as the belief that it is a contraceptive method and religious views (29). The results of our study show that vaginal douching is also a common practice in Northern Cyprus and is a risk factor for vaginal infections. It is thought that this practice, which is adopted as a hygiene practice, is actually an application that affects the pH of the vagina and increases the exposure to infection, and they are not aware of its potential harm.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

It is also thought that another factor that can increase the risk of vaginal infection is the family planning method they use as 10.5% of the women use intrauterine devices (IUD), while 15.9% use condoms as protection. It is known that IUDs change the vaginal flora and create a suitable environment for anaerobic bacteria causing infection (4). It is thought that the very low rate of condom use will increase the risk of sexually transmitted infections in women whose spouses do not use condoms, and may pave the way for the transmission of other infections that may cause abnormal vaginal discharge.

The mean score that the women obtained in this study from the Genital Hygiene Behaviours Inventory (GHBI) was determined to be 78.24 ± 12.07 (min:49-max:108). When it is considered that the highest score that can be obtained from the scale is 108, it can be said that the genital hygiene behaviours of the women are at a required level. In this study, it was found that the GHBI scores significantly increased with age (82.14 ± 10.15) ($p < 0.001$). It is known that most women experience genital infections at least once in their lifetime. Considering that 66.9% of the women participating in the study were young women under the age of 35, it is clear that these women will have a higher risk of contracting genital tract infections during their lifetime. The reason that the older women received higher scores from the inventory may be due to them going to the hospital more often after experiencing birth, discharge and infection and trying to apply the information that they have obtained from health workers.

It was found that the GHBI scores (80.77 ± 11.76) of the women with university or above level of education were higher than the women in the other groups ($p < 0.001$). Similar to our study, in a study conducted with women who applied to a health centre, it was reported that the GHBI mean score was 80.28 (min:48-max:107). It has been reported that hygiene behaviour is affected by education, being employed and increased income (8,30). It has been proven that raising the level of education and making women more productive in working life have positive effects on women's health. Educated women will have the awareness to seek health advice from the right sources, and to apply to a health institution instead of using traditional practices in order to receive the right treatment in abnormal situations.

It was found that women who obtained a higher score in the GHBI could identify abnormalities in their discharge better than those who received lower scores ($p < 0.001$), understood the importance of keeping their vagina dry ($p < 0.03$), did not apply vaginal douche ($p < 0.007$), and avoided applying products such as soap, deodorant and perfume to the perineal area ($p < 0.03$). Women with a previous history of sexually transmitted diseases had significantly higher GHDE scores than other women ($p < 0.005$), and this caused these women to be uncomfortable due to the negative effects that the infections had on their sexual health, reproductive health and quality of life, or they were afraid of the

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

possibility the infection would reoccur resulting in them having a more careful attitude towards genital hygiene behaviours.

As a result, it was determined that about half of the women experienced abnormal vaginal discharge. Genital tract infections are diseases that can be prevented and quickly treated with correct hygiene habits and timely treatment. However, when most women are faced with abnormal vaginal discharge, they do not apply to health establishments due to neglect, lack of education, embarrassment or economic reasons. Within the scope of primary health care services, it is recommended that midwives and nurses question the genital hygiene behaviours of women by both planning home visits and during visits to health institutions, and by providing accurate information on reproductive health, genital hygiene, sexually transmitted diseases and family planning with brochures or training booklets.

Limitations

Our study is limited to women who live in Northern Cyprus and agree to participate in the study and increasing the sample size of the study will increase the power of the study.

Main Points

- About half of the women participating in the study complained of abnormal vaginal discharge.
- About one-third of women use the traditional method for abnormal vaginal discharge.
- Vaginal douche is a common practice among women.
- Increasing age and educational status of women are associated with an increase in the scores they get from the "Genital Hygiene Behavior Inventory".
- In this study, the GHBI mean scores of the women participating in the study was determined to be high, and their Genital Hygiene behaviors were determined to be at the desired level.

REFERENCES

1. Daşikan Z, Kılıç B, Baytok C, Kocairi H, Kuzu S. The Genital Hygiene Practices of Women Who Have Genital Discharge Gynecology Outpatient Clinic. Gümüşhane University Journal of Health

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

Sciences. 2015 Jan 1;4(1):113–24.

2. Koştı N, Taşçı Beydağ K. Genital Hygiene Practices of Women who Present to a Gynecology Clinic. Ataturk University School of Nursing Journal. 2009;12(1):66–71.
3. Karatay G, Özvarış B. Evaluation of Applications Regarding the Genital Hygiene of Women Living in Barrel Houses Within a Region Existing a Health Center. Journal of the School of Nursing. 2006;10(1):7–14.
4. Zincir H, Temel AB. IUD Insertion According to Special IUD Consultation Principle and The Relation Between Genital Hygiene Training and The Incidence of Vulvovaginal Infections. J Heal Sci. 2010;19(1):60–7.
5. Arslan Özkan İ, Kulakaç Ö. Genital Hygiene Behaviors of Female Prisoners. J Anatolia Nurs Heal Sci. 2011;14(2):31–8.
6. Hacıaloğlu N, Nazik E, Kılıç M. A descriptive study of douching practices in Turkish women. Int J Nurs Pract. 2009;15(2):57–64.
7. Ünsal A, Özyazıcıoğlu N, Sezgin S. Attitudes About Genital Hygiene of Individuals Living in One Town and Nine Villages East Black Sea Region. J Anatolia Nurs Heal Sci. 2010;13(2):12–9.
8. Yağmur Y. The Genital Hygiene Behaviors of the Females Aged 15-49 Living at the Firat Health Clinic Neighborhood in Malatya. TAF Prev Med Bull. 2007;6(5):325–30.
9. Gözüyeşil E. Investigation of genital hygiene behavior: An example of a slum area. Ortadoğu Med J. 2020;12(2):186–93.
10. Özkan S, Demir Ü. Determining the effectiveness of the nurse in the diagnosis of vajinitis and finding out the factors causing the formation of vaginitis in the fertile age women between 15-49. Heal Soc. 2002;12(4):54–61.
11. Hotun Şahin N, Bilgiç D, Kızılkaya Beji N. Women's Health and Diseases for Nurses and Midwives. In: Reproductive System Infections. 1st ed. Istanbul: Nobel Medicine Bookstores; 2015.
12. Can Gürkan Ö, Arslan Özkan H. Reproductive system infections and sexually transmitted infections. In: Women's Health and Diseases for Nursing and Midwifery. 1st ed. Ankara: Academician Bookstore; 2019.
13. Yağmur Y, Ergin İO. Life Conditions and Genital Hygiene Practices In Seasonal Agricultural Women. J Int Soc Res. 2017;10(51):614–20.
14. Cangöl E, Tokuç B. The Evaluation of Genital Infections and Genital Hygiene Practices of Women

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

Who Applied to Gynecology Polyclinic. *Florance Nightingale Nurs J*. 2013;21(2):85–91.

15. Kisa S, Taskin L. Behavioral risk factors that predispose women to vaginal infections in Turkey. *Pak J Med Sci [Internet]*. 2010 [cited 2021 Jul 20];26(4):800–4. Available from: www.pjms.com.pk
16. Ege E, Eryılmaz G. The Effect of Planned Education Given to the Women on Genital Hygiene Behaviours. *Ataturk University School of Nursing Journal*. 2006;9(3):8–16.
17. Kisa S, Taşkın L. Behavioral risk factors that predispose women to vaginal infections in Turkey. *Pakistan J Med Sci*. 2010;26(4):800–4.
18. Dalbudak S, Bilgili N. GATA The genital hygiene behaviors of women who applied to the obstetrics and gynecology outpatient clinic and the effect of these behaviors on vaginal infection. *Gulhane Medical Journal*. 2013;55:281–7.
19. Yılmaz Y, Kahraman S. The knowledge about the adolescent girls' genitals and hygiene who live in Sanliurfa, applications and factors that affect. *J Hum Sci*. 2019;16(3):823–32.
20. Srivastava L. Reproductive Tract Infections among Women of Rural Community in Mewat, India: *J Health Manag*. 2010;12(4):519–38.
21. Okumuş F, Demirci N. Health-seeking behaviors of women living in rural areas in case of vaginal discharge. *Anadolu Nursing and Health Science Journal*. 2015;18(3):204–13.
22. Özcan H, Arık S, Esen ÜG, Aslan N. Perception of Vaginal Discharge of Young Women and Traditional Practices for Vaginal Discharge. *Gumushane University Health Science Journal*. 2020;9(3):272–9.
23. Darvishi M, Jahdi F, Hamzegardeshi Z, Goodarzi S, Vahedi M. The Comparison of vaginal cream of mixing yogurt, honey and clotrimazole on symptoms of vaginal candidiasis. *Glob J Health Sci*. 2015;7(6):108–16.
24. Felix T, Röder Brito D, Santos Pedroso R. Alternative and complementary therapies for vulvovaginal candidiasis. *Folia Microbiol (Praha)*. 2019;64(2):133–41.
25. Li C, Han H, Lee J, Lee M, Lee Y, Kim M. Knowledge, behaviors and prevalence of reproductive tract infections: a descriptive study on rural women in hunchun, china. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2010;4(3):122–9.
26. Çalışkan D. Should Traditional Intravaginal Application “Vaginal Shower, Lavage” Be Done? Shouldn't it be done? *TTB Journal of Continuing Medical Education*. 2005;14(1):15–8.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. *Cyprus J Med Sci* 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

27. Yılmaz N, Saracoglu P, Egilmez P. Feminine Hygiene Practice. Turkish Gynecology Journal. 2003;2(10):10–5.
28. Büşra Ç, Akar Y. Prevalence of Vaginitis Among 15-49 Age Women Registered in a Family Health Center. J Midwifery Heal Sci. 2020;3(2):100–14.
29. Çalışkan D, Çöl M, Akdur R, Yavuzdemir Ş, Yavuz Y. Study on Vaginal Shower Frequency and Effective Factors in Women aged 15-49 in the Park Health Center Region. Ankara University Faculty of Medicine Journalı. 1996;49(2):73–80.
30. Erbil N, Bölükbaş N, Kocabaş E. Genital Hygiene Behaviors of Married Women and Determination of Affecting Factors. In: Reproductive Health and Family Planning Congress Book. 2009. p. 199–120.

Table 1. Socio-demographic Characteristics of Women (n=408)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

Sociodemographic Characteristics		n	%
Age group	18-24 age	124	30,4
	25-34 age	149	36,5
	35-49 age	135	33,1
Marital status	Married	244	59,8
	Single	164	40,2
Educational Status	Middle school and below	74	18,1
	High	110	27,0
	University and above	224	54,9
Working Status	Working	234	57,4
	Not working	174	42,6
Economic Status	Low	19	4,7
	Medium	268	65,7
	High	121	29,7
Child Presence	Yes	201	49,3
	No	207	50,7

Table 2. Obstetric and Gynaecological Details of the Women (n=408)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

Obstetric and Gynecological Features		n	%
Family Planning Usage Status	Yes	191	46,8
	No	217	53,2
Used Family Planning Method	Condom	65	15,9
	Intrauterin device	43	10,5
	Tablet	32	7,8
	Tube ligation	25	6,1
	Other (monthly injection and traditional methods)	26	6,5
	No	217	53,2
Past sexually transmitted disease status	Yes	34	8,3
	No	374	91,7
Vaginal douche application status	Yes	146	35,8
	No	262	64,2
Abnormal vaginal discharge	Yes	191	46,8
	No	217	53,2
The case of applying to the health service for abnormal vaginal discharge	Yes	158	38,7
	No	250	61,3
Traditional method for abnormal vaginal discharge	Yes	123	30,2
	No	285	69,8

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

Used traditional method for abnormal vaginal discharge	Clean vagina with vinegar, soda or lemon juice	47	11,5
	Drink with onion, parsley, camomile tea	28	6,8
	Clean vagina with yoghurt	38	9,4
	Other (clean garlic or nettle's water)	10	2,5
	No	285	69,8

Table 3. Comparison of GHBI Mean Scores According to Some Socio-Demographic Specifications of Participants. (n=408)

Socio-demographic Characteristics		GHDE Mean±SD	F	p
Age groups	18-24 age	73,61±12,19	17,53	0,001
	25-34 age	78,57±12,32		
	35-49 age	82,14±10,15		
Educational status	Middle school and below	75,70±11,62	11,581	0,001
	High school	74,80±11,87		
	University and above	80,77±11,76		

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>

p<0,05 ; F: One-way ANOVA

Table 4. Comparison of GHBI Mean Scores of Participants according to Certain Obstetric and Gynaecological Specifications and Vaginal Discharge Situation. (n=408)

Vaginal Discharge Condition				GHDE Mean±SD	t	p
		n	%			
Distinguishing normality of vaginal discharge	Yes	283	69,4	82,55±10,02	12,619	0,001
	No	125	30,6	68,48±10,54		
Use of perfume, deodorant, soap to prevent vaginal discharge	Yes	52	12,7	74,84±9,92	-2,184	0,03
	No	356	87,3	78,74±12,28		
In abnormal vaginal discharge, the vagina should be kept dry	Yes	220	53,9	79,47±10,06	2,233	0,03
	No	188	46,1	76,80±13,94		
Abnormal vaginal discharge	Yes	191	46,8	79,51±10,27	1,940	0,05
	No	217	53,2	77,19±13,31		
Past sexually transmitted disease status	Yes	34	8,3	83,79±1,46	2,824	0,005
	No	374	91,7	77,74±12,09		
Vaginal douche application status	Yes	146	35,8	76,11±11,53	-2,679	0,007
	No	262	64,2	79,43±12,21		

p<0,05 ; t: Independent t Test

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please how to cite: Yarıcı F, Mamadov B, Necipoğlu D. Bakır M, Evaluation of Vaginal Discharge and Genital Hygiene Behaviors of Women aged 15-49 living in TRNC. Cyprus J Med Sci 2021; DOI: 10.5152/cjms.2021.224-328 - Available online at <https://cyprusjmedsci.com/EN>