1 **Supplementary information for manuscript**

2
3 **Organic aerosol in the summertime Southeastern United States: Components and**
4 **their link to volatility distribution, oxidation state and hygroscopicity**

5
6 E. Kostenidou^{1,2}, E. Karnezi³, J. R. Hite Jr⁴, A. Bougiatioti^{4,6}, K. Cerully^{5a}, L. Xu^{5,b}, N. L.
7 Ng^{5,4}, A. Nenes^{1,4,5,6*} and S. N. Pandis^{1,2,3*}

8
9 ¹Institute of Chemical Engineering Sciences, Foundation for Research and Technology,
10 Hellas, Patras, Greece

11 ²Department of Chemical Engineering, University of Patras, Patras, Greece

12 ³Department of Chemical Engineering, Carnegie Mellon University, Pittsburgh, USA

13 ⁴School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta,
14 GA, USA

15 ⁵School of Chemical and Biomolecular Engineering, Georgia Institute of Technology,
16 Atlanta, GA, USA

17 ⁶Institute for Environmental Research and Sustainable Development, National
18 Observatory of Athens, Palea Penteli, Greece

19 ^anow at: TSI, Inc., Shoreview, MN, USA

20 ^bnow at: Division of Geological and Planetary Sciences, California Institute of
21 Technology, Pasadena, CA, USA

22
23 *correspondence to athanasios.nenes@gatech.edu, spyros@chemeng.upatras.gr
24
25
26
27

Table S1. Volatility distribution, average volatility and vaporization enthalpy for each PMF factor and for the total OA.

OA Type	Saturation Concentration C^* ($\mu\text{g m}^{-3}$)				ΔH_{vap} (kJ mol^{-1})
	10^{-1}	1	10	Average C^*	
$a_m=1$					
MO-OOA	0.44	0.14	0.42	0.95 ± 0.31	89 ± 10
LO-OOA	0.27	0.19	0.54	1.88 ± 0.32	58 ± 13
Isoprene-OA	0.41	0.16	0.43	1.05 ± 0.30	63 ± 15
BBOA	0.47	0.29	0.24	0.59 ± 0.22	55 ± 11
Total OA	0.54	0.19	0.17	0.55 ± 0.29	86 ± 9
$a_m=0.1$					
MO-OOA	0.23	0.17	0.60	2.36 ± 3.33	100
LO-OOA	0.52	0.18	0.30	0.59 ± 3.17	98 ± 6
Isoprene-OA	0.60	0.10	0.30	0.49 ± 3.53	96 ± 8
BBOA	0.64	0.19	0.17	0.34 ± 2.48	86 ± 9
Total OA	0.65	0.09	0.25	0.40 ± 3.53	100
$a_m=0.01$					
MO-OOA	0.46	0.11	0.43	0.92 ± 3.74	150
LO-OOA	0.40	0.06	0.54	1.41 ± 3.32	121 ± 25
Isoprene-OA	0.32	0.12	0.56	1.72 ± 2.93	113 ± 22
BBOA	0.32	0.35	0.32	1.00 ± 2.47	100
Total OA	0.40	0.19	0.41	1.00 ± 3.16	100
$a_m=1, \text{CE}_{\text{TD}}=0.9*\text{CE}_{\text{BP}}$					
MO-OOA	0.41	0.24	0.35	0.86 ± 2.97	86 ± 9
LO-OOA	0.42	0.29	0.28	0.73 ± 3.69	63 ± 15
Isoprene-OA	0.39	0.25	0.36	0.95 ± 2.70	54 ± 10
BBOA	0.52	0.46	0.02	0.32 ± 1.89	54 ± 10
Total OA	0.19	0.19	0.62	2.64 ± 4.5	58 ± 13

Table S2: Volatility distribution and average volatility for specific vaporization enthalpies for each PMF factor and for the total OA.

OA Type	Saturation Concentration C^* ($\mu\text{g m}^{-3}$)			
	10^{-1}	1	10	Average C^*

$\Delta H_{\text{vap}}=50 \text{ kJ mol}^{-1}$				
MO-OOA	0.00	0.05	0.95	9.00±3.96
LO-OOA	0.25	0.05	0.70	2.80±3.15
Isoprene-OA	0.35	0.05	0.60	1.78±3.15
BBOA	0.65	0.05	0.30	0.45±3.15
Total OA	0.05	0.05	0.9	7.08±3.15

$\Delta H_{\text{vap}}=80 \text{ kJ mol}^{-1}$				
MO-OOA	0.35	0.15	0.50	1.41±3.15
LO-OOA	0.64	0.16	0.20	0.26±3.15
Isoprene-OA	0.75	0.05	0.20	0.28±3.15
BBOA	0.87	0.03	0.10	0.17±3.17
Total	0.45	0.25	0.3	0.71±3.15

$\Delta H_{\text{vap}}=100 \text{ kJ mol}^{-1}$				
MO-OOA	0.60	0.04	0.36	0.58±3.95
LO-OOA	0.90	0.04	0.06	0.14±3.96
Isoprene-OA	0.95	0	0.05	0.13±3.54
BBOA	0.96	0.04	0.00	0.11±3.15
Total OA	0.76	0.04	0.2	0.27±3.16

Table S3. Saturation vapor pressure (P^o) and saturation concentration (C^*) (at 298 K) for various acids.

Organic acid	$P^o(298K)$ (Pa*10⁻⁵)	$C^*(298K)$ (µg m⁻³)
Adipic acid		
Bilde et al. (2003)	0.73 (0.4)	0.43
Riipinen et al. (2007)	0.78 (0.43)	0.46
Saleh et al. (2009)	0.52 (0.3)	0.31
Yaws et al. (2003)	0.78 (0.43)	0.46
Azelaic acid		
Bilde et al. (2003)	0.44 (0.18)	0.34
Yaws et al. (2003)	0.81 (0.3)	0.62
Malonic acid		
Bilde et al. (2003)	8.35 (1.78)	3.5
Hyvärinen et al. (2006)	7.45 (1.58)	3.1
Suberic acid		
Bilde et al. (2003)	0.05 (0.14)	0.036
Pimelic acid		
Saleh et al. (2008)	7.2 (1.7)	4.65
Oxalic acid		
Booth et al. (2010)	2150 (860)	780
Glutaric acid		
Bilde and Pandis (2001)	75 (37)	39.9
Bilde (2003)	88 (44)	46.8
Succinic acid		
Bilde 2003	3.93	1.9
Hyvärinen et al. 2006	4.97	2.4
Saleh 2009	4.31	2.1
Yaws 2003	4.77	2.3
Levoglucosan		
May et al. (2012)		13 (2)
Pinonic acid		
Bilde (2001)	7	5.19
Salo et al. (2010)	0.42 (0.15)	0.312

54 **Table S4.** Hygroscopicity values (κ) for the same acids as in Table S4. Values in
55 parenthesis represent standard deviations.

Organic acid	κ
Adipic acid	
Cerully et al. (2014)	0.022 (0.002)
Kuwata et al. (2013)	0.002 (0.001)
Rissman et al. (2007)	0.059 (+0.021; -0.014)
Huff Hartz et al. (2006)	0.03 (+0.002; -0.001)
Broekhuizen et al. (2004)	0.096 (n/a)
Raymond and Pandis (2002)	0.02 (+0.018; -0.008)
Prenni et al. (2001)	0.014 (n/a)
Corrigan and Novakov (1999)	0.03 (n/a)
Cruz and Pandis (1997)	0.099 (+0.048; -0.029)
Azelaic acid	
Cerully et al. (2014)	0.061 (0.007)
Kuwata et al. (2013)	0.03 (0.01)
Huff Hartz et al. (2006)	0.022 (+0.018; -0.009)
Malonic acid	
Cerully et al. (2014)	0.281 (0.034)
Kumar et al. (2003)	0.227 (0.028)
Prenni et al. (2001)	0.237 (n/a)
Suberic acid	
Cerully et al. (2014)	0.007 (0.000)
Kuwata et al. (2013)	0.001 (n/a)
Pimelic acid	
Cerully et al. (2014)	0.213 (0.016)
Kuwata et al. (2013)	0.15 (0.01)
Frosch et al. (2010)	0.15 (0.04)
Huff Hartz et al. (2006)	0.14 (+0.109; -0.054)
Oxalic acid	
Sullivan et al. (2009)	0.5 (0.05)
Glutaric acid	
Reymond and Pandis (2002)	0.195 (0.082)
Koehler et al. (2006)	0.2 (0.08)
Succinic acid	
Cerruly 2014	0.285 (0.029)
Hori (2003)	0.231 (0.065)
Prenni (2001)	0.310
Corrigan and Novakov (1999)	0.225
Levoglucosan	
Svenningsson et al. (2006)	0.208 (0.015)
Koehler et al. (2006)	0.165 (0.015)
Pinonic acid	
Raymond and Pandis (2002)	0.106 (0.09)

Figure S1. Diagnostic plots of the PMF analysis: (a) Q/Q_{expected} versus the number of the examined factors, (b) Q/Q_{expected} versus the f_{peak} for the optimum solution (4 factors), (c) mass fraction of PMF factors versus the f_{peak} , (d) correlations of time series and mass spectra among the 4 PMF factors, (e) distribution of scaled residuals for each m/z and (f) time series of the measured and the reconstructed organic mass.

Figure S2. Model residuals $E = X - GF$ calculated using the PMF evaluation tool, PET (Ulbrich et al., 2009).

Figure S3. Comparison between the ambient PMF factors (crosses) as reported by Xu et al. (2015) and ambient plus TD OA PMF factors (bars) found in this work.

Figure S4. MFRs of the loss-corrected PMF OA factors and total OA for a lower TD CE. The circles denote the measurements with the one standard deviation of the mean for a 10% lower TD CE, the grey solid lines stand for the optimum solution if CE TD was 10% lower and the dash lines correspond to the predicted base case.

Figure S5. Predicted volatility distributions of the OA PMF factors and total OA for a lower TD CE. The error bars are estimated using the approach of Karnezi et al. (2014). The grey solid bars represent the results for a 10% lower TD CE. The green, blue, orange, red and purple bars stand for the base case solutions of MO-OOA, LO-OOA, Isoprene-OA, BBOA and total OA.

Figure S6. (a) and (b) Comparison between volatility compositions of OA for various studies: MILAGRO (Mexico City), MEGAPOLI (Paris winter and summer), Athens winter and FAME-08 (Finokalia, Crete). (c) and (d) Comparison between volatility distributions of Isoprene-OA and total OA estimated by other groups in Centreville during SOAS.

Figure S7. MFRs of the loss-corrected Isoprene-OA factor. The circles correspond to the measurements with the one standard deviation of the mean. The blue line is the predicted solution using as input the volatility distribution of IEPOX SOA of Lopez-Hilfiker et al. (2016).

Figure S8. O:C ratios versus the average volatility as $\log_{10}C^*$. The black isolines correspond to the theoretically intrinsic κ suggested by Nakao et al. (2017). The circles stand for the volatility and hygroscopicity measurements of known compounds.

References:

- Bilde, M., and Pandis, S. N.: Evaporation rates and vapor pressures of individual aerosol species formed in the atmospheric oxidation of α - and β -pinene, *Environ. Sci. Technol.*, 35, 3344-3349, 2001.
- Bilde, M., Svenningsson, B., Mønster, J., and Rosenørn, T.: Even-odd alternation of evaporation rates and vapor pressures of C3-C9 dicarboxylic acid aerosols, *Environ. Sci. Tech.*, 37, 1371, 2003.
- Booth, A. M., Barley, M. H., Topping, D. O., McFiggans, G., Garforth, A., and Percival, C. J.: Solid state and sub-cooled liquid vapour pressures of substituted dicarboxylic acids using Knudsen Effusion Mass Spectrometry (KEMS) and Differential Scanning Calorimetry, *Atmos. Chem. Phys.*, 10, 4879-4892, 2010.
- Broekhuizen, K., Kumar, P. P., and Abbatt, J. P. D.: Partially soluble organics as cloud condensation nuclei: Role of trace soluble and surface active species, *Geophys. Res. Lett.*, 31, L01107, doi: 10.1029/2003GL018203, 2004.
- Cerully, K. M., Hite, J., McLaughlin, M., and Nenes, A.: Towards the determination of joint volatility-hygroscopicity distributions: instrument development and response characterization for single-component aerosol, *Aerosol. Sci. Tech.*, 48, 296–312, 2014.
- Corrigan, C. E., and Novakov, T.: Cloud condensation nucleus activity of organic compounds: a laboratory study, *Atmos. Environ.*, 33, 2661-2668, 1999.
- Cruz, C. N., and Pandis, S. N.: A study of the ability of pure secondary organic aerosol to act as cloud condensation nuclei, *Atmos. Environ.*, 31, 2205-2214, 1997.
- Frosch, M., Zardini, A. A., Platt, S. M., Müller, L., Reinnig, M.-C., Hoffmann, T., and Bilde, M.: Thermodynamic properties and cloud droplet activation of a series of oxo-acids, *Atmos. Chem. Phys.*, 10, 5873-5890, 2010.
- Hori, M., Ohta, S., Murao, N., and Yamagata, S.: Activation capability of water soluble organic substances as CCN, *J. Aerosol. Sci.*, 34, 419–448, 2003.
- Huff Hartz, K. E. H., Tischuk, J. E., Chan, M. N., Chan, C. K., Donahue, N. M., and Pandis, S. N.: Cloud condensation nuclei activation of limited solubility organic aerosol. *Atmos. Environ.*, 40, 605–617, 2006.

186 Hyvärinen, A.-P., Lihavainen, H., Gaman, A., Vairila, L., Ojala, H., Kulmala, M., and
 187 Viisanen, Y.: Surface tensions and densities of oxalic, malonic, succinic, maleic,
 188 malic, and cis-pinonic acids. *J. Chem. Eng. Data*, 51, 255, doi:10.1021/je050366x,
 189 2006.

190 Koehler, K. A., Kreidenweis, S. M., DeMott, P. J., Prenni, A. J., Carrico, C. M., Ervens,
 191 B., and Feingold, G.: Water activity and activation diameters from hygroscopicity
 192 data - Part II: Application to organic species, *Atmos. Chem. Phys.*, 6, 795-809,
 193 2006.

194 Kumar, P. P., Broekhuizen, K., and Abbatt, J. P. D.: Organic acids as cloud condensation
 195 nuclei: Laboratory studies of highly soluble and insoluble species. *Atmos. Chem.*
 196 *Phys.*, 3, 509–520, 2003.

197 Kuwata, M., Shao, W., Lebouteiller, R., and Martin, S. T.: Classifying organic materials
 198 by oxygen-to-carbon elemental ratio to predict the activation regime of Cloud
 199 Condensation Nuclei (CCN), *Atmos. Chem. Phys.*, 13, 5309-5324, 2013.

200 Lopez-Hilfiker, F. D., Mohr, C., D'Ambro, E. L., Lutz, A., Riedel, T. P., Gaston, C. J.,
 201 Iyer, S., Zhang, X., Gold, A., Surratt, J. D., Lee, B. H., Kurten, T., Hu, W. W.,
 202 Jimenez, J., Hallquist, M., and Thornton, J. A.: Molecular composition and
 203 volatility of organic aerosol in the Southeastern U.S.: Implications for IEPOX
 204 Derived SOA. *Environ. Sci. Technol.*, 50, (5), 2200-2209, 2016.

205 May, A. A., Saleh, R., Hennigan, C. J., Donahue, N. M., and Allen L. Robinson, A. L.:
 206 Volatility of organic molecular markers used for source apportionment analysis:
 207 Measurements and implications for atmospheric lifetime, *Environ. Sci. Technol.*,
 208 46, 12435–12444, 2012.

209 Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., and Sherman, D. E.: The effects of low
 210 molecular weight dicarboxylic acids on cloud formation. *J. Phys. Chem. A.*, 105,
 211 11240–11248, 2001.

212 Raymond, T. and Pandis, S. N.: Cloud activation of single-component organic aerosol
 213 particles, *J. Geophys. Res.*, 107, 4787, doi: 10.1029/2002JD002159, 2002.

214 Riipinen, I., Koponen, I. K., Frank, G. P., Hyvarinen, A.-P., Vanhanen, J., Lihavainen, H.,
 215 Lehtinen, K. E. J., Bilde, M., and Kulmala, M.: Adipic and malonic acid aqueous

216 solutions: surface tensions and saturation vapor pressures. *J. Phys. Chem.*, 111,
 217 12995–13002, 2007.

218 Rissman, T.A., Varutbangkul, V., Surratt, J.D., Topping, D.O., McFiggans, G., Flagan,
 219 R.C., and Seinfeld, J.H.: Cloud condensation nucleus (CCN) behavior of organic
 220 aerosol particles generated by atomization of water and methanol solutions,
 221 *Atmos. Chem. Phys.*, 7, 2949-2971, 2007.

222 Saleh, R., Walkerb, J., and Khlystov, A.: Determination of saturation pressure and
 223 enthalpy of vaporization of semi-volatile aerosols: The integrated volume method,
 224 *Aerosol Science*, 39, 876–887, 2008.

225 Saleh, R., Shihadeh A., and Khlystov, A.: Determination of evaporation coefficients of
 226 semi-volatile organic aerosols using an integrated volume—tandem differential
 227 mobility analysis (IV-TDMA) method. *J. Aerosol Sci.*, 40, 1019-1029, 2009.

228 Salo, K., Jonsson, A. M., Andersson, P. U., and Hallquist, M.: Aerosol volatility and
 229 enthalpy of sublimation of carboxylic acids. *J. Phys. Chem. A.*, 114, 4586–4594,
 230 2010.

231 Sullivan, R. C., Moore, M. J. K., Petters, M. D., Kreidenweis, S. M., Roberts, G. C., and
 232 Prather, K. A.: Effect of chemical mixing state on the hygroscopicity and cloud
 233 nucleation properties of calcium mineral dust particles, *Atmos. Chem. Phys.*, 9,
 234 3303-3316, 2009.

235 Svenningsson, B., Rissler, J., Swietlicki, E., Mircea, M., Bilde, M., Facchini, M. C.,
 236 Decesari, S., Fuzzi, S., Zhou, J., Mønster, J., and Rosenørn, T.: Hygroscopic
 237 growth and critical supersaturations for mixed aerosol particles of inorganic and
 238 organic compounds of atmospheric relevance, *Atmos. Chem. Phys.*, 6, 1937-1952,
 239 2006.

240 Yaws, C. L.: *Yaws' Handbook of thermodynamic and physical properties of chemical*
 241 *compounds*, 2003.