Supporting information for

Field characterization of the PM_{2.5} Aerosol Chemical Speciation Monitor: insights into the composition, sources and processes of fine particles in Eastern China

Yunjiang Zhang¹⁻⁴, Lili Tang^{1,2}, Philip L. Croteau⁵, Olivier Favez³, Yele Sun⁶, Manjula R. Canagaratna⁵, Zhuang Wang¹, Florian Couvidat³, Alexandre Albinet³, Hongliang Zhang⁷, Jean Sciare⁸, André S. H. Prévôt⁹, John T. Jayne⁵, Douglas R. Worsnop⁵

¹Jiangsu Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Nanjing University of Information Science and Technology, Nanjing 210044, China

²Jiangsu Environmental Monitoring Center, Nanjing 210036, China

³Institut National de l'Environnement Industriel et des Risques, Verneuil-en-Halatte, 60550, France

⁴Laboratoire des Sciences du Climat et de l'Environnement, CNRS-CEA-UVSQ, Université Paris-Saclay, Gif sur Yvette, 91191, France

⁵Aerodyne Research, Inc., Billerica, Massachusetts 01821, United States

⁶State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

⁷Nanjing Handa Environmental Science and Technology Limited, Nanjing 211102, China

⁸The Cyprus Institute, Environment Energy and Water Research Center, Nicosia, Cyprus

⁹Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen PSI 5232, Switzerland

Correspondence to: L. L. Tang (lily3258@163.com) and Y. L. Sun (sunyele@mail.iap.ac.cn)

Pages 9

Figures 13

Figure S1. Time series and mass spectral profiles of 2-factor PMF solution of the PM₁ ACSM dataset at three different f_{peak} values.

Figure S2. Time series and mass spectral profiles of 2-factor PMF solution of the PM_{2.5} ACSM dataset at three different f_{peak} values.

Figure S3. Summary of key diagnostic plots of the PM₁-ACSM PMF results for 2-factor solution: (a) Q/Q_{exp} as a function of number of factors, (b) mass fraction of OOA and HOA as a function of FPEAK, (c) the box and whiskers plot showing the distributions of scaled residuals for each m/z, (d) a comparison of the measured mass with the PMF reconstructed mass, (e) time series of the residual diagnostics and Q/Q_{exp} for each point in time.

Figure S4. Summary of key diagnostic plots of the PM_{2.5}-ACSM PMF results for 2-factor solution: (a) Q/Q_{exp} as a function of number of factors, (b) mass fraction of OOA and HOA as a function of FPEAK, (c) the box and whiskers plot showing the distributions of scaled residuals for each m/z, (d) a comparison of the measured mass with the PMF reconstructed mass, (e) time series of the residual diagnostics and Q/Q_{exp} for each point in time.

Figure S5. Relationship between measured NH₄ and predicted NH₄ for both the PM_{2.5} and PM₁ ACSMs, respectively. The points in plots are colored by the ratio of [SO₄] / [SO₄/NO₃]. Note that predicted NH₄ is estimated by $18 \times (2 \times [SO_4/96] + [NO_3/62] + [Cl/35.5])$.

Figure S6. Ion balance of the water-soluble ions measured by the PM_{2.5} MARGA. Note that: anion equivalents = $[NH_4^+/18] + [Na^+/23] + [K^+/39] + [Mg^{2+}/12] + [Ca^{2+}/20]$, and cation equivalents = $[SO_4^{2-}/48] + [NO_3^-/62] + [Cl^-/35.5]$, in which chemical ions are in the unit of μg/m³.

Figure S7. The relationship of (a) predicted NO_3^- (estimated from NaNO_3 and $\text{Mg}(\text{NO}_3)_2$) vs measured MARGA NO_3^- ; (b) the sum of predicted NO_3^- and measured $\text{PM}_{2.5}$ ACSM NO_3^- vs measured MARGA NO_3^- ; (c) predicted Cl^- (estimated from NaCl) vs measured MARGA Cl^- ; and (d) the sum of predicted Cl^- and measured $\text{PM}_{2.5}$ ACSM Cl^- vs measured MARGA Cl^- .

Figure S8. Relationships between (a) PM_1 (measured by Met one BAM1020) and total $\text{PM}_{2.5}$ (measured by TEOM-FDMS and Met one BAM1020 respectively) mass loadings; and (b) non-refractory NR- PM_1 (measured by the PM_1 ACSM) and $\text{PM}_{2.5}$ (NR- $\text{PM}_{2.5}$ measured by the $\text{PM}_{2.5}$ -ACSM) for the entire study.

Figure S9. Sized-segregated diurnal variations of fine aerosol species and organic components.

Figure S10. Averaged mass spectra (MS) of OA for PM_1 and $\text{PM}_{2.5}$ ACSM during the new particle formation (NPF, Episode 2) and the fog event (Fog, Episode 5) periods, respectively.

Figure S11. The mass concentrations of NO₂, SO₂, and NH₃ as a function of the NR-PM_{2.5} SOA bins the total particle surface area (S_a) bins, respectively. The median (middle horizontal line), mean (solid squares), 25th and 75th percentiles (lower and upper box), and 10th and 90th percentiles (lower and upper whiskers) are shown for each bin.

Figure S12. The spatial distribution of monthly mean NO₂ column density (<http://www.temis.nl/airpollution/no2.html>) in October (a) and November (b) 2015, respectively.

Figure S13. 48 h trajectory arriving at for the new particle formation and growth (a, Episode 2) and foggy (b, Episode 5) events.