



Observations of the summertime atmospheric pollutants vertical distributions and the corresponding ozone production in Shanghai, China

Chengzhi Xing^{1#}, Cheng Liu^{1,2,3,9#*}, Shanshan Wang^{4*}, Ka Lok Chan^{5*}, Yang Gao⁶, Xin Huang⁷,
5 Wenjing Su¹, Chengxin Zhang¹, Yunsheng Dong³, Guangqiang Fan³, Tianshu Zhang³, Zhenyi Chen³,
Qihou Hu³, Hang Su^{8,10}, Zhouqing Xie^{1,2,3,9}, Jianguo Liu^{2,3}

¹School of Earth and Space Sciences, University of Science and Technology of China, Hefei, 230026, China

²Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, 361021, China

10 ³Key Lab of Environmental Optics & Technology, Anhui Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, Hefei, 230031, China

⁴Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention (LAP³), Department of Environmental Science and Engineering, Fudan University, Shanghai, 200433, China

⁵Meteorological Institute, Ludwig-Maximilians Universität München, Munich, Germany

15 ⁶College of Environmental Science and Engineering, Ocean University of China, Qingdao, 266100, China

⁷School of Atmospheric Sciences, Nanjing University, Nanjing, 210093, China

⁸Institute for Environmental and Climate Research, Jinan University, Guangzhou 511443, China

⁹Anhui Province Key Laboratory of Polar Environment and Global Change, USTC, Hefei, 230026, China

20 ¹⁰Biogeochemistry Department, Max Planck Institute for Chemistry, Mainz, 55020, Germany

[#]This two authors contributed equally

*Correspondence to: Shanshan Wang (shanshanwang@fudan.edu.cn), Ka Lok Chan (lok.chan@lmu.de), Cheng Liu(chliu81@ustc.edu.cn)

Abstract. Ground based Multi-Axis Differential Optical Absorption Spectroscopy (MAX-DOAS) and lidar measurements
25 were performed in Shanghai, China during May 2016 to investigate the summertime atmospheric pollutants vertical
distribution. In this study, vertical profiles of aerosol extinction coefficient, nitrogen dioxide (NO₂) and formaldehyde
(HCHO) concentrations were retrieved from MAX-DOAS measurement using the Heidelberg Profile (HeiPro) algorithm,
while vertical distribution of ozone (O₃) was obtained from an ozone lidar. Sensitivity study of the MAX-DOAS aerosol
profile retrieval shows that the a priori aerosol profile shape has significant influences on the aerosol profile retrieval.
30 Aerosol profiles retrieved from MAX-DOAS measurements with Gaussian a priori demonstrate the best agreements with
simultaneous lidar measurements and vehicle-based tethered-balloon observations among all a priori aerosol profiles. MAX-
DOAS measured tropospheric NO₂ Vertical Column Densities (VCDs) show a good agreement with OMI satellite
observations with Pearson correlation coefficient (R) of 0.95. In addition, measurements of the O₃ vertical distribution
indicate that the ozone productions do not only occur at surface level but also at higher altitudes (about 1.1 km). Planetary
35 boundary layer (PBL) height, horizontal and vertical wind fields information were integrated to discuss the ozone formation



at upper altitudes. The results reveal that enhanced ozone concentrations at ground and upper altitudes are not directly related to horizontal and vertical transportations. Similar patterns of O₃ and HCHO vertical distributions were observed during this campaign, which implies that the ozone productions near to the surface and at higher altitudes are mainly influenced by the abundance of volatile organic compounds (VOCs) in the lower troposphere.

40 1 Introduction

Air pollution has become one of the major environmental problems around the world. It is particularly serious in China due the rapid development of economy and industrialization. This problem directly affected the ecological environment and earth's radiation budget. It also has a series of adverse impacts on human health. Regional ozone pollution caused by photochemical reaction during summertime becomes a serious problem in China in the recent years. Previous studies of the
45 O₃ vertical distribution and temporal variation of O₃ within the boundary layer in northern China indicated the O₃ levels are closely correlated with the abundance of O₃ precursors (Tang et al., 2017a and 2017b). The major primary O₃ precursors in China are nitrogen oxides (NO_x), defined as the sum of nitric oxide (NO) and nitrogen dioxide (NO₂), and volatile organic compounds (VOCs) (Geng et al., 2007). Nitrogen dioxide and formaldehyde (HCHO) are the important constituents in the atmosphere playing key roles in both tropospheric and stratospheric chemistry (Seinfeld et al., 1998; Chan et al., 2015; Wang
50 et al., 2017). NO₂ contributes to the formation of secondary aerosols and participate in the catalytic formation of ozone in the troposphere (Crutzen 1975), while HCHO is one of the most important VOCs which can be used as a proxy for the total VOCs (Sillman, 1995; Duncan et al., 2010).

In the presence of sunlight, VOCs and NO_x contribute together to the formation of ozone in the troposphere (Crutzen, 1975; Seinfeld et al., 1998). The ambient ozone level strongly depends on both the relative and absolute amounts of VOCs
55 and NO_x (Geng et al., 2007; Tang et al., 2009; Tang et al., 2012). Moreover, there are many studies using HCHO/NO₂ column ratio as an indicator to determine surface O₃-NO_x-VOCs sensitivity in previous (Martin et al., 2004; Tang et al., 2012; Mahajan et al., 2015). However, there is a lack of observation for NO_x and HCHO vertical distribution in order to investigate the O₃ formation and atmospheric chemistry in the lower troposphere. Modelling studies of boundary layer ozone over northern China show different sensitivity of ozone regime in vertical, which is also impacted by the vertical circulation
60 (Tang et al., 2017a and 2017b). In addition, aerosols, particularly fine particles, are one of the major air pollutants in China. They play a key role in the Earth's climate and weather system. The chemical and physical properties of aerosols are strongly dependent on their compositions and sources. Previous studies show that secondary inorganic aerosols (sulfate and nitrate) are the dominating composition of the fine particles in both northern and eastern China (i.e., Shanghai and Beijing) (Du et al., 2011; Zhu et al., 2016). Therefore, we focus on analyzing the secondary formation process of aerosols and photochemical
65 pollution in this study.

Multi-Axis Differential Optical Absorption Spectroscopy (MAX-DOAS) is a passive remote sensing technique measuring spectra of scattered sunlight at different elevation angles. Combining with forward radiative transfer simulations,



MAX-DOAS measurements can provide vertical distribution information of aerosol extinction and trace gases concentration in the lower troposphere (Hönninger and Platt, 2002; Bobrowski et al., 2003; Hönninger et al., 2004; Wagner et al., 2004; Wittrock et al., 2004; Platt and Stutz, 2008). In the past decade, ground based MAX-DOAS has been widely used for atmospheric aerosol and trace gases measurements (Frieß et al., 2006; Irie et al., 2008; Li et al., 2010; Li et al., 2013). MAX-DOAS measurements are often used to validate satellite observations of atmospheric trace gases, e.g. NO₂, SO₂ and HCHO (Irie et al., 2008; Kramer et al., 2008; Ma et al., 2013; Wang et al., 2017). Light detection and ranging (lidar) is an active remote sensing measurement technique providing the quantitative range-resolved aerosol parameters. Aerosol vertical distribution obtained from Lidar measurements are useful for the validation of MAX-DOAS retrieval of aerosol extinction profiles (Irie et al., 2008; Lee et al., 2011). Advanced lidar systems can also provide vertical profiles of different atmospheric species, such as ozone and water vapour.

Shanghai is one of the four direct-controlled municipalities in China, with population of over 24 million. Owing to the rapid urbanization of Shanghai and its surrounding cities in Yangtze River Delta (YRD) region, air quality has become deteriorated and much more attractive (Geng et al., 2007; Chan et al., 2015). Shanghai has a large number of vehicles in China. Vehicle emissions were reported to contribute about 35% of the overall NO_x emission in Shanghai (Li et al., 2011; Hao et al., 2011). FengXian, where the experiment was performed, is one of the sub-districts of Shanghai (see Fig. S1) located in the north of Hangzhou gulf. The measurement site is mainly surrounded by agriculture area with only few industrial and traffic emissions. During the measurement campaign, this area was mainly affected by the East Asian Monsoon with prevailing winds being mainly south-easterly. Previous study shows that ozone pollution is particularly serious in rural areas of Shanghai during summer. Biogenic VOCs was found to be one of the major ozone precursors near the ground surface (Geng et al., 2007). However, there is a lack of observations for the investigation on the ozone formation in vertical.

In this paper, we present the MAX-DOAS and lidar measurements for the vertical distribution of aerosol and trace gases in FengXian during May 2016. The vertical profiles of aerosol and NO₂ were retrieved and also validated by balloon-based measurements. Vertical distributions of NO₂ and HCHO, as well as ozone profile, were used to investigate the ozone formation. In addition, horizontal and vertical fluxes of ozone were calculated from WRF-Chem simulations to estimate the ozone production at different altitudes.

2 Measurements and methodology

2.1 The MAX-DOAS measurements

The MAX-DOAS instrument operated at the measurement site in FengXian consists of a telescope, two spectrometers with temperature stabilized at 20°C and a computer acting as controlling and data acquisition unit. The viewing elevation angle of the telescope is controlled by a stepping motor. Scattered sunlight collected by the telescope is redirected by a prism reflector and quartz fibers to the spectrometer for spectral analysis. Two spectrometers (Ocean Optic HR2000+ and a



100 Maya2000Pro spectrometer) were used to cover both the UV (303-370 nm) and VIS (390-608 nm) wavelength ranges. The full width half maximum (FWHM) spectral resolution of the UV and Vis spectrometers are 0.5 and 0.3 nm, respectively. The field of view (FOV) of the instrument is estimated to be less than 1 °.

During the measurement period, the viewing azimuth direction was adjusted to the north. A full measurement sequence consists of 7 elevation angles, i.e. 3 °, 5 °, 8 °, 10 °, 15 °, 30 ° and 90 °, each with 100 scans. The exposure time is automatically
105 calculated depending on the intensity of the received scattered sunlight in order to achieving similar intensities for each elevation. The full measurement sequence takes about 5-15 mins depending on the scattered sunlight intensity. Dark current and offset spectra were measured by blocking incoming light by a mechanical shutter, and were subtracted from the measurement spectra before spectral analysis. In this study, data measured with solar zenith angle (SZA) of less than 75 ° were used to avoid strong influence from the stratosphere (Wang et al., 2014; Wang et al., 2017).

110 2.1.2 Spectral analysis

The MAX-DOAS measured spectra were analyzed using the software QDOAS which is developed by BIRA-IASB (<http://uv-vis.aeronomie.be/software/QDOAS/>). The DOAS fit results are the differential slant column densities (DSCDs), i.e. the difference of the slant column density (SCD) between the off-zenith spectrum and the corresponding zenith reference spectrum. Details of the DOAS fit settings are listed in Table 1. A typical DOAS retrieval example for the oxygen dimer (O₄),
115 NO₂ and HCHO is shown in Fig. S2. The stratospheric contribution was approximately eliminated by taking the zenith spectra of each scan as reference in the DOAS analysis. Before profile retrieval, DOAS fit results with root mean square (RMS) of residuals larger than 0.003 were filtered.

2.1.3 HeiPro algorithm description and retrieval parameter settings

Aerosol and trace gases (i.e., NO₂, HCHO) vertical profiles are retrieved from MAX-DOAS measurements using
120 HeiPro (Heidelberg Profile, developed by IUP Heidelberg) retrieval algorithm (Frieß et al., 2006; Frieß et al., 2011). The inversion algorithm is developed based on the Optical Estimation Method (OEM) (Rodgers, 2000), which employs the radiative transfer model SCIATRAN as the forward model. In general, the maximum a posteriori (MPA) solution \hat{x} is determined by minimizing the cost function χ^2 (Frieß et al., 2006; Frieß et al., 2011; Frieß et al., 2016), which can be expressed as

$$125 \chi^2(x) = [y - F(x, b)]^T S_e^{-1} [y - F(x, b)] + [x - x_a]^T S_a^{-1} [x - x_a] \quad (1)$$

The radiative transfer model or forward model $F(x, b)$ describes the measurement vector y (DSCDs at different elevation angles) as a function of the atmospheric state x (aerosol or trace gas profiles) and meteorological parameter b (i.e. pressure, temperature). The employed atmospheric pressure and temperature profiles were adapted from a climatology database, which contains various monthly and latitudinal dependent trace gases vertical profiles (Clémer et al., 2010; Hendrick et al.,
130 2014; Wang et al., 2014; Wang et al., 2016).



We assume a fix set of aerosol optical properties with asymmetry parameter of 0.69, a single scattering albedo of 0.90 and ground albedo of 0.05. These values are considered realistic for Shanghai according to (some measurements or previous studies, i.e. Chan et al., 2015). The lowest 4.0 km of the troposphere were divided into 20 layers, each with a thickness of 200 m. A fixed temporal interval of 15 min was used in the inversion. x_a denotes the a priori state vector serving as an additional constraint in the optimization. In order to investigate the impacts of a priori profile shape on the aerosol inversion, four different a priori extinction profiles available in the HeiPro algorithm, i.e. linearly decreasing, exponential decreasing, Boltzmann distribution (smoothed box-shaped) and Gaussian distribution (peaking shape), were used for the sensitivity study (Wang et al., 2016). Details of the sensitivity study are presented in Sect. 3.1. For NO₂ and HCHO retrieval, both the a priori profile x_a are exponentially decreasing with scaling height of 3 km, in which the surface concentration of NO₂ and HCHO is set to 10.5 and 1.5 ppb, respectively.

The covariance matrixes S_a and S_e describes the uncertainties and the cross correlation between different layers in the a priori and between measurements at different elevation angles, respectively. Another important quantity is the Jacobian matrix $A = \partial x / \partial x$. It represents the sensitivity of the retrieval to the true state. In addition, A provides the degree of freedom of signal (DFS), corresponding to the number of independent pieces contained in the measurement.

2.2 Lidar measurements

A polarization backscatter lidar was installed at the same experiment site collocating with the MAX-DOAS instrument. The lidar system is equipped with a diode-pumped Nd:YAG laser emitting laser pulses at 532 and 355 nm by doubling and tripling the laser frequency. The typical pulse energy of the laser is about 20 mJ with a pulse repetition frequency of 20 Hz. The laser beam is emitted with divergence of 0.25 mrad and 158 mm off-axis to the receiving telescope with a field of view of 0.5 mrad, resulting in an overlap height of about 195 m. A constant lidar ratio (S_p , extinction to backscatter ratio) of 50 sr was assumed in the lidar retrieval.

Another ozone lidar was applied to detect the O₃ profiles at the same time. The differential absorption lidar system emits laser pulse at 316 nm. The typical pulse energy of the laser is about 90 mJ with a pulse repetition frequency of 10 Hz. The laser beam is emitted with divergence of 0.3 mrad and 120 mm off-axis to the receiving telescope with a field of view of 0.5 mrad, resulting in an overlap height of about 300 m.

2.3 Ancillary data

Vehicle-based tethered-balloon observations were also preformed regularly at the measurement site during the campaign. The balloon measurement provides information of several meteorological parameters, including temperature, pressure, relative humidity, wind speed, as well as the atmospheric pollutants, i.e. PM_{2.5}, PM₁₀, NO₂ and O₃, from ground level up to 900 m above (Li et al., 2015).



In addition, a dynamical/chemical model (WRF-Chem) was used to study the temporal development and the formation of ozone. A detailed description of the model can be found in *Grell et al., 2005*. The simulation domain was set to cover an area of $1200 \times 1200 \text{ km}^2$ ($114^\circ\text{--}127^\circ \text{ E}$, $25^\circ\text{--}36^\circ \text{ N}$) in order to include a number of large cities in the YRD area. The horizontal resolution of the simulation is set to $12 \times 12 \text{ km}^2$ while vertical direction of the model is divided into 26 hybrid
165 pressure-sigma levels extending from the ground up to 17 km. This setting allows a better reconstruction of the atmospheric status and less impacts due to the diverse emissions of these cities.

3 Results and discussion

3.1 Dependency of retrieval on a priori profile

The inversion of aerosol extinction profiles was achieved by fitting the O_4 DSCD measurements to the forward model
170 simulations. Previous studies show that there is a systematic uncertainty on the O_4 absorption, in which the uncertainty of the O_4 absorption was estimated to be $\sim 25\%$ (Cl ner et al., 2010; Gro mann et al., 2013; Vlemmix et al., 2015). However, reason of the uncertainty is not yet well understood. The uncertainty is typically corrected by multiplying the O_4 absorption cross section with a constant correction factor (Wagner et al., 2009; Cl ner et al., 2010; Wagner et al., 2011; Chan et al., 2015; Wang et al., 2016). By comparing the measured and modelled O_4 absorptions, we estimated that the literature O_4
175 absorption cross section was underestimated by 20%. Therefore, a scaling factor of 1.2 was selected to multiply with the O_4 cross section for the O_4 retrieval band between 425 and 470 nm.

In order to investigate the impacts of different a priori on aerosol retrieval, a cloud-free day of 17 May 2016 was selected for the aerosol a priori sensitivity study. Fig. 1(a) shows the available configuration of aerosol a priori profile in the HeiPro algorithm, including linear, exponential, Boltzmann and Gaussian shapes. The corresponding retrieved aerosol
180 extinction profile is shown in the Fig. 1(b) together with lidar observation. The retrieval results using linear, exponential and Boltzmann a priori aerosol profile are very similar displaying the maximum aerosol extinction close to the ground. Aerosol extinction profile retrieved using the Gaussian a priori profile shows the best agreement with simultaneous lidar measurements exhibiting an elevated layer during this period. Furthermore, the diurnal aerosol extinction profiles retrieved with different a priori and the lidar measurements are shown in the right panel of Fig. 1. The retrieval with the Gaussian a
185 priori profile also shows a better consistency with lidar results during the whole day. The in situ measurements of particle mass concentrations can be also used to semi-quantitative validate the MAX-DOAS retrieval of aerosol extinction coefficients (Wang et al., 2016). We compared the retrieved aerosol extinction profiles to the balloon-based $\text{PM}_{2.5}$ measurements. As shown in Fig. S3(a), Aerosol profile retrieved using Gaussian a priori shows the best agreement with the balloon-based measurements, both measurements show a peak value at about 0.75 km above ground level.

190 Table 2 summarized the parameters of aerosol retrieval performance. The retrieval errors and resulting cost functions using Gaussian a priori are the smallest among all a priori profiles. Moreover, the DFS is about 2.96 when a Gaussian a priori was used. The DFS value suggests at least two independent pieces of information can be determined from the



195 measurements. So the sensitivity study indicates that the Gaussian a priori profile is the most realistic option for aerosol retrieval during this campaign. As a consequence, Gaussian aerosol profile is selected as the a priori for all aerosol retrieval in this study.

Besides, aerosols strongly influence the effective light path of scattered sun-light in the atmosphere and the Slant Column Densities (SCDs) of trace gases. Therefore, we have examined the sensitivity of trace gas retrieval to aerosol profile. Aerosol profiles retrieved with different a priori profile shapes were used in the differential air mass factor (Δ AMF) calculation for the NO_2 profile retrieval in the visible band of 425 to 470 nm. NO_2 profiles retrieved with different aerosol profiles are shown in Fig. 2(a). The result shows the retrieved vertical distributions of NO_2 are easily impacted by the introduced aerosol vertical distributions. The NO_2 profile retrieved by the Gaussian aerosol a priori is significantly different from the others. Using balloon-based measurements as a reference, the NO_2 profile retrieved with the Gaussian aerosol a priori shows the best agreement. For the other three retrievals, NO_2 concentrations at upper layers are significantly lower than the balloon measurement. Moreover, the NO_2 profile retrieved using Gaussian aerosol a priori profile as inputs is correlated better ($R=0.93$) with balloon-based NO_2 concentration profiles than others. In Fig. 2(b), the mean difference and standard deviations between the NO_2 profile retrieved using Gaussian aerosol a priori profile as inputs and balloon-based measured NO_2 profile ($26.14\% \pm 41.34\%$) is smaller than the other three retrieved NO_2 profiles. All these results indicate that aerosol profile scenarios are very important for the trace gas retrieval.

3.2 Temporal variations of NO_2

210 Time series of NO_2 concentrations profiles were retrieved from MAX-DOAS measurement using the HeiPro algorithm. In order to convert the NO_2 SCDs to tropospheric vertical column densities (VCDs), MAX-DOAS retrieved NO_2 profiles, lidar aerosol profiles, averaged temperature and pressure profiles measured by in situ instruments on the balloon were introduced as inputs in the radiative transfer model for the NO_2 Air Mass Factors (AMFs) calculation. Fig. 3 shows the temporal variations of hourly averaged tropospheric NO_2 VCDs and vertical profiles. During the measurement campaign, significant NO_2 VCD peaks were observed on 10 and 16 May 2016, respectively. By extracting NO_2 concentration at the lowest layer of the retrieval, the averaged ground level NO_2 mixing ratio of 12.4 ppb in FengXian area is generally much lower than that in city center of Shanghai (63.3 ppb) (available from <http://www.shanghaiair.sinnapp.com/>). The sectoral NO_x emissions in FengXian area can be divided into three major types, i.e. transportation, industrial + residential and power generation. These three emission sources contributed 83% (transportation), 15% (industrial + residential) and 2% (power generation) of the total NO_x emissions in FengXian (Chan et al., 2015). The emission inventory indicates that transportation emission plays a dominant role on the local NO_x concentration. 24 hours Air Mass Backward Trajectories (AMBTs) from 8:00 to 17:00 LST (Local Standard Time) at altitude of 500 m over the experimental site were calculated to assess the role of air mass transport during the NO_2 episode periods. As shown in Fig. S4, the peak values of tropospheric NO_2 VCDs are closely related to the wind direction. Increased NO_2 levels mainly occurred during northwesterly or northerly wind conditions, especially during the episodes on 10 and 16 May. Transportations of NO_2 can be also observed from OMI



satellite measurements during these episode days. Figure 4 shows the spatial distribution of tropospheric NO₂ VCDs from USTC OMI products on 10 and 16 May, respectively. Major industrial areas such as BaoShan, JiaDing, northern part of PuDongXi in Shanghai and a heavy industrial city (i.e., ZhangJiaGang) are located along the backward trajectories during the NO₂ episode days. In contrast, lower tropospheric NO₂ VCDs were observed during southerly and easterly wind conditions where the air masses were coming from unpolluted regions and the East China Sea.

The MAX-DOAS NO₂ measurements are also used to valid the USTC OMI NO₂ product (Liu et al., 2016). MAX-DOAS NO₂ VCDs were temporally averaged over the OMI overpass time of 12:00 to 13:00 LST, while the OMI NO₂ data are spatially averaged over pixels within 15 km of the experimental site. Previous study shows that a better approximation of trace gas and aerosol profiles for the tropospheric AMFs calculation can significantly improve the OMI NO₂ VCDs over polluted area by 35-40% and bring them closer to the ground based observations (Lin et al., 2014). Therefore, we have re-computed the OMI NO₂ tropospheric AMFs by using the combined NO₂ profiles, in which the lowest 3 km were adopted from the MAX-DOAS NO₂ profile retrieval, while NO₂ profiles above 3 km were taken from WRF-Chem simulations. Daily tropospheric NO₂ VCDs from MAX-DOAS measurement, NASA and USTC OMI product are shown in Fig. 5(a). The temporal trends of MAX-DOAS and USTC OMI data show similar characteristic. However, the MAX-DOAS measurements are systematically higher than OMI observations by 23% on average. The discrepancy can be explained by the averaging effect over the large OMI pixel which includes the neighboring clean areas. The correlations between MAX-DOAS and two different OMI products are shown in Fig. 5(b). The USTC OMI products agree better with the MAX-DOAS observations with Pearson correlation coefficient (R) of 0.95 (slope of 0.74 and offset of -2.09×10^{15} molec/cm²), while the correlation between MAX-DOAS and NASA OMI product is 0.71 with slope of 0.17 and offset of 2.69×10^{15} molec/cm². Compared to MAX-DOAS measurements and USTC OMI product, The NASA OMI NO₂ products report much lower NO₂ VCDs especially during these two NO₂ episode days. The result suggests that adopting local measurement of atmospheric parameters i.e. aerosols and trace gases profiles in AMF calculation could improve the accuracy of satellite VCDs products.

3.3 Ozone vertical distribution

Enhanced surface O₃ concentrations were found over rural areas of Shanghai compared to the city center (Geng et al., 2008; Xing et al., 2011). This is probably resulted from the significant contribution of anthropogenic emissions of NO in the city center which consumes ambient ozone through NO-titration (i.e. $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$). In this study, we focused on the formation pathways and the vertical distribution of O₃ in rural areas. In order to validate the vertical ozone distribution from lidar measurements, ozone profiles were compared to simultaneous ozone balloon-based measurements. The comparison result shows a good agreement with each other (see Fig. S3(b)-(d)). Time series of ozone vertical distributions measured by the lidar and surface ozone concentrations measured by the in situ monitor are shown in Fig. 6(a) and (b), respectively. The measurement result shows significant lower ozone concentrations on 15 and 20 May, 2016. Meteorological data shows that the solar irradiance was relatively low and associated with occasional rain on these two days. Decreases of ozone concentration were probably due to lower solar irradiance affecting the photochemical formation of ozone and rainy



condition favouring the wet removal pathway of atmospheric ozone. High ozone concentrations were observed during the
260 noontime (12:00-14:00) on 16 and 17 May (case 1), as well as all day of 18 May (case 2).

For Case 1 indicated in Fig. 6(a), surface O₃ concentrations measured by the in situ monitor correlated well with the
lidar observations at low altitudes, particularly for peak ozone values of 75-80 ppb during these two periods. On 16 May,
enhanced ozone values not only occurred at surface level, but also found at altitude of about 1.1 km. In contrast, high ozone
concentrations were only found at surface level but no extension to the high altitude at the same time period of 17 May. To
265 further investigate the causes of enhanced O₃ levels at upper altitudes on the 16 May, we calculated the O₃ fluxes in both
horizontal and vertical directions using WRF-Chem simulations (Jiang et al., 2008). The horizontal O₃ fluxes reckoned by
multiplying the horizontal wind speed with the O₃ concentration of the corresponding grid were illustrated in Fig. 7(a) to (d).
The vertical flux, shown in Fig. 7(e), is defined as the product of vertical wind speed and O₃ concentration at the
corresponding layer. Positive values represent upward transportation. To evaluate the accuracy of model simulations, the
270 simulated ozone concentrations were validated by the ozone lidar measurements. Figure 8 shows the correlation of model
simulation and lidar measurement from 0.09-1.5 km. Both datasets agree well with each other with Pearson correlation
coefficient (R) of 0.87 (slope of 0.97 and offset of 7.39 ppb). In addition, the root mean square error (RMSE), root mean
square error systematic (RMSEs), root mean square error unsystematic (RMSEu) and the index of agreement (d) in different
altitudes were analysed to quantify the differences between the measured and simulated values (Willmott, 1981; Geng et al.,
275 2007). A summary of the statistical analysis is listed in Table 3. The result indicates that the model is able to reproduce the
reality. The flux analysis in Fig. 7 shows that horizontal transportations at 900, 1000 and 1300 m were trivial (2.1, 11.7 and
1.1 ppb m s⁻¹ averaged net values) between 12:00 and 14:00 on 16 May. The result indicates that the horizontal
transportations only show negligible effects on the enhanced ozone concentrations at upper altitudes during this time period.
Since the vertical wind speed is relatively low (~0.01 m/s, in Fig. 6(c)), the vertical transportations was considered only to
280 play a minor role. Both the horizontal and vertical transportations were not significant, thus, enhanced ozone levels at upper
altitudes were mainly due to local formation.

In case 2, high O₃ concentrations were observed from ground surface to higher altitudes on 18 May. Ozone was mainly
concentrated at a layer of 0.9 km high from 03:00 to 8:00 LST. Then the ozone layer began to disperse to adjacent layers
from 0.5 to 1.3 km. And the ozone concentration gradually increased to more than 115 ppb from 0.5 to 0.9 km after 12:30.
285 Time series of the planetary boundary layer (PBL) height retrieved from the Mie lidar measurement is shown in Fig. 6(a)
(He et al., 2006). It is found that the PBL is relatively stable (at about 0.9 km) from 0:00 to 8:00 am on 18 May. It suggests
that the ozone layer was constrained above PBL height at that moment. Afterwards, the PBL height was subsequently rising
after sunrise due to the increase of air temperature. Owing to the rise of PBL height and the downward wind, ozone at upper
altitudes was gradually mixed and spread throughout the PBL from 9:00 to 12:30. After 12:30, the horizontally averaged net
290 flux of O₃ at 500, 900, 1000 and 1300 m are -1.9, 1.3, -0.44, and 1.1 ppb m s⁻¹, respectively. The vertical wind speed at
different altitudes is extremely low (<0.005 m/s). The above analysis implies that the increased ozone in the PBL after 12:30



were probably related to local formation. More details about the gas chemical analysis during the promoting O₃ formation were discussed in Sect. 3.4.

3.4 O₃-NO₂-HCHO in vertical

295 As discussed in Sect. 3.3, two high ozone concentration episodes was mainly locally formed. VOCs are the important precursors for the O₃ formation in urban areas (Kleinman et al., 2001; Zhang et al., 2004; Geng et al., 2007). Previous studies show that the formation of surface O₃ is mainly under VOC-sensitive regime in Shanghai (Geng et al., 2008). The production of O₃ is not only due to the abundance of VOCs, but also related to the reactions with OH radicals and solar irradiance. As HCHO is one of the major VOCs and strongly correlated with peroxy radicals (Sillman, 1995; Duncan et al., 2010), HCHO
300 measurement results were used as an indicator to represent the total VOCs here. Observations of NO_x and VOCs vertical distribution can provide indispensable information to investigate the ozone formation pathways.

Vertical distributions of NO₂ and HCHO were retrieved from the MAX-DOAS observations during the campaign. Fig. 9 shows the time series of NO₂, HCHO and O₃ vertical distributions during 15 to 20 May. Missing data is due to cloud filtering and/or low signal to noise ratio of the measurements. The distribution patterns of HCHO and O₃ above 500 m were
305 very similar. However, NO₂ were mainly concentrated below 500 m. Here we focus on 18 May to investigate the causes of high ozone concentration. As discussed in the Case 2 of Sect. 3.3, it is found that transportation contribution is trivial compared to the total O₃ concentration observed on this day. Moreover, we found the higher HCHO concentration (> 8 ppb) occurred prior to the increase of O₃ concentration. High HCHO levels during the time are mainly contributed by the oxidation of biogenic emissions of VOCs from plants, i.e., isoprene. Therefore, high abundance of VOCs and relatively
310 strong radiance contribute to higher formation rates of O₃ for Case 2 on 18 May.

3.5 Aerosol profile and evolution

Aerosols depolarization profiles were measured by the Mie-elastic backscatter polarization lidar during the campaign. Depolarization ratio is an indicator of the sphericity of aerosols. Low depolarization ratio indicates aerosols are spherical (Burton et al., 2012; Wong et al., 2017). Lidar results show that depolarization ratios of over 70% of measurements during
315 the campaign are between 0.001 and 0.03 (see Fig. S5). It demonstrates that aerosols in this area were dominated by spherical particles. In general, there are five major spherical particles in the atmosphere, namely black carbon, organic carbon, sulphate, nitrate aerosols and sea salt. Major sources of atmospheric aerosols in FengXian area could be the sea salt aerosols because the measurement site is close to the coast.

Fig. 10(a) shows an enhancement of aerosol extinction from 19:00 to 22:00 on 9 May, while depolarization ratios at 200
320 m were decreasing during the same period in Fig. 10(b). The growth of extinction coefficient is due to the emission of biofuel/biomass burning in the surrounding agriculture areas (Du et al., 2011). However, the decrease of depolarization ratios is most likely due the secondary processes in the atmosphere as the emission sources do not change rapidly. In Fig. 10(c), the reduction of depolarization ratios is observed prior to the enhancement of aerosol extinction coefficient from 19:00 to 22:00



on 9 of May. Decrease of aerosol depolarization ratio indicates that the aerosols became more spherical during that time. It is
325 related to the aerosol aging process which accompanied with the mixing between primary and secondary aerosols, as well as
the interactions among aerosols, trace gases and moisture in the atmosphere. In addition, NO₂ is an important precursor for
atmospheric nitrates particles formation (Myoseon and Kamens, 2001; Wang et al., 2017). Correlation between NO₂
concentrations and aerosol extinctions from 100m to 1000 m above ground level on 9 May is shown in Fig. 11. Strong
correlation is observed between ambient NO₂ and aerosol (R=0.63) indicating the significant contribution of NO₂ to the
330 secondary aerosol formation. Under high atmospheric ozone conditions, ambient NO₂ is oxidized rapidly to form nitrate
aerosols in the atmosphere of Shanghai (Du et al., 2011).

4 Summary and conclusions

In this paper, we present measurements of NO₂ and HCHO vertical profiles using ground based MAX-DOAS, while
aerosol and O₃ profiles were measured by lidar at Shanghai from 5 to 23 May 2016. Sensitivity study shows that the a priori
335 profile is playing an important role in the aerosol profiles retrieval. During the period of this campaign, the shapes of aerosol
profiles are similar to Gaussian vertical distribution in Shanghai. Accurate aerosol extinction profiles were found to be very
important for the retrieval of NO₂ and HCHO vertical distribution. Simultaneous measurements of NO₂ profiles obtained
from balloon-based in situ instrument agrees well with the MAX-DOAS data. In order to validate the OMI NO₂ VCDs, the
OMI satellite products from USTC and NASA were compared to the ground based MAX-DOAS observations. USTC OMI
340 data, using corresponding local trace gases profiles for the AMF calculation, present better correlation (R=0.95) than NASA
OMI's (R=0.71) with ground based MAX-DOAS measurements. According to the AMBTs analysis and the spatial
distribution of averaged OMI tropospheric NO₂ VCDs, the NO₂ pollution at FengXian were mainly influenced by
transportations from industrial areas located in the north and northwest of Shanghai (BaoShan and JiaDing) and south of
Jiangsu province (e.g. ZhangJiaGang).

345 O₃ vertical profiles were measured by lidar. Based on the analysis of horizontal and vertical fluxes of ozone at different
altitudes, we know that transportation is not a major influencing factor causing the increase of O₃. Similar vertical
distributions of HCHO and O₃ indicate the local formation was the dominant ozone source during the time. Moreover,
secondary aerosol formation process was found based on the analysis of aerosol extinction coefficient and depolarization
ratios. A positive correlation between NO₂ and aerosols during the campaign indicates the significant contribution of NO₂ to
350 total aerosols in the atmosphere.

Acknowledgements

This research was supported by grants from National Key Project of MOST (2016YFC0203302), National Natural Science
Foundation of China (41575021, 91544212, 41405117) and the Key Project of CAS (KJZD-EW-TZ-G06-01). We



acknowledge the NOAA Air Resources Laboratory (ARL) for making the HYSPLIT transport and dispersion model
355 available on the Internet (<http://ready.arl.noaa.gov/>). We thank Shanghai Environment Monitoring Center, Nanjing
University and East China University of Science and Technology of contributing to the balloon-based measurements. We
would like also to thank Hefei Institute of Physical Science, Chinese Academy of Sciences for the technical support of lidar
measurement.

360 References

- Aliwell, S., Van Roozendaal, M., Johnston, P., Richter, A., Wagner, T., Arlander, D., Burrows, J., Fish, D., Jones, R., and
Tørnkvist, K.: Analysis for BrO in zenith-sky spectra: An intercomparison exercise for analysis improvement, *J. Geophys.*
Res., 107, 4199, doi: 10.1029/2001JD000329, 2002.
- Bobrowski, N., Hönninger, G., Galle, B., and Platt, U.: Detection of bromine monoxide in a volcanic plume, *Nature*, 423,
365 273-276, 2003.
- Burton, S., Ferrare, R., Hostetler, C., Hair, J., Rogers, R., Obland, M., Butler, C., Cook, A., Harper, D., and Froyd, K.:
Aerosol classification using airborne High Spectral Resolution Lidar measurements-methodology and examples, *Atmos.*
Meas. Tech., 5, 73-98, doi: 10.5194/amt-5-73-2012, 2012.
- Chan, K. L., Hartl, A., Lam, Y. F., Xie, P. H., Liu, W. Q., Cheung, H. M., Lampel, J., Pöhler, D., Li, A., Xu, J., Zhou, H. J.,
370 Ning, Z., and Wenig, M. O.: Observations of tropospheric NO₂ using ground based MAX-DOAS and OMI measurements
during the Shanghai World Expo 2010, *Atmos. Environ.*, 119, 45-58, 2015.
- Chan, K. L., Wang, S. S., Liu, C., Zhou, B., Wenig, M. O., and Saiz-Lopez, A.: On the summertime air quality and related
photochemical processes in megacity Shanghai, China, *Sci. Total Environ.*, 580, 974-983, doi:
10.1016/j.scitotenv.2016.12.052, 2017.
- 375 Clémer, K., Van Roozendaal, M., Fayt, C., Hendrick, F., Hermans, C., Pinardi, G., Spurr, R., Wang, P., and De Mazière, M.:
Multiple wavelength retrieval of tropospheric aerosol optical properties from MAXDOAS measurements in Beijing, *Atmos.*
Meas. Tech. 3, 863-878, doi:10.5194/amt-3-863-2010, 2010.
- Crutzen, P.: Physical and chemical processes which control the production, destruction and distribution of ozone and some
other chemically active minor constituents, *WMO The Phys. Basis of Climate and Climate Modelling* 236-243 (SEE N 76-
380 19675 10-47), 1975.
- Du, H., Kong, L., Cheng, T., Chen, J., Du, J., Li, L., Xia, X., Leng, C., and Huang, G.: Insights into summertime haze
pollution events over Shanghai based on online water-soluble ionic composition of aerosols, *Atmos. Environ.*, 45, 5131-
5137, doi: 10.1016/j.atmosenv.2011.06.027, 2011.



- 385 Duncan, B. N., Yoshida, Y., Olson, J. R., Sillman, S., Martin, R. V., Lamsal, L., Hu, Y., Pickering, K. E., Retscher, C., Allen, D. J., and Crawford, J. H.: Application of OMI observations to a space-based indicator of NO_x and VOC controls on surface ozone formation, *Atmos. Environ.*, 44, 2213-2223, 2010.
- Fleischmann, O. C., Hartmann, M., Burrows, J. P., and Orphal, J.: New ultraviolet absorption cross-sections of BrO at atmospheric temperatures measured by time-windowing Fourier transform spectroscopy, *J. Photoch. Photobio. A*, 168, 117-132, 2004.
- 390 Friß U., Monks, P. S., Remedios, J. J., Rozanov, A., Sinreich, R., Wagner, T., and Platt, U.: MAX-DOAS O₄ measurements: A new technique to derive information on atmospheric aerosols: 2. Modeling studies, *J. Geophys. Res.*, 111, D14203, doi: 10.1029/2005jd006618, 2006.
- Friß U., Sihler, H., Sander, R., Pöhler, D., Yilmaz, S., and Platt, U.: The vertical distribution of BrO and aerosols in the Arctic: Measurements by active and passive differential optical absorption spectroscopy, *J. Geophys. Res.*: 116, D00R04, doi: 395 10.1029/2011JD015938, 2011.
- Friß U., Klein Baltink, H., Beirle, S., Clémer, K., Hendrick, F., Henzing, B., Irie, H., de Leeuw, G., Li, A., Moerman, M. M., van Roozendaal, M., Shaiganfar, R., Wagner, T., Wang, Y., Xie, P., Yilmaz, S., and Zieger, P.: Intercomparison of aerosol extinction profiles retrieved from MAX-DOAS measurements, *Atmos. Meas. Tech.*, 9, 3205-3222, doi: 10.5194/amt-9-3205-2016, 2016.
- 400 Geng, F., Zhao, C., Tang, X., Lu, G., and Tie, X.: Analysis of ozone and VOCs measured in Shanghai: A case study, *Atmos. Environ.*, 41, 989-1001, 2007.
- Geng, F., Tie, X., Xu, J., Zhou, G., Peng, L., Gao, W., Tang, X., and Zhao, C.: Characterizations of ozone, NO_x, and VOCs measured in Shanghai, China, *Atmos. Environ.*, 42, 6873-6883, 2008.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” 405 chemistry within the WRF model, *Atmos. Environ.*, 39, 6957-6975, 2005.
- Großmann, K., Friß U., Peters, E., Wittrock, F., Lampel, J., Yilmaz, S., Tschritter, J., Sommariva, R., von Glasow, R., Quack, B., Krüger, K., Pfeilsticker, K., and Platt, U.: Iodine monoxide in the Western Pacific marine boundary layer, *Atmos. Chem. Phys.*, 13, 3363-3378, doi: 10.5194/acp-13-3363-2013, 2013.
- Hao, N., Valks, P., Loyola, D., Cheng, Y., and Zimmer, W.: Space-based measurements of air quality during the World 410 Expo 2010 in Shanghai, *Environ. Res. Lett.*, 6, 044004, doi: 10.1088/1748-9326/6/4/044/004, 2011.
- He, Q. S., Mao, J. T., Chen, J. Y., and Hu, Y. Y.: Observational and modeling studies of urban atmospheric boundary-layer height and its evolution mechanisms, *Atmos. Environ.*, 40, 1064-1077, 2006.
- Hendrick, F., Müller, J. F., Clémer, K., Wang, P., De Mazière, M., Fayt, C., Gielen, C., Hermans, C., Ma, J. Z., Pinardi, G., Stavrakou, T., Vlemmix, T., and Van Roozendaal, M.: Four years of ground-based MAX-DOAS observations of HONO and 415 NO₂ in the Beijing area, *Atmos. Chem. Phys.*, 14, 765-781, doi: 10.5194/acp-14-765-2014, 2014.
- Hönninger, G., and Platt, U.: Observations of BrO and its vertical distribution during surface ozone depletion at Alert, *Atmos. Environ.*, 36, 2481-2489, 2002.



- Hönninger, G., von Friedeburg, C., and Platt, U.: Multi axis differential optical absorption spectroscopy (MAX-DOAS), *Atmos. Chem. Phys.*, 4, 231-254, doi: 10.5194/acp-4-231-2004, 2004.
- 420 Irie, H., Kanaya, Y., Akimoto, H., Iwabuchi, H., Shimizu, A., and Aoki, K.: First retrieval of tropospheric aerosol profiles using MAX-DOAS and comparison with lidar and sky radiometer measurements, *Atmos. Chem. Phys.*, 8, 341-350, doi: 10.5194/acp-8-314-2008, 2008.
- Jiang, F., Wang, T., Wang, T., Xie, M., and Zhao, H.: Numerical modeling of a continuous photochemical pollution episode in Hong Kong using WRF-chem, *Atmos. Environ.*, 42, 8717-8727, 2008.
- 425 Kleinman, L. I., Daum, P. H., Lee, Y.-N., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J., and Rudolph, J.: Sensitivity of ozone production rate to ozone precursors, *Geophys. Res. Lett.*, 15, 2903-2906, doi: 10.1029/2000GL012597, 2001.
- Kramer, L. J., Leigh, R. J., Remedios, J. J., and Monks, P. S.: Comparison of OMI and ground-based in situ and MAX-DOAS measurements of tropospheric nitrogen dioxide in an urban area, *J. Geophys. Res.*: 113, D16S39, doi: 430 10.1029/2007JD009168, 2008.
- Lee, H., Irie, H., Gu, M., Kim, J., and Hwang, J.: Remote sensing of tropospheric aerosol using UV MAX-DOAS during hazy conditions in winter: utilization of O₄ absorption bands at wavelength intervals of 338–368 and 367–393 nm, *Atmos. Environ.*, 45, 5760-5769, 2011.
- Li, J., Fu, Q., Huo, J., Wang, D., Yang, W., Bian, Q., Duan, Y., Zhang, Y., Pan, J., Lin, Y., Huang, K., Bai, Z., Wang, S.-H., 435 Fu, J. S., and Louie, P. K. K.: Tethered balloon-based black carbon profiles within the lower troposphere of Shanghai in the 2013 East China smog, *Atmos. Environ.*, 123, 327-338, 2015.
- Li, L., Chen, C. H., Fu, J. S., Huang, C., Streets, D. G., Huang, H. Y., Zhang, G. F., Wang, Y. J., Jang, C. J., Wang, H. L., Chen, Y. R., and Fu, J. M.: Air quality and emissions in the Yangtze River Delta, China, *Atmos. Chem. Phys.*, 11, 1621-1639, doi: 10.5194/acp-11-1621-2011, 2011.
- 440 Li, X., Brauers, T., Shao, M., Garland, R., Wagner, T., Deutschmann, T., and Wahner, A.: MAX-DOAS measurements in southern China: retrieval of aerosol extinctions and validation using ground-based in-situ data, *Atmos. chem. phys.*, 10, 2079-2089, doi: 10.5194/acp-10-2079-2010, 2010.
- Li, X., Brauers, T., Hofzumahaus, A., Lu, K., Li, Y. P., Shao, M., Wagner, T., and Wahner, A.: MAX-DOAS measurements of NO₂, HCHO and CHOCHO at a rural site in Southern China, *Atmos. Chem. Phys.*, 13, 2133-2151, doi: 10.5194/acp-13-445 2133-2013, 2013.
- Lin, J. T., Martin, R. V., Boersma, K. F., Sneep, M., Stammes, P., Spurr, R., Wang, P., Van Roozendaal, M., Cléner, K., and Irie, H.: Retrieving tropospheric nitrogen dioxide from the Ozone Monitoring Instrument: effects of aerosols, surface reflectance anisotropy, and vertical profile of nitrogen dioxide, *Atmos. Chem. Phys.*, 14, 1441-1461, doi: 10.5194/acp-14-1441-2014, 2014.
- 450 Liu, H., Liu, C., Xie, Z., Li, Y., Huang, X., Wang, S., Xu, J., and Xie, P.: A paradox for air pollution controlling in China revealed by "APEC Blue" and "Parade Blue", *Sci. Rep.*, 6, 34408, doi: 10.1038/srep34408, 2016.



- Ma, J., Beirle, S., Jin, J., Shaiganfar, R., Yan, P., and Wagner, T.: Tropospheric NO₂ vertical column densities over Beijing: results of the first three years of ground-based MAX-DOAS measurements (2008–2011) and satellite validation, *Atmos. Chem. Phys.*, 13, 1547–1567, doi: 10.5194/acp-13-1547-2013, 2013.
- 455 Martin, R. V., Fiore, A. M., and Van Donkelaar, A.: Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions, *Geophys. Res. Lett.*, 31, L06120, doi:10.1029/2004GL01941, 2004.
- Mahajan, A. S., De Smedt, I., Biswas, M. S., Ghude, S., Fadnavis, S., Roy, C., and van Roozendael, M.: Inter-annual variations in satellite observations of nitrogen dioxide and formaldehyde over India, *Atmos. Environ.*, 116, 194–201, doi: 10.1016/j.atmosenv.2015.06.004, 2015.
- 460 Meller, R., and Moortgat, G. K.: Temperature dependence of the absorption cross sections of formaldehyde between 223 and 323 K in the wavelength range 225–375 nm, *J. Geophys. Res.*, 105, 7089–7101, 2000.
- Myoseon, J., and Kamens Richards, M.: Characterization of secondary aerosol from the photooxidation of toluene in the presence of NO_x and 1-Propene, *Environ. Sci. Tech.*, 35, 3626–3639, 2001.
- Platt, U., and Stutz, J.: *Differential Optical Absorption Spectroscopy*, Springer, Heidelberg, Berlin, 229–375, 2008.
- 465 Rodgers, C. D.: *Inverse methods for atmospheric sounding: theory and practice*, World Scientific Publishing, Singapore-New Jersey-London-Hong Kong, 2000.
- Rothman, L., Gordon, I., Barber, R., Dothe, H., Gamache, R., Goldman, A., Perevalov, V., Tashkun, S., and Tennyson, J.: HITEMP, the high-temperature molecular spectroscopic database, *J. Quant. Spec. Ra.*, 111, 2139–2150, 2010.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics - From Air Pollution to Climate Change*, John Wiley, New York, 1998.
- 470 Serdyuchenko, A., Gorshchev, V., Weber, M., Chehade, W., and Burrows, J.: High spectral resolution ozone absorption cross-sections, *Atmos. Meas. Tech.*, 7, 625–636, doi: 10.5194/amt-7-625-2014, 2014.
- Sillman, S.: The use of NO_y, H₂O₂, and HNO₃ as indicators for ozone-NO_x-hydrocarbon sensitivity in urban locations, *J. Geophys. Res.*, 100, 14175–14188, doi: 10.1029/49JD02953, 1995.
- 475 Tang, G., Li, X., Wang, Y., Xin, J., and Ren, X.: Surface ozone trend details and interpretations in Beijing, 2001–2006, *Atmos. Chem. Phys.*, 9, 8813–8823, doi:10.5194/acp-9-8813-2009, 2009.
- Tang, G., Wang, Y., Li, X., Ji, D., Hsu, S., and Gao, X.: Spatial-temporal variations in surface ozone in Northern China as observed during 2009–2010 and possible implications for future air quality control strategies, *Atmos. Chem. Phys.*, 12, 2757–2776, doi:10.5194/acp-12-2757-2012, 2012.
- 480 Tang, G., Zhu, X., Xin, J., Hu, B., Song, T., Sun, Y., Zhang, J., Wang, L., Cheng, M., Chao, N., Kong, L., Li, X., and Wang, Y.: Modelling study of boundary-layer ozone over northern China - Part I: Ozone budget in summer, *Atmos. Res.*, 187, 128–137, doi: 10.1016/j.atmosres.2016.10.017, 2017a.
- Tang, G., Zhu, X., Xin, J., Hu, B., Song, T., Sun, Y., Wang, L., Wu, F., Sun, J., Cheng, M., Chao, N., Li, X., and Wang, Y.: Modelling study of boundary-layer ozone over northern China - Part II: Responses to emission reductions during the Beijing
- 485 Olympics, *Atmos. Res.*, 193, 83–93, doi: 10.1016/j.atmosres.2017.02.014, 2017b.



- Thalman, R., and Volkamer, R.: Temperature dependent absorption cross-sections of O₂-O₂ collision pairs between 340 and 630 nm and at atmospherically relevant pressure, *Phys. Chem. Chem. Phys.*, 15, 15371-15381, 2013.
- Vandaele, A. C., Hermans, C., Simon, P. C., Carleer, M., Colin, R., Fally, S., Merienne, M.-F., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 42 000 cm⁻¹ to 10 000 cm⁻¹ (238–1000 nm) at 220 K and 294
490 K, *J. Quant. Spectrosc. Ra.*, 59, 171-184, 1998.
- Vlemmix, T., Hendrick, F., Pinardi, G., De Smedt, I., Fayt, C., Hermans, C., PETERS, A., Wang, P., Levelt, P., and Van Roozendael, M.: MAX-DOAS observations of aerosols, formaldehyde and nitrogen dioxide in the Beijing area: comparison of two profile retrieval approaches, *Atmos. Meas. Tech.*, 8, 941-963, doi: 10.5194/amt-8-941-2015, 2015.
- Wagner, T., Dix, B. v., Friedeburg, C. v., Frieß, U., Sanghavi, S., Sinreich, R., and Platt, U.: MAX-DOAS O₄ measurements:
495 A new technique to derive information on atmospheric aerosols—Principles and information content, *J. Geophys. Res.*, 109, D22205, doi: 10.1029/2004JD004904, 2004.
- Wagner, T., Deutschmann, T., and Platt, U.: Determination of aerosol properties from MAX-DOAS observations of the Ring effect, *Atmos. Meas. Tech.*, 2, 495-512, doi: 10.5194/amt-2-495-2009, 2009.
- Wagner, T., Beirle, S., Brauers, T., Deutschmann, T., Frieß, U., Hak, C., Halla, J. D., Heue, K. P., Junkermann, W., Li, X.,
500 Platt, U., and Pundt-Gruber, I.: Inversion of tropospheric profiles of aerosol extinction and HCHO and NO₂ mixing ratios from MAX-DOAS observations in Milano during the summer of 2003 and comparison with independent data sets, *Atmos. Meas. Tech.*, 4, 2685-2715, doi: 10.5194/amt-4-2685-2011, 2011.
- Wang, S., Cuevas, C. A., Frieß, U., and Saiz-Lopez, A.: MAX-DOAS retrieval of aerosol extinction properties in Madrid, Spain, *Atmos. Meas. Tech.*, 9, 5089-5101, doi: 10.5194/amt-9-5089-2016, 2016.
- 505 Wang, T., Hendrick, F., Wang, P., Tang, G., Clémer, K., Yu, H., Fayt, C., Hermans, C., Gielen, C., Müller, J. F., Pinardi, G., Theys, N., Brenot, H., and Van Roozendael, M.: Evaluation of tropospheric SO₂ retrieved from MAX-DOAS measurements in Xianghe, China, *Atmos. Chem. Phys.*, 14, 11149-11164, doi: 10.5194/acp-14-11149-2014, 2014.
- Wang, Y., Lampel, J., Xie, P., Beirle, S., Li, A., Wu, D., and Wagner, T.: Ground-based MAX-DOAS observations of tropospheric aerosols, NO₂, SO₂ and HCHO in Wuxi, China, from 2011 to 2014, *Atmos. Chem. Phys.*, 17, 2189-2215, doi:
510 10.5194/acp-17-2189-2017, 2017.
- Willmott, C. J.: On the validation of models, *Physical geography*, 2, 184-194, doi: 10.1080/02723646.1981.10642213, 1981.
- Wittrock, F., Oetjen, H., Richter, A., Fietkau, S., Medeke, T., Rozanov, A., and Burrows, J.: MAX-DOAS measurements of atmospheric trace gases in Ny-Ålesund-Radiative transfer studies and their application, *Atmos. Chem. Phys.*, 4, 955-966, doi: 10.5194/acp-4-955-2004, 2004.
- 515 Wong, M. S., Qin, K., Lian, H., Campbell, J. R., Lee, K. H., and Sheng, S.: Continuous ground-based aerosol Lidar observation during seasonal pollution events at Wuxi, China, *Atmos. Environ.*, 154, 189-199, doi: 10.1016/j.atmosenv.2017.01.051, 2017.



Xing, J., Wang, S. X., Jang, C., Zhu, Y., and Hao, J. M.: Nonlinear response of ozone to precursor emission changes in China: a modeling study using response surface methodology, Atmos. Chem. Phys., 11, 5027-5044, doi: 10.5194/acp-11-5027-2011, 2011.

Zhang, R., Lei, W., Tie, X., and Hess, P.: Industrial emissions cause extreme urban ozone diurnal variability, Proc. Natl. Acad. Sci. USA, 101, 6346-6350, doi: 10.1073/pnas.0401484101, 2004.

Zhu, X., Tang, G., Hu, B., Wang, L., Xin, J., Zhang, J., Liu, Z., Münkler, C., and Wang, Y.: Regional pollution and its formation mechanism over North China Plain: A case study with ceilometer observations and model simulations, J. Geophys. Res. Atmos., 121, 14574-14588, doi: 10.1002/2016JD025730, 2016.

530

535

540

545

550

**Table 1. Setting for the O₄, NO₂, and HCHO DOAS spectral analysis**

Parameter	Data source	Fitting interval		
		O ₄	NO ₂	HCHO
Wavelength range		425-490nm	425-490nm	336.5-359nm
NO ₂	298K, I ₀ * correction (SCD of 10 ¹⁷ molecules/cm ²), Vandaele et al. (1998)	√	√	√
NO ₂	220K, I ₀ correction (SCD of 10 ¹⁷ molecules/cm ²), Pre-orthogonalized Vandaele et al. (1998)	√	√	×
O ₃	223K, I ₀ correction (SCD of 10 ²⁰ molecules/cm ²), Serdyuchenko et al. (2014)	√	√	√
O ₃	243K, I ₀ correction (SCD of 10 ²⁰ molecules/cm ²), Pre-orthogonalized Serdyuchenko et al. (2014)	×	√	√
O ₄	293K, Thalman and Volkamer (2013)	√	√	√
HCHO	297K, Meller and Moortgat (2000)	×	√	×
BrO	223K, Fleischmann et al. (2004)	×	×	√
H ₂ O	296K, HITEMP, Rothman et al. (2010)	√	√	×
Ring	Calculated with QDOAS	√	√	√
Polynomial degree		Order 5	Order 5	Order 5
Intensity offset		Constant	Constant	Constant

*Solar I₀ correction, Aliwell et al., 2002


555 **Table 2. Cost function, DFS and retrieved errors using different a priori profile in the aerosol retrieval**

The shape of a priori	Chi square	DFS	Retrieved error (<5%)	Smooth error (<5%)	Noise error (<5%)
Linearly shape	48.534512	2.957806	65%	65%	100%
Exponential shape	22.907515	2.9162457	75%	75%	100%
Boltzmann shape	28.533862	3.0331712	65%	65%	80%
Gaussian shape	2.9297998	2.9614936	100%	100%	100%



Table 3. Statistical analysis for the simulation of wind speed and ozone concentrations in different altitudes


	Altitudes(m)	RMSE	RMSEs	RMSEu	d
Wind	580	0.61	0.31	0.49	0.83
Speed	670	0.58	0.28	0.53	0.69
(m.s ⁻¹)	800	0.70	0.40	0.59	0.79
Ozone	580	6.6	3.2	6.5	0.63
(ppbv)	670	8.1	4.1	7.0	0.57
	800	7.3	3.7	6.7	0.59



565 Figure 1. Different aerosol extinction a priori and corresponding retrieval on 17 May 2016. (a) different a priori aerosol extinction profiles used in HeiPro. (b) shows four kinds of retrieved aerosol profiles and the lidar measured profile at 8:30. Diurnal aerosol extinction coefficient at 477 nm from (c) Mie-Scattering polarization lidar and retrieved using different a priori (d) Gaussian, (e) exponential, (f) linear and (g) Boltzmann.

570

575




580 **Figure 2.** Comparison of NO₂ profiles retrieved using four different aerosol a priori profiles as inputs and measured
by tethered-balloon. (a) shows the four retrieved vertical NO₂ concentrations and balloon-based measurement, as well
as the correlation coefficients. (b) shows the mean differences and standard deviations between four different
retrieved NO₂ profiles and the balloon-based NO₂ profiles.

585

590

595

600




605 **Figure 3. Time series of hourly averaged (a) NO₂ VCDs and (b) NO₂ vertical profiles from MAX-DOAS measurements.**

610

615

620



625

Figure 4. Spatial distribution of OMI tropospheric NO₂ VCDs on (a) 10 and (b) 16 May, 2016.


630

635

640

645

650



655 **Figure 5. Comparison of tropospheric NO₂ VCDs between ground based MAX-DOAS measurement and OMI satellite observation. (a) shows time series of daily averaged tropospheric NO₂ VCDs. MAX-DOAS data are averaged around the OMI overpass time. (b) shows the correlation of daily averaged tropospheric NO₂ VCDs measured by MAX-DOAS with USTC OMI and NASA OMI satellite data. The OMI measurements are spatially averaged over the grid cells within 15 km of ground location around the campaign site.**

660

665

670

675




Figure 6. Time series of (a) ozone vertical distributions measured by ozone lidar, (b) surface O₃ concentrations detected from in situ ozone instrument and (c) vertical wind profiles simulated by WRF in FengXian from 15 to 20
680 May, 2016.

685

690

695




Figure 7. Ozone horizontal transport fluxes at (a) 500 m, (b) 900 m, (c) 1000 m and (d) 1300 m, as well as (e) vertical transport fluxes.

700

705

710





Figure 8. Correlation of O_3 concentration at different altitudes between WRF-Chem simulation and lidar measurement.

715

720

725

730




735

Figure 9. Time series of retrieved (a) NO_2 and (b) HCHO vertical profiles from MAX-DOAS, as well as (c) O_3 vertical profiles measured by lidar.

740

745

750




755

Figure 10. Aerosol extinction coefficient (a) and depolarization ratio (b) on 9 May. (c) shows time series of aerosol extinction and depolarization ratios at 200 m from 19:00 to 22:00 on 9 May, 2016.

760

765

770



775 **Figure 11.** Scatter plots of aerosol extinction coefficient measured by Mie-Scattering depolarization lidar versus MAX-DOAS measurement of NO₂ from 100 to 1000 m above ground level.