1   **Concentrations and source regions of light absorbing**
2   **impurities in snow/ice in northern Pakistan and their**
3   **impact on snow albedo**

4   Chaman Gul^{1,2,5}, Siva Praveen Puppala², Shichang Kang^{1,3,5}, Bhupesh Adhikary²,
5   Yulan Zhang¹, Shaukat Ali⁴, Yang Li³, Xiaofei Li¹

6   ¹State Key Laboratory of Cryosphere Science, Northwest Institute of Eco-Environment and Resources,
7   Chinese Academy of Sciences, Lanzhou 73000, China

8   ²International Centre for Integrated Mountain Development (ICIMOD), G.P.O. Box 3226, Kathmandu,
9   Nepal

10   ³CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing, 100101, China

11   ⁴Global Change Impact Studies Centre (GCISC), Ministry of Climate Change, Islamabad, Pakistan

12   ⁵University of Chinese Academy of Sciences, Beijing, China

13   *Correspondence to:* Chaman Gul (chaman.gul@icimod.org; chaman@lzb.ac.cn)

14

15

16 **Figure S1. CALPISO extinction profile for a) winter (December, January), and b) summer (May, June)**

a. Debris covered glacier

b. Dirty glacier surface

c. Aged snow thickness

d. Clean and dirty snow

Figure S2. Appearance of different summer snow/ice sampling sites on a) b) Gulkin glacier, c) near Sachin glacier, and d) Sachin glacier

20

21

22 **Figure S3. a) Average concentration of EC at sites S1 to S6 sampled in winter and glaciers sampled in summer; b) individual sample concentration in winter**
23 **samples.**

24

25

26

27 **Figure S4. Contribution of subtype aerosols (%) in the atmosphere along the track of CALIPSO during the selected month June in 2006-2014**

28

a. Low snow thickness (0 – 8 cm)

b. Fresh snow (Kalam-S6)

c. Aged snow (Sost-S1)

d. Snow on slope

29

30 **Figure S5. Appearance of different snow areas sampled in winter**

31

32

33 **Figure S6. Albedo reduction (%) range between sulfate coated and uncoated black carbon calculated for the low concentration site S6.**

Figure S7. Concentration of black carbon (BC1, BC2 and BC) on the Sachin glacier calculated using the WRF-STEM model: a) summer, b) winter.

37 **Tables**

38 **Table S1 Parameters used for sensitivity analysis with SNICAR model for winter snow samples under the input parameters of direct incident radiation and mid-**
39 **latitude winter, clear-sky conditions.**

40 1 = solar zenith angle

41 2 = snow grain effective radius (μm)

42 3 = snowpack thickness (m)

43 4 = snowpack density (kg m^{-3})

44 5 = albedo of underlying ground (a. visible, 0.3-0.7 μm , b. near-infrared, 0.7-5.0 μm)

45 6 = MAC scaling factor (experimental) for BC

46 7= BC concentration (ppb, uncoated)

47 8 = dust concentration (ppm, 5.0–10.0 μm diameter)

48 9 = volcanic ash concentration (ppm)

49 10 = experimental particle 1 concentration (ppb)

50

Site	1	2	3	4	5a	5b	6	7	8	9	10
Kalam-S6	80.34	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	71.25	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	63.96	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	59.21	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	57.66	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	59.54	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	64.57	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0

Kalam-S6	72.07	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	81.31	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Kalam-S6	88.95	150	0.08	200	0.2	0.4	7.5/11/15	100	5.5	0	0
Sost-S1	82.61	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	73.66	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	66.45	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	61.64	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	59.66	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	61.36	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	65.93	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	72.98	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	81.80	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0
Sost-S1	88.95	350	0.04	320	0.19	0.35	7.5/11/15	1000	57	0	0

52 **Table S2. EC concentrations in glacier ice and snow and plains snow in north Pakistan, and other regions**

Location	Lat/long	Elevation (masl)	Min. EC (ngg ⁻¹)	Max. EC (ngg ⁻¹)	Time period (years)	Sample type/snow age	Reference
Mera Glacier, Nepal	27.72°N 86.8°E	6376	3	13	1999–2010	Ice core (rBC)	Ginot et al., 2014
Numanani, China	30.45°N 81.27°E	5900	4	14	Annual mean 2004	Ss (EC)	Xu et al., 2006
Greenland	72.6°N, 38.5°W	3209	4.2	30.1	1994 – 1996		Slater et al., 2002
NCO-P, Nepal	27.95°N, 86.82°E	5079	26	68.2	March – May 2006		Yasunari et al., 2010
Kangwure, China	28.47°N 85.82°E	6000	22	122	Annual mean 2001	Ss (EC)	Xu et al., 2007
Qiangyong, China	28.83°N 90.25°E	5400	43	143	Annual mean 2001	Ss (EC)	Xu et al., 2008
Zhadang, China	30.47°N 90.50°E	5802	334	473	2001–2010		Qu et al., 2014
Muji glacier, China	39.19°N 73.74° E	5062	25	731	2012		Yang et al., 2015
Zhadang, China	30.47°N 90.5°E	5802	114	1114	2005–2006		Ming et al., 2009
Tien Shan, China	43.1°N 86.82°E	4050		3000	2011	bottom firn-pack	Xu et al., 2012
Mera glacier, Nepal	27.72°N 86.8°E	5400		3535	2009		Kaspari et al., 2014
Northern China			60	4020	Jan-14		Wang et al., 2016
Urumqi glacier, China	43.10°N 86.80°E	4040	16	4093	2013		Ming et al., 2016
LHG glacier, China	39.17°N 96.17°E	5480		28,636	summer 2013-14	superimposed ice	Li et al., 2016
Present study							
Passu glacier	36.45°N 74.85°E	2838	87	734	May-June 2015	DCI (EC)	This study
Sachin glacier	35.32°N 74.76°E	3666	492	1789	May-June 2015	DCI & snow (EC)	This study
Sachin glacier	35.32°N 74.76°E	3666	543	3478	Oct-16	DCI & snow	This study
Gulkin glacier	36.42°N 74.77°E	3066	266	3574	Oct-16	DCI	This study
Mear glacier	36.15°N 74.82°E	3281	222	3656	May-June 2015	DCI (EC)	This study

Gulkin glacier	36.42°N 74.77°E	3066	81	5676	May-June 2015	DCI (EC)	This study
Barpu glacier	36.18°N 74.08°E	3055	877	5994	May-June 2015	DCI (EC)	This study
Henarche glacier	36.05°N 74.57°E	2941	778	10,502	May-June 2015	Surface ice (EC)	This study
S6-Kalam	35.43°N 72.6°E	1958	79	123	8-Jan-16	1–2 days	This study
S3-Tawas	36.39°N 73.36°E	2437		650	2-Jan-16	8–17 days	This study
S2- Hopar	36.22°N 74.75°E	2698	229	1064	1-Jan-16	1–15 days	This study
S5-Shangla	34.83°N 74.82°E	2348	367	1111	6–7 Jan 2016	8–9 days	This study
S4-Astore	35.37°N 74.80°E	2302	450	2640	3–4 Jan 2016	4–7 days	This study
S1- Sost	36.78°N 74.93°E	2951	482	5957	30–31 Dec 2015	2–17 days	This study

54 Table S3. Broadband albedo values at 0.975 μm wavelength for selected solar zenith angles, and daily mean albedo reduction (%)

Pollutant	Site S6 (low EC and dust)				Site S1 (high EC and dust)			
	SZA	MAC 7.5	MAC 11	MAC 15	SZA	MAC 7.5	MAC 11	MAC 15
EC only	80.34	0.79	0.78	0.77	82.61	0.59	0.56	0.52
	71.25	0.77	0.75	0.74	73.66	0.54	0.50	0.46
	63.96	0.75	0.74	0.72	66.45	0.50	0.46	0.42
	59.21	0.74	0.72	0.71	61.64	0.48	0.43	0.40
	57.66	0.74	0.72	0.71	59.66	0.47	0.43	0.39
	59.54	0.74	0.72	0.71	61.36	0.48	0.43	0.39
	64.57	0.75	0.74	0.72	65.93	0.50	0.46	0.42
	72.07	0.77	0.76	0.74	72.98	0.54	0.49	0.46
	81.31	0.80	0.78	0.77	81.80	0.59	0.55	0.51
	88.95	0.82	0.81	0.80	88.95	0.64	0.60	0.57
Daily mean reduction (%)		1.77	2.35	2.87		8.70	10.49	12.03
Dust only	80.34	0.83	0.83	0.83	82.61	0.76	0.76	0.76
	71.25	0.81	0.81	0.81	73.66	0.73	0.73	0.73
	63.96	0.80	0.80	0.80	66.45	0.71	0.71	0.71
	59.21	0.79	0.79	0.79	61.64	0.70	0.70	0.70
	57.66	0.79	0.79	0.79	59.66	0.70	0.70	0.70
	59.54	0.79	0.79	0.79	61.36	0.70	0.70	0.70
	64.57	0.80	0.80	0.80	65.93	0.71	0.71	0.71
	72.07	0.81	0.81	0.81	72.98	0.73	0.73	0.73

	81.31	0.83	0.83	0.83	81.80	0.76	0.76	0.76
	88.95	0.85	0.85	0.85	88.95	0.78	0.78	0.78
	Daily mean reduction (%)	0.07	0.07	0.07		0.05	0.05	0.05
EC and dust	80.34	0.79	0.78	0.77	82.61	0.59	0.55	0.52
	71.25	0.77	0.75	0.74	73.66	0.54	0.50	0.46
	63.96	0.75	0.73	0.72	66.45	0.50	0.46	0.42
	59.21	0.74	0.72	0.71	61.64	0.48	0.43	0.40
	57.66	0.74	0.72	0.70	59.66	0.47	0.42	0.39
	59.54	0.74	0.72	0.71	61.36	0.48	0.43	0.39
	64.57	0.75	0.74	0.72	65.93	0.50	0.45	0.42
	72.07	0.77	0.76	0.74	72.98	0.54	0.49	0.46
	81.31	0.79	0.78	0.77	81.80	0.59	0.55	0.51
	88.95	0.82	0.81	0.80	88.95	0.64	0.60	0.57
	Daily mean reduction (%)	1.80	2.36	2.90		8.75	10.51	12.03
No EC/ no dust	80.34	0.83	0.83	0.83	82.61	0.77	0.77	0.77
	71.25	0.81	0.81	0.81	73.66	0.75	0.75	0.75
	63.96	0.80	0.80	0.80	66.45	0.73	0.73	0.73
	59.21	0.79	0.79	0.79	61.64	0.72	0.72	0.72
	57.66	0.79	0.79	0.79	59.66	0.71	0.71	0.71
	59.54	0.79	0.79	0.79	61.36	0.72	0.72	0.72
	64.57	0.79	0.80	0.80	65.93	0.73	0.73	0.73
	72.07	0.81	0.81	0.81	72.98	0.75	0.75	0.75
	81.31	0.83	0.83	0.83	81.80	0.77	0.77	0.77

55	88.95	0.85	0.85	0.85	88.95	0.79	0.79	0.79

56   **Table S4. Radiative forcing for selected solar zenith angles (Wm⁻²) and daily mean (%)**

Pollutant	Site S6 (low EC and dust)				Site S1 (high EC and dust)			
	SZA	MAC 7.5	MAC 11	MAC 15	SZA	MAC 7.5	MAC 11	MAC 15
EC only	80.34	4.90	6.47	8.00	82.61	23.87	29.10	33.71
	71.25	5.86	7.72	9.53	73.66	27.50	33.25	38.21
	63.96	6.57	8.65	10.65	66.45	29.96	36.01	41.16
	59.21	7.00	9.70	11.33	61.64	31.39	37.58	42.82
	57.66	7.14	9.38	11.54	59.66	31.93	38.18	43.44
	59.54	6.97	9.16	11.29	61.36	31.47	37.67	42.91
	64.57	6.51	8.58	10.56	65.93	30.13	36.19	41.35
	72.07	5.78	7.61	9.40	72.98	27.75	33.53	38.52
	81.31	4.79	6.33	7.83	81.80	24.23	29.51	34.16
	88.95	3.93	5.20	6.44	88.95	20.88	25.62	29.86
	Daily mean	2.40	3.18	3.90	-	10.74	12.95	14.85
Dust only	80.34	0.20	0.20	0.20	82.61	1.43	1.43	1.43
	71.25	0.24	0.24	0.24	73.66	1.71	1.71	1.71
	63.96	0.27	0.27	0.27	66.45	1.91	1.91	1.91
	59.21	0.29	0.29	0.29	61.64	2.03	2.03	2.03
	57.66	0.30	0.30	0.30	59.66	2.08	2.08	2.08
	59.54	0.29	0.29	0.29	61.36	2.04	2.04	2.04
	64.57	0.27	0.27	0.27	65.93	1.93	1.93	1.93
	72.07	0.24	0.24	0.24	72.98	1.73	1.73	1.73

	81.31	0.20	0.20	0.20	81.80	1.38	1.38	1.38
	88.95	0.16	0.16	0.16	88.95	1.21	1.21	1.21
	Daily mean	0.10	0.10	0.10	-	0.67	0.67	0.67
	80.34	5.00	6.55	8.07	82.61	24.03	29.16	33.71
	71.25	5.98	7.82	9.61	73.66	27.70	33.33	38.22
	63.96	6.70	8.75	10.75	66.45	30.17	36.09	41.17
	59.21	7.14	9.32	11.43	61.64	31.61	37.68	42.83
EC and	57.66	7.28	9.49	11.64	59.66	32.15	38.27	43.45
dust	59.54	7.11	9.28	11.38	61.36	31.69	37.76	42.92
	64.57	6.65	8.68	10.66	65.93	30.34	36.27	41.36
	72.07	5.89	7.71	9.48	72.98	27.95	33.61	38.52
	81.31	4.89	6.41	7.90	81.80	24.39	29.58	34.16
	88.95	4.01	5.26	6.49	88.95	21.01	25.67	29.85
	Daily mean	2.45	3.20	3.93	-	10.81	12.98	14.86

57

58

59 **Table S5. Potential source countries for pollutants: a) identified using wind vector analysis; b) identified using trajectory analysis; and c) identified using the WRF-**
60 **STEM model**

Method	Region	Countries
a) Wind vector	East Asia	Thailand, Myanmar
	Central Asia	Tajikistan, Kyrgyzstan, Uzbekistan, Azerbaijan, Turkmenistan
	South Asia	Bay of Bengal, India, Pakistan, Afghanistan
	Middle East	Syria, Iraq, Lebanon, Armenia.
	West Asia	Iran, Turkey
b) Trajectory analysis	Central Asia	Uzbekistan, Western part Turkmenistan, Kazakhstan, Azerbaijan
	South Asia	Pakistan
	Middle East	Iraq, Armenia, Syria, United Arab Emirates, Jordan
	West Asia	Iran, Georgia, Turkey, Cyprus
	Russian Federation	Western Russia.
	Europe	Ukraine
	Africa	Egypt, Libya, Tunisia
c) WRF-STEM	East Asia	Indonesia, Singapore, Malaysia, Thailand, Myanmar, Sri Lanka
	Central Asia	Mongolia, Tajikistan, Kyrgyzstan, Kazakhstan, Uzbekistan, Turkmenistan
	South Asia	Bangladesh, India, Bhutan, Nepal
	Middle East	Oman, United Arab Emirates, Qatar, Bahrain, Kuwait, Iraq, Syria, Lebanon, Palestinian Territory, Israel, Jordan, Saudi Arabia, Yemen
	Russian	Russia

Federation	
West Asia	Azerbaijan, Armenia, Georgia, Turkey and Cyprus
Europe	Europe
Africa	Kenia, Somalia, Egypt, Congo
Other	China, Pakistan, Afghanistan, Iran