**Retrieving the global distribution of threshold of wind erosion from satellite data
and implementing it into the GFDL AM4.0/LM4.0 model**

Bing Pu^{1,2,*}, Paul Ginoux², Huan Guo^{2,3}, N. Christine Hsu⁴, John Kimball⁵, Beatrice Marticorena⁶, Sergey Malyshev², Vaishali Naik², Norman T. O'Neill⁷, Carlos Pérez García-Pando⁸, Joseph M. Prospero⁹, Elena Shevliakova², Ming Zhao²

¹Atmospheric and Oceanic Sciences Program, Princeton University,
Princeton, New Jersey 08544

²NOAA Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey 08540

³Cooperative Programs for the Advancement of Earth System Science, University
Corporation for Atmospheric Research, Boulder, Colorado, 80301

⁴NASA Goddard Space Flight Center, Greenbelt, Maryland, 20771

⁵Department of Ecosystem and Conservation Sciences, University of Montana,
Missoula, Montana 59812

⁶LISA, Universités Paris Est-Paris Diderot-Paris

⁷Département de géomatique appliquée, Université de Sherbrooke

⁸Barcelona Supercomputing Center, Barcelona, Spain, 08034

⁹Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, Miami,
Florida, 33149

* Current affiliation: Department of Geographical and Atmospheric Science, the
University of Kansas, Lawrence, Kansas, 66045

Supplement

Table S1 Name and location of 16 RSMAS sites used in Figure 8.

No.	Sites	Lag/Long
1	Cape Grim	40.68°S, 144.68°E
2	Marsh	62.18°S, 58.3°W
3	Mawson	67.6°S, 62.5°E
4	Palmer Station	64.77°S, 64.05°W
5	Norfolk Island	29.08°S, 167.98°E
6	American Samoa	14.25°S, 170.58°W
7	Midway Island	28.22°N, 177.35°W
8	Oahu Hawaii	21.33°N, 157.7°W
9	Cheju\Korea\Pem	33.52°N, 126.48°E
10	Fanning Island	3.92°N, 159.33°W
11	Ragged Point	13.17°N, 59.43°W
12	Izaña Tenerife	28.3°N, 16.5°W
13	Bermuda West & East	32.27°N, 64.87°W
14	Heimaey Iceland	63.4°N, 20.3°W
15	Mace Head	53.32°N, 9.85°W
16	RSMAS	25.75°N, 80.25°W

Table S2 Annual mean dust emission (Tg yr^{-1}) averaged over nine regions and global land from the Control, $V_{\text{thresh}}\text{Ann}$, and $V_{\text{thresh}}12\text{mn}$ simulations.

Regions	Control	$V_{\text{thresh}}\text{Ann}$	$V_{\text{thresh}}12\text{mn}$
Sahel	148	1185	970
Sahara	216	487	488
Arabian Peninsula	130	536	482
Northern China	74	78	73
India	55	438	239
US	4	78	37
South Africa	8	12	7
South America	25	169	103
Australia	44	556	434
Global land	978	5190	3562

One grid point in the U.S. (34.5N, 105W)

(a) Cumulative Frequency of DOD

(b) Cumulative Frequency of V_{10m}

Figure S1. A schematic of deriving $V_{threshold}$ at one grid point in the U.S. in March. (a) Determining the threshold percentage of cumulative frequency of DOD at $DOD_{thresh} = 0.02$ level; (b) matching the threshold percentage in (a) with the percentage of cumulative frequency distribution of surface wind speed to get the threshold of wind erosion.

Dust source function

S

S'

Figure S2. Dust source function S in Eq. 3 used in the Control simulation (top) and unmasked source function S' used in the $V_{\text{thresh}}12\text{mn}$ and $V_{\text{thresh}}\text{Ann}$ simulations (bottom).

Figure S3. (a)-(d) Model simulated LAI from the V_{thresh} 12mn run, average from 2001 to 2015 for four seasons. (e)-(h) The difference between the V_{thresh} 12mn run and the Control run. Global land averages are listed at the bottom of each plot.

Figure S4. Same as Fig. S3, but for the $V_{\text{thresh}}^{\text{Ann}}$ simulation.

AERONET SDA sites (N. Africa)

Figure S5. Location of 12 AERONET SDA COD sites (filled circles with site numbers) and three LISA sites (black triangles with site names in purple). For SDA site, the filled color indicates the climatological values of COD during 2003-2015, and the names of the sites are listed on the right corner of the plot.

Figure S6. Daily anomalies (with reference to the 2000-2015 mean) of 850 hPa geopotential height (gpm; only values over land are plotted) and wind vectors (m s^{-1}) from Oct. 17 to Oct. 19, 2012 from the $V_{\text{thresh}}12\text{mn}$ simulation. Elevations higher than 850 hPa are masked in grey.

Figure S7. (a)-(e) The differences between the $V_{\text{threshold}}$ and default V_t (i.e., $V_t = 6 \text{ m s}^{-1}$ in Eq. 3) in the AM4.0/LM4.0. (f)-(j) $V_{\text{threshold}}$ over the area extend beyond the domain of the default dust source function (S). Unit: m s^{-1} .

Figure S8. The differences of dust emission ($\text{g m}^{-2} \text{yr}^{-1}$) between the $V_{\text{thresh}} 12\text{mn}$ and Control runs are decomposed into two parts: (a)-(e) within the domain of the default dust source function (S) and (f)-(j) over the extended area outlined by $V_{\text{threshold}}$. Black boxes denote nine dusty regions as listed in Table 1.