Supplement to: Long-term INP measurements from four stations across the globe

Jann Schrod¹, Erik S. Thomson², Daniel Weber^{1,3}, Jens Kossmann¹, Christopher Pöhlker⁴, Jorge Saturno^{4,5}, Florian Ditas^{4,6}, Paulo Artaxo⁷, Valérie Clouard^{8,9}, Jean-Marie Saurel⁸, Martin Ebert¹⁰, Joachim Curtius¹, and Heinz G. Bingemer¹

¹Institute for Atmospheric and Environmental Sciences, Goethe University Frankfurt, Frankfurt am Main, Germany

²Department of Chemistry and Molecular Biology, Atmospheric Science, University of Gothenburg, Gothenburg, Sweden

³now at: Federal Waterways Engineering and Research Institute, Karlsruhe, Germany

⁴Max Planck Institute for Chemistry, Biogeochemistry and Multiphase Chemistry Departments, Mainz, Germany

⁵now at: Physikalisch-Technische Bundesanstalt, Braunschweig, Germany

⁶now at: Hessisches Landesamt für Naturschutz, Umwelt und Geologie, Wiesbaden, Germany

⁷Physics Institute, University of São Paulo, São Paulo, Brazil

⁸Université de Paris, Institut de physique du globe de Paris, Paris, France

⁹now at: Géosciences Environnement Toulouse, Toulouse, France

¹⁰Institute for Applied Geosciences, Technical University of Darmstadt, Darmstadt, Germany

Correspondence: Jann Schrod (schrod@iau.uni-frankfurt.de)

Figure S1. INP concentrations with respect to ice supersaturation for the sites SB (a), TO (b), MQ (c) and AZ (d). Boxes represent the interquartile range of observations, whiskers represent the 5–95 % range, squares indicate the median and stars give the arithmetic mean.

Figure S2. Contour plots of the probability density distributions of the INP concentrations for the complete spectrum of measurement conditions for the sites SB (a), TO (b), MQ (c) and AZ (d).

Figure S3. Scatter plot of the PM₁₀ concentration (MadininAir) and the INP concentration at MQ at $RH_{water} = 101\%$ and $T = -30^\circ\text{C}$.